

Full wwPDB X-ray Structure Validation Report (i)

Apr 3, 2023 – 01:18 pm BST

PDB ID	:	8A1I
Title	:	Crystal structure of murine Armc8 isoform beta
Authors	:	van gen Hassend, P.M.; Schindelin, H.
Deposited on	:	2022-06-01
Resolution	:	2.69 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.32.2
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.32.2

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 2.69 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
	$(\# { m Entries})$	$(\# { m Entries}, { m resolution} { m range}({ m \AA}))$
R_{free}	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069(2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	А	401	6% 80%	12%	8%
1	Ι	401	5%	11%	10%
1	Р	401	3% 7 9%	12%	9%



2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 17596 atoms, of which 8912 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues			Atc	oms				ZeroOcc	AltConf	Trace
1	Λ	270	Total	С	Η	Ν	Ο	\mathbf{S}	Se	76	0	0
	A	370	5896	1844	2995	498	536	13	10	10	0	0
1	т	260	Total	С	Η	Ν	0	S	Se	75	0	0
1	1	300	5708	1788	2896	480	521	13	10	15	0	0
1	D	266	Total	С	Η	Ν	0	S	Se	75	0	0
	1	300	5836	1826	2965	491	531	13	10	15	0	0

• Molecule 1 is a protein called Isoform 2 of Armadillo repeat-containing protein 8.

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	379	ILE	-	insertion	UNP Q9DBR3-2
А	380	ILE	-	insertion	UNP Q9DBR3-2
Ι	379	ILE	-	insertion	UNP Q9DBR3-2
Ι	380	ILE	-	insertion	UNP Q9DBR3-2
Р	379	ILE	-	insertion	UNP Q9DBR3-2
Р	380	ILE	-	insertion	UNP Q9DBR3-2

• Molecule 2 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: C₆H₁₃NO₄S).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf			
2	А	1	Total 25	C 6	Н 13	N 1	0 4	S 1	0	0

• Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	1	Total Cl 1 1	0	0
3	Ι	2	Total Cl 2 2	0	0

• Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	Δ	1	Total C H	0	2	0
4	Л	T	14 3 8	3		0
4	Δ	1	Total C H	0	0	0
т	11	I	12 3 6	3	0	0
4	Δ	1	Total C H	0	0	0
т	11	I	12 3 6	3	0	0
4	т	1	Total C H	0	1	0
	1	T	12 3 6	3	1	0
4	т	1	Total C H	0	1	0
	1	T	13 3 7	3	1	0
4	Р	1	Total C H	0	0	0
т	I	I	11 3 5	3	0	0
1	р	1	Total C H	0	0	0
	L L		11 3 5	3	0	0



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	
5	Ι	1	Total 5	0 4	S 1	0	0

• Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	А	10	Total O 10 10	0	0
6	Ι	12	Total O 12 12	0	0



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	Р	16	Total O 16 16	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Isoform 2 of Armadillo repeat-containing protein 8









PRO PRO VAL LEU LEU ALA ALA GLN GLN CLY VAL THR SER THR



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	65.64Å 148.18Å 67.14Å	Deperitor
a, b, c, α , β , γ	90.00° 90.57° 90.00°	Depositor
$\mathbf{P}_{\text{oscolution}}(\hat{\mathbf{A}})$	19.97 - 2.69	Depositor
Resolution (A)	47.17 - 2.69	EDS
% Data completeness	73.9 (19.97-2.69)	Depositor
(in resolution range)	73.9(47.17-2.69)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$1.74 (at 2.69 \text{\AA})$	Xtriage
Refinement program	PHENIX 1.14_3260	Depositor
B B.	0.214 , 0.248	Depositor
It, It _{free}	0.222 , 0.253	DCC
R_{free} test set	1310 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	75.7	Xtriage
Anisotropy	0.022	Xtriage
Bulk solvent $k_{sol}(e/A^3), B_{sol}(A^2)$	0.37, 48.3	EDS
L-test for $twinning^2$	$< L >=0.48, < L^2>=0.31$	Xtriage
	0.002 for l,k,-h	
Estimated twinning fraction	0.046 for h,-k,-l	Xtriage
	0.027 for l,-k,h	
F_o, F_c correlation	0.93	EDS
Total number of atoms	17596	wwPDB-VP
Average B, all atoms $(Å^2)$	93.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.10% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CL, SO4, MES

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond	lengths	Bond	angles
	Unam	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.28	0/2939	0.46	0/3964
1	Ι	0.28	0/2848	0.45	0/3842
1	Р	0.28	0/2906	0.46	0/3918
All	All	0.28	0/8693	0.46	0/11724

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	2901	2995	2994	28	0
1	Ι	2812	2896	2891	25	0
1	Р	2871	2965	2963	24	1
2	А	12	13	13	0	0
3	А	1	0	0	0	0
3	Ι	2	0	0	0	0
4	А	18	20	24	0	0
4	Ι	12	13	16	1	0
4	Р	12	10	16	0	0
5	Ι	5	0	0	0	0



Mol	Chain Non-H H(mode		H(model)	H(added) Clashes		Symm-Clashes			
6	А	10	0	0	0	0			
6	Ι	12	0	0	0	0			
6	Р	16	0	0	0	0			
All	All	8684	8912	8917	76	1			

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (76) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom 1	Atom 2	Interatomic	Clash	
Atom-1	Atom-2	distance (Å)	overlap (Å)	
1:I:298:VAL:HG23	1:I:360:ALA:HB2	1.60	0.83	
1:A:248:LYS:HD2	1:A:252:MSE:HE3	1.68	0.75	
1:I:308:ILE:HD11	1:I:315:GLN:HA	1.72	0.71	
1:P:17:VAL:HG21	1:P:60:LEU:HD11	1.75	0.69	
1:I:154:MSE:HE1	1:I:192:ASN:HB2	1.76	0.68	
1:P:298:VAL:HG23	1:P:360:ALA:HB2	1.77	0.66	
1:I:250:ILE:CG2	1:I:287:MSE:HE1	2.26	0.66	
1:P:250:ILE:CG2	1:P:287:MSE:HE1	2.26	0.64	
1:P:54:LYS:O	1:P:58:ILE:HG23	2.00	0.62	
1:I:237:PRO:HG3	1:I:263:MSE:HE1	1.81	0.61	
1:A:176:LYS:HB3	1:A:180:HIS:CE1	2.37	0.60	
1:A:250:ILE:CG2	1:A:287:MSE:HE1	2.32	0.60	
1:P:282:PRO:O	1:P:285:VAL:HG22	2.02	0.59	
1:P:250:ILE:HG23	1:P:287:MSE:HE1	1.84	0.59	
1:A:101:LEU:HB3	1:A:106:ILE:HD13	1.86	0.58	
1:A:250:ILE:HG23	1:A:287:MSE:HE1	1.87	0.57	
1:I:116:SER:O	1:I:122:ILE:HD11	2.05	0.56	
1:I:277:VAL:HG23	1:I:278:LEU:HG	1.88	0.56	
1:P:28:LEU:HD21	1:P:43:MSE:HE1	1.87	0.56	
1:A:116:SER:O	1:A:122:ILE:HD11	2.06	0.55	
1:P:197:LEU:HD12	1:P:236:LEU:HD11	1.88	0.54	
1:P:27:ARG:NH2	1:P:42:ASP:OD2	2.41	0.54	
1:A:282:PRO:O	1:A:285:VAL:HG22	2.09	0.53	
1:A:257:ALA:O	1:A:261:THR:HG23	2.08	0.53	
1:P:116:SER:O	1:P:122:ILE:HD11	2.08	0.53	
1:A:239:ILE:HG22	1:A:243:MSE:HE2	1.89	0.53	
1:A:180:HIS:HB3	1:A:183:ILE:HD12	1.91	0.52	
1:P:285:VAL:HG21	1:P:321:THR:HG22	1.93	0.51	
1:P:277:VAL:HG23	1:P:278:LEU:HG	1.92	0.51	
1:P:28:LEU:CD2	1:P:43:MSE:HE1	2.41	0.50	



	lo ao pagom	Interatomic	Clash		
Atom-1	Atom-2	distance (Å)	overlap (Å)		
1:I:254:LEU:HB2	1:I:287:MSE:HE2	1.93	0.49		
1:I:334:TYR:HB2	1:I:335:PRO:HD3	1.94	0.49		
1:A:197:LEU:HD12	1:A:236:LEU:HD11	1.95	0.49		
1:A:101:LEU:O	1:A:106:ILE:HG23	2.12	0.49		
1:I:239:ILE:HG22	1:I:243:MSE:HE2	1.93	0.49		
1:I:250:ILE:HG22	1:I:287:MSE:HE1	1.95	0.49		
1:P:143:LEU:O	1:P:143:LEU:HD23	2.13	0.49		
1:I:209:LEU:HD13	1:I:243:MSE:HE1	1.93	0.48		
1:A:144:TYR:CE2	1:A:180:HIS:HB2	2.48	0.48		
1:I:154:MSE:HE1	1:I:192:ASN:CB	2.43	0.48		
1:P:328:LEU:HD21	1:P:364:LEU:HD23	1.95	0.48		
1:I:282:PRO:O	1:I:285:VAL:HG22	2.13	0.48		
1:A:140:GLU:OE1	1:A:176:LYS:HB2	2.14	0.48		
1:I:43:MSE:HE1	1:I:66:LEU:HD21	1.96	0.48		
1:A:227:VAL:HG21	1:A:268:ALA:HB1	1.95	0.48		
1:I:308:ILE:CD1	1:I:315:GLN:HA	2.43	0.47		
1:I:257:ALA:O	1:I:261:THR:HG23	2.15	0.47		
1:A:254:LEU:HB2	1:A:287:MSE:HE2	1.95	0.47		
1:A:240:PHE:HD1	1:A:243:MSE:CE	2.28	0.47		
1:I:107:ILE:HD12	1:I:142:LEU:HD13	1.98	0.46		
1:A:227:VAL:CG2	1:A:268:ALA:HB1	2.46	0.46		
1:P:107:ILE:HB	1:P:108:PRO:HD3	1.98	0.46		
1:A:365:TYR:HE2	1:A:383:LEU:HD21	1.81	0.45		
1:I:308:ILE:HG23	1:I:367:SER:HB3	1.99	0.45		
1:A:37:LEU:O	1:A:40:VAL:HG12	2.17	0.45		
1:A:119:LEU:HD23	1:P:35:LYS:HE2	1.99	0.44		
1:P:56:ASN:O	1:P:60:LEU:HD13	2.18	0.44		
1:I:214:VAL:HG22	4:I:504:GOL:H12	2.00	0.44		
1:A:17:VAL:HG22	1:A:56:ASN:OD1	2.17	0.43		
1:A:70:LEU:HD23	1:A:70:LEU:O	2.18	0.43		
1:P:70:LEU:HD23	1:P:70:LEU:O	2.19	0.42		
1:P:253:GLN:OE1	1:P:286:ARG:NH1	2.51	0.42		
1:A:81:THR:O	1:A:85:VAL:HG23	2.19	0.42		
1:P:220:PRO:O	1:P:224:MSE:HG2	2.18	0.42		
1:I:225:THR:O	1:I:229:VAL:HG23	2.19	0.42		
1:I:281:LEU:HB3	1:I:282:PRO:HD3	2.01	0.42		
1:A:277:VAL:HG23	1:A:278:LEU:HG	2.02	0.42		
1:P:182:THR:HG22	1:P:186:ASN:OD1	2.20	0.41		
1:A:150:ILE:HB	1:A:151:PRO:HD3	2.02	0.41		
1:I:273:ASP:OD1	1:I:274:SER:N	2.53	0.41		
1:P:55:ALA:O	1:P:59:VAL:HG13	2.20	0.41		



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)	
1:A:203:LYS:HE3	1:A:203:LYS:H	1.86	0.41	
1:A:143:LEU:O	1:A:143:LEU:HD23	2.21	0.41	
1:I:28:LEU:HD11	1:I:43:MSE:HE2	2.03	0.41	
1:I:133:PHE:CD1	1:I:174:CYS:SG	3.14	0.41	
1:P:241:VAL:HG13	1:P:279:LYS:HG2	2.02	0.40	

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-1 Atom-2		Clash overlap (Å)	
1:P:16:GLU:OE1	1:P:378:LYS:NZ[1_556]	2.13	0.07	

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	А	366/401~(91%)	364 (100%)	2(0%)	0	100	100
1	Ι	356/401~(89%)	352~(99%)	4 (1%)	0	100	100
1	Р	362/401~(90%)	359~(99%)	3 (1%)	0	100	100
All	All	1084/1203~(90%)	1075 (99%)	9 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric Outliers		Percentiles		
1	А	329/344~(96%)	317~(96%)	12 (4%)	35 64		
1	Ι	318/344 (92%)	310~(98%)	8 (2%)	47 76		
1	Р	325/344~(94%)	314~(97%)	11 (3%)	37 66		
All	All	972/1032 (94%)	941 (97%)	31 (3%)	39 68		

All (31) residues with a non-rotameric sidechain are listed below:

\mathbf{Mol}	Chain	\mathbf{Res}	Type
1	А	14	LEU
1	А	44	LYS
1	А	92	MSE
1	А	130	ARG
1	А	203	LYS
1	А	209	LEU
1	А	246	ARG
1	А	252	MSE
1	А	265	ARG
1	А	292	ARG
1	А	344	ILE
1	А	345	LYS
1	Ι	16	GLU
1	Ι	51	ASN
1	Ι	78	GLU
1	Ι	130	ARG
1	Ι	265	ARG
1	Ι	292	ARG
1	Ι	333	LYS
1	Ι	358	ARG
1	Р	17	VAL
1	Р	34	GLN
1	Р	60	LEU
1	Р	73	GLU
1	Р	130	ARG
1	Р	159	ARG
1	Р	219	ASN
1	Р	265	ARG
1	Р	292	ARG
1	Р	346	ARG
1	Р	347	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.



5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 12 ligands modelled in this entry, 3 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal Type Chain Bee		Tink	Bond lengths		Bond angles		les			
	туре	Unam	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	GOL	Р	502	-	$5,\!5,\!5$	0.86	0	$5,\!5,\!5$	0.99	0
4	GOL	А	504	-	5,5,5	0.98	0	$5,\!5,\!5$	0.91	0
2	MES	А	501	-	12,12,12	2.13	1 (8%)	14,16,16	1.59	3 (21%)
4	GOL	Ι	504	-	$5,\!5,\!5$	0.91	0	$5,\!5,\!5$	0.94	0
4	GOL	Р	501	-	5,5,5	0.86	0	$5,\!5,\!5$	0.97	0
4	GOL	Ι	505	-	5,5,5	0.99	0	$5,\!5,\!5$	0.96	0
5	SO4	Ι	503	-	4,4,4	0.14	0	$6,\!6,\!6$	0.09	0
4	GOL	А	503	-	5,5,5	0.97	0	$5,\!5,\!5$	0.93	0
4	GOL	А	505	-	$5,\!5,\!5$	0.93	0	$5,\!5,\!5$	0.83	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

4 GOL P 502 - - $0/4/4/4$ -	Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
	4	GOL	Р	502	-	-	0/4/4/4	-



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	А	504	-	-	2/4/4/4	-
2	MES	А	501	-	-	1/6/14/14	0/1/1/1
4	GOL	Ι	504	-	-	1/4/4/4	-
4	GOL	Р	501	-	-	0/4/4/4	-
4	GOL	Ι	505	-	-	2/4/4/4	-
4	GOL	А	503	-	-	0/4/4/4	-
4	GOL	А	505	-	-	2/4/4/4	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	А	501	MES	C8-S	-7.09	1.67	1.77

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	А	501	MES	O1S-S-C8	2.95	110.47	106.92
2	А	501	MES	C5-N4-C3	2.44	114.31	108.83
2	А	501	MES	O3S-S-C8	2.27	109.44	105.77

There are no chirality outliers.

Mol	Chain	Res	Type	Atoms
2	А	501	MES	N4-C7-C8-S
4	А	505	GOL	O1-C1-C2-C3
4	А	505	GOL	O1-C1-C2-O2
4	А	504	GOL	O1-C1-C2-C3
4	Ι	504	GOL	O1-C1-C2-O2
4	Ι	505	GOL	O2-C2-C3-O3
4	Ι	505	GOL	C1-C2-C3-O3
4	А	504	GOL	O2-C2-C3-O3

All (8) torsion outliers are listed below:

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	Ι	504	GOL	1	0



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	360/401~(89%)	0.56	26 (7%) 15 13	49, 80, 132, 171	0
1	Ι	350/401~(87%)	0.57	20 (5%) 23 22	47, 83, 136, 207	0
1	Р	356/401~(88%)	0.40	12 (3%) 45 45	49, 79, 129, 166	0
All	All	1066/1203~(88%)	0.51	58 (5%) 25 24	47, 80, 132, 207	0

All (58) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	Ι	351	LEU	9.2
1	Ι	349	HIS	8.2
1	Ι	334	TYR	7.5
1	Ι	348	ASP	5.8
1	Р	344	ILE	5.2
1	Р	347	LEU	5.1
1	А	334	TYR	4.8
1	А	344	ILE	4.7
1	Ι	114	LEU	4.3
1	А	347	LEU	4.0
1	А	190	VAL	4.0
1	Ι	74	THR	3.9
1	Ι	336	SER	3.7
1	Ι	324	LEU	3.6
1	Р	334	TYR	3.4
1	А	142	LEU	3.3
1	Р	13	VAL	3.3
1	А	351	LEU	3.2
1	Ι	335	PRO	3.2
1	Р	346	ARG	3.1
1	Р	342	THR	3.1
1	A	48	ILE	3.0
1	А	349	HIS	2.9



Mol	Chain	Res	Type	RSRZ
1	А	47	VAL	2.8
1	Ι	350	ASP	2.8
1	А	345	LYS	2.8
1	Р	26	ASP	2.8
1	Ι	159	ARG	2.8
1	Р	28	LEU	2.7
1	А	174	CYS	2.7
1	Р	32	ASP	2.7
1	Ι	107	ILE	2.6
1	Ι	37	LEU	2.6
1	А	183	ILE	2.6
1	Р	351	LEU	2.6
1	А	195	HIS	2.6
1	А	60	LEU	2.6
1	А	102	LEU	2.6
1	Р	14	LEU	2.6
1	А	107	ILE	2.5
1	Ι	383	LEU	2.5
1	А	15	SER	2.5
1	А	14	LEU	2.3
1	А	140	GLU	2.3
1	А	335	PRO	2.3
1	Ι	81	THR	2.3
1	Ι	51	ASN	2.3
1	А	196	LEU	2.2
1	Ι	71	GLN	2.2
1	Р	35	LYS	2.2
1	А	184	LEU	2.1
1	А	336	SER	2.1
1	А	294	LEU	2.1
1	А	90	LEU	2.1
1	Ι	28	LEU	2.1
1	Ι	298	VAL	2.0
1	Ι	48	ILE	2.0
1	А	106	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(A^2)$	Q<0.9
4	GOL	Ι	505	6/6	0.79	0.26	89,93,112,112	1
4	GOL	Ι	504	6/6	0.81	0.31	95,98,118,118	1
4	GOL	А	503	6/6	0.82	0.26	93,112,117,117	2
4	GOL	Р	501	6/6	0.82	0.24	106,108,130,130	0
2	MES	А	501	12/12	0.84	0.18	120,130,152,156	0
4	GOL	А	504	6/6	0.85	0.21	97,100,121,121	0
4	GOL	А	505	6/6	0.88	0.30	74,75,90,90	0
3	CL	А	502	1/1	0.89	0.09	89,89,89,89	0
3	CL	Ι	501	1/1	0.89	0.15	89,89,89,89	0
4	GOL	Р	502	6/6	0.91	0.18	84,85,102,103	0
5	SO4	Ι	503	5/5	0.93	0.08	120,121,122,122	0
3	CL	Ι	502	1/1	0.97	0.16	84,84,84,84	0

6.5 Other polymers (i)

There are no such residues in this entry.

