



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 16, 2023 – 05:32 PM EDT

PDB ID : 2A40
Title : Ternary complex of the WH2 domain of WAVE with Actin-DNAse I
Authors : Chereau, D.; Kerff, F.; Dominguez, R.
Deposited on : 2005-06-27
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	375	 3% 87% 10%
1	D	375	 2% 84% 13%
2	B	260	 94% 6%
2	E	260	 92% 7%
3	C	32	 3% 56% 9% 31%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	32	 3% 62% 6% 31%
4	G	2	 100%
4	H	2	 100%

2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 11357 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Actin, alpha skeletal muscle.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	362	2899	1843	481	552	23	0	15	0
1	D	362	2878	1829	477	549	23	0	12	0

- Molecule 2 is a protein called Deoxyribonuclease-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	260	2071	1312	345	406	8	0	5	0
2	E	260	2109	1340	351	410	8	0	12	0

- Molecule 3 is a protein called Wiskott-Aldrich syndrome protein family member 2.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	C	22	181	109	36	36	0	1	0
3	F	22	181	109	36	36	0	1	0

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	G	2	28	16	2	10	0	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	H	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 5 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Ca	0	0
			1	1		
5	B	1	Total	Ca	0	0
			1	1		
5	D	1	Total	Ca	0	0
			1	1		
5	E	1	Total	Ca	0	0
			1	1		

- Molecule 6 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: C₁₀H₁₆N₅O₁₃P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
6	D	1	Total	C	N	O	P	0	0
			31	10	5	13	3		

- Molecule 7 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total C O 6 3 3	0	0
7	B	1	Total C O 6 3 3	0	0

- Molecule 8 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	B	1	Total Mg 1 1	0	0
8	E	1	Total Mg 1 1	0	0

- Molecule 9 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	288	Total O 288 288	0	0
9	B	167	Total O 167 167	0	0
9	C	7	Total O 7 7	0	0
9	D	253	Total O 253 253	0	0
9	E	183	Total O 183 183	0	0


Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	F	4	Total	O	0	0
			4	4		



- Molecule 3: Wiskott-Aldrich syndrome protein family member 2



- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucofuranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucofuranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	118.72Å 41.60Å 153.05Å 90.00° 108.79° 90.00°	Depositor
Resolution (Å)	47.51 – 1.80 47.49 – 1.80	Depositor EDS
% Data completeness (in resolution range)	80.6 (47.51-1.80) 80.6 (47.49-1.80)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.48 (at 1.79Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.166 , 0.215 0.167 , 0.217	Depositor DCC
R_{free} test set	5386 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	22.2	Xtrriage
Anisotropy	0.327	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 43.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	11357	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 84.91 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.4638e-07. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CA, NAG, HIC, ATP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.63	0/2988	0.71	0/4043
1	D	0.57	0/2961	0.65	0/4007
2	B	0.65	2/2132 (0.1%)	0.73	1/2902 (0.0%)
2	E	0.60	0/2192	0.73	1/2981 (0.0%)
3	C	0.48	0/184	0.51	0/244
3	F	0.40	0/184	0.47	0/244
All	All	0.61	2/10641 (0.0%)	0.70	2/14421 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	122	SER	C-O	6.28	1.35	1.23
2	B	173	CYS	CB-SG	5.49	1.91	1.82

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	123	THR	CA-CB-CG2	-5.40	104.85	112.40
2	E	166	MET	CG-SD-CE	5.37	108.79	100.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2899	0	2912	30	0
1	D	2878	0	2886	35	0
2	B	2071	0	2014	8	0
2	E	2109	0	2074	15	0
3	C	181	0	182	2	0
3	F	181	0	182	1	0
4	G	28	0	25	0	0
4	H	28	0	25	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
5	D	1	0	0	0	0
5	E	1	0	0	0	0
6	A	31	0	12	0	0
6	D	31	0	12	0	0
7	A	6	0	8	0	0
7	B	6	0	8	0	0
8	B	1	0	0	0	0
8	E	1	0	0	0	0
9	A	288	0	0	8	0
9	B	167	0	0	2	1
9	C	7	0	0	0	0
9	D	253	0	0	4	0
9	E	183	0	0	4	0
9	F	4	0	0	0	0
All	All	11357	0	10340	90	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (90) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:12[B]:ASN:ND2	9:A:1461:HOH:O	1.62	1.30
2:E:121[B]:HIS:HD2	9:E:1433:HOH:O	1.19	1.21
1:D:299:MET:HE3	1:D:313[B]:MET:HG3	1.30	1.11
1:D:299:MET:CE	1:D:313[B]:MET:HG3	1.80	1.11
1:D:37:ARG:NH2	1:D:84:LYS:HE3	1.67	1.08
1:D:216:LEU:HD22	1:D:238:LYS:HE3	1.43	1.01
2:E:92[A]:LEU:HD11	2:E:119:SER:HB3	1.57	0.86
1:D:37:ARG:HH22	1:D:84:LYS:HE3	1.47	0.77
2:B:103:SER:OG	9:B:1502:HOH:O	1.96	0.76
2:B:226:VAL:CG2	2:B:260:THR:HG23	2.17	0.73

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:360:GLN:NE2	1:A:360:GLN:H	1.87	0.73
1:D:244:ASP:OD1	1:D:245:GLY:N	2.22	0.72
1:A:115[A]:ASN:ND2	9:A:1640:HOH:O	2.23	0.72
2:B:17[B]:SER:OG	9:B:1523:HOH:O	2.11	0.67
1:A:176:MET:HE2	1:A:284[B]:LYS:HD2	1.77	0.67
1:D:291:LYS:HE3	9:D:1622:HOH:O	1.94	0.67
1:D:216:LEU:HD22	1:D:238:LYS:CE	2.21	0.67
1:D:216:LEU:CD2	1:D:238:LYS:HE3	2.25	0.65
2:E:121[B]:HIS:ND1	9:E:1434:HOH:O	2.15	0.60
2:E:92[A]:LEU:HD11	2:E:119:SER:CB	2.31	0.59
1:D:299:MET:HE1	1:D:313[B]:MET:HG3	1.78	0.58
1:D:275:HIS:CD2	1:D:275:HIS:H	2.23	0.56
1:A:326:LYS:NZ	9:A:1571:HOH:O	2.39	0.54
1:D:116[B]:ARG:CZ	9:D:1499:HOH:O	2.55	0.54
1:A:360:GLN:H	1:A:360:GLN:HE21	1.56	0.53
2:B:226:VAL:HG22	2:B:260:THR:HG23	1.90	0.53
1:D:37:ARG:HH21	1:D:84:LYS:HE3	1.66	0.53
1:D:35:VAL:HG11	1:D:84:LYS:HD2	1.91	0.52
1:A:216:LEU:HD22	1:A:238[A]:LYS:HE3	1.91	0.52
1:A:106:THR:HB	1:A:137:GLN:HG3	1.91	0.52
3:C:453:GLU:HA	3:C:453:GLU:OE2	2.10	0.52
1:D:272:ALA:HB1	1:D:276:GLU:HB2	1.92	0.51
1:A:115[B]:ASN:ND2	9:A:1440:HOH:O	2.43	0.51
1:A:176:MET:CE	1:A:284[B]:LYS:HD2	2.41	0.50
1:D:299:MET:HE3	1:D:313[B]:MET:CG	2.22	0.50
1:A:70:PRO:HG3	1:A:81:ASP:HB3	1.94	0.50
1:D:80:ASP:O	1:D:84:LYS:HE2	2.12	0.49
1:A:12[B]:ASN:CG	9:A:1461:HOH:O	2.26	0.49
2:E:260:THR:HG22	9:E:1344:HOH:O	2.12	0.49
1:D:37:ARG:NH2	1:D:84:LYS:CE	2.58	0.49
1:A:176:MET:HE1	1:A:284[B]:LYS:HE3	1.95	0.49
1:A:272:ALA:HB1	1:A:276[B]:GLU:HB2	1.94	0.48
1:D:172:PRO:HA	1:D:175:ILE:HD12	1.95	0.48
2:E:121[B]:HIS:CD2	9:E:1433:HOH:O	2.11	0.48
1:A:12[A]:ASN:HB3	1:A:71:ILE:CD1	2.43	0.48
1:A:37:ARG:NH2	1:A:81:ASP:OD1	2.36	0.48
3:C:437:SER:O	3:C:441[A]:SER:OG	2.30	0.48
2:E:24:TYR:HE1	2:E:27[B]:ARG:HH21	1.59	0.48
1:A:41:GLN:HB2	2:B:64:HIS:CD2	2.49	0.48
1:D:318:THR:HA	1:D:327:ILE:HD13	1.96	0.47
2:E:226:VAL:CG2	2:E:260:THR:HG23	2.45	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:326:LYS:HB3	1:D:326:LYS:HE3	1.69	0.47
1:D:149:THR:OG1	1:D:167:GLU:OE2	2.30	0.47
1:A:180[B]:LEU:HD13	1:A:267:ILE:HD11	1.97	0.46
1:D:275:HIS:H	1:D:275:HIS:HD2	1.63	0.46
1:A:69:TYR:HE1	9:A:1564:HOH:O	1.97	0.46
1:D:99:GLU:H	1:D:99:GLU:CD	2.19	0.46
2:E:106:ASN:HA	2:E:109:PHE:O	2.15	0.46
1:A:360:GLN:NE2	1:A:360:GLN:N	2.60	0.46
1:D:335:ARG:HA	1:D:338:SER:OG	2.17	0.45
1:D:105:LEU:HD11	1:D:123:MET:HG3	1.98	0.45
2:E:24:TYR:HE1	2:E:27[B]:ARG:NH2	2.14	0.45
1:D:336:LYS:HG2	9:D:1545:HOH:O	2.16	0.44
2:E:230:ALA:HA	2:E:256:GLU:O	2.17	0.44
1:A:318:THR:HA	1:A:327:ILE:CD1	2.47	0.44
1:D:208:ILE:HD13	1:D:242:LEU:HD22	1.99	0.44
1:A:176:MET:HE1	1:A:284[B]:LYS:CE	2.47	0.44
1:A:336:LYS:HE2	9:A:1632:HOH:O	2.18	0.44
3:F:454:GLU:H	3:F:454:GLU:HG3	1.56	0.44
1:D:70:PRO:HG3	1:D:81:ASP:HB3	2.00	0.43
1:A:113:LYS:HB3	1:A:113:LYS:HE3	1.91	0.43
1:A:124:PHE:O	1:A:128[A]:ASN:HA	2.19	0.43
1:A:124:PHE:O	1:A:128[B]:ASN:HA	2.19	0.43
1:D:190:MET:HE1	9:D:1512:HOH:O	2.18	0.42
2:B:220:LEU:HD12	2:B:220:LEU:HA	1.92	0.42
1:D:178:LEU:HG	1:D:180[B]:LEU:HB3	2.02	0.42
1:A:335:ARG:HA	1:A:338:SER:OG	2.20	0.42
2:E:115:VAL:HG12	2:E:158:TRP:CZ3	2.55	0.42
1:D:131:ALA:HB1	1:D:356:TRP:HB3	2.01	0.41
1:D:190:MET:CE	1:D:206:ARG:NH2	2.84	0.41
2:B:92:LEU:HD11	2:B:119:SER:HB2	2.02	0.41
2:B:230:ALA:HA	2:B:256:GLU:O	2.19	0.41
1:A:269[A]:MET:HG2	1:A:271:SER:HB3	2.02	0.41
2:E:30[B]:ARG:HH22	2:E:57:GLN:HE21	1.69	0.41
2:E:216:VAL:HG23	2:E:222:GLN:HG2	2.03	0.41
1:A:208:ILE:HD11	1:A:243:PRO:HG2	2.03	0.40
1:A:336:LYS:HG2	9:A:1441:HOH:O	2.21	0.40
1:D:275:HIS:CD2	1:D:275:HIS:N	2.89	0.40
1:D:239:SER:HB3	1:D:247:VAL:CG1	2.52	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
9:B:1464:HOH:O	9:B:1550:HOH:O[2_556]	2.18	0.02

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	374/375 (100%)	369 (99%)	5 (1%)	0	100	100
1	D	371/375 (99%)	365 (98%)	6 (2%)	0	100	100
2	B	263/260 (101%)	255 (97%)	8 (3%)	0	100	100
2	E	270/260 (104%)	266 (98%)	4 (2%)	0	100	100
3	C	21/32 (66%)	21 (100%)	0	0	100	100
3	F	21/32 (66%)	21 (100%)	0	0	100	100
All	All	1320/1334 (99%)	1297 (98%)	23 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	319/317 (101%)	314 (98%)	5 (2%)	62	54
1	D	316/317 (100%)	310 (98%)	6 (2%)	57	46
2	B	234/229 (102%)	232 (99%)	2 (1%)	78	75
2	E	241/229 (105%)	238 (99%)	3 (1%)	71	65

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	C	20/29 (69%)	17 (85%)	3 (15%)	3	0
3	F	20/29 (69%)	19 (95%)	1 (5%)	24	10
All	All	1150/1150 (100%)	1130 (98%)	20 (2%)	60	51

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	132	MET
1	A	225	ASN
1	A	250	ILE
1	A	288	ASP
1	A	326	LYS
2	B	2	LYS
2	B	101	CYS
3	C	441[A]	SER
3	C	441[B]	SER
3	C	454	GLU
1	D	50	LYS
1	D	128	ASN
1	D	225	ASN
1	D	250	ILE
1	D	288	ASP
1	D	360	GLN
2	E	101	CYS
2	E	119	SER
2	E	156	GLN
3	F	453	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (18) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	92	ASN
1	A	225	ASN
1	A	280	ASN
1	A	360	GLN
2	B	61	ASN
2	B	155	GLN
2	B	161	ASN
2	B	236	GLN
2	B	243	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	92	ASN
1	D	128	ASN
1	D	275	HIS
1	D	280	ASN
2	E	57	GLN
2	E	61	ASN
2	E	74	ASN
2	E	155	GLN
2	E	161	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	HIC	A	73	1	8,11,12	1.14	2 (25%)	6,14,16	0.65	0
1	HIC	D	73	1	8,11,12	1.07	1 (12%)	6,14,16	0.70	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	HIC	A	73	1	-	0/5/6/8	0/1/1/1
1	HIC	D	73	1	-	0/5/6/8	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	73	HIC	CD2-NE2	-2.19	1.34	1.38
1	A	73	HIC	CD2-NE2	-2.10	1.35	1.38
1	A	73	HIC	CD2-CG	2.07	1.39	1.36

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates (i)

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	G	1	2,4	14,14,15	0.50	0	17,19,21	1.03	1 (5%)
4	NAG	G	2	4	14,14,15	0.58	0	17,19,21	1.18	2 (11%)
4	NAG	H	1	2,4	14,14,15	0.65	0	17,19,21	1.00	1 (5%)
4	NAG	H	2	4	14,14,15	0.45	0	17,19,21	1.26	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	G	1	2,4	-	2/6/23/26	0/1/1/1
4	NAG	G	2	4	-	1/6/23/26	0/1/1/1
4	NAG	H	1	2,4	-	0/6/23/26	0/1/1/1
4	NAG	H	2	4	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	G	2	NAG	C1-O5-C5	2.91	116.14	112.19
4	H	1	NAG	O5-C1-C2	-2.88	106.75	111.29
4	G	2	NAG	C4-C3-C2	-2.61	107.20	111.02
4	H	2	NAG	O5-C5-C6	2.37	110.92	107.20
4	H	2	NAG	C1-O5-C5	2.25	115.24	112.19
4	G	1	NAG	O5-C5-C6	2.01	110.36	107.20

There are no chirality outliers.


All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	H	2	NAG	O5-C5-C6-O6
4	H	2	NAG	C4-C5-C6-O6
4	G	1	NAG	C8-C7-N2-C2
4	G	1	NAG	O7-C7-N2-C2
4	G	2	NAG	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 6 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	GOL	A	1383	-	5,5,5	0.36	0	5,5,5	0.88	0
7	GOL	B	1384	-	5,5,5	0.45	0	5,5,5	0.61	0
6	ATP	A	1379	5	26,33,33	1.04	1 (3%)	31,52,52	1.39	4 (12%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	ATP	D	1380	5	26,33,33	1.04	0	31,52,52	1.43	3 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	GOL	A	1383	-	-	2/4/4/4	-
7	GOL	B	1384	-	-	4/4/4/4	-
6	ATP	A	1379	5	-	1/18/38/38	0/3/3/3
6	ATP	D	1380	5	-	1/18/38/38	0/3/3/3

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	1379	ATP	C5-C4	2.23	1.46	1.40

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	D	1380	ATP	N3-C2-N1	-3.64	122.98	128.68
6	A	1379	ATP	O4'-C1'-C2'	-3.37	102.00	106.93
6	A	1379	ATP	O3G-PG-O2G	2.81	118.38	107.64
6	A	1379	ATP	N3-C2-N1	-2.79	124.32	128.68
6	D	1380	ATP	O3G-PG-O2G	2.77	118.24	107.64
6	A	1379	ATP	N6-C6-N1	2.11	122.95	118.57
6	D	1380	ATP	O4'-C1'-C2'	-2.04	103.94	106.93

There are no chirality outliers.


All (8) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
7	A	1383	GOL	C1-C2-C3-O3
7	B	1384	GOL	O1-C1-C2-C3
7	B	1384	GOL	C1-C2-C3-O3
7	B	1384	GOL	O2-C2-C3-O3
7	A	1383	GOL	O2-C2-C3-O3
6	A	1379	ATP	PB-O3B-PG-O1G
7	B	1384	GOL	O1-C1-C2-O2
6	D	1380	ATP	PB-O3B-PG-O1G

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	361/375 (96%)	-0.20	10 (2%) 53 47	19, 26, 40, 61	0
1	D	361/375 (96%)	-0.14	7 (1%) 66 63	21, 30, 44, 61	0
2	B	260/260 (100%)	-0.29	0 100 100	23, 28, 41, 55	0
2	E	260/260 (100%)	-0.43	0 100 100	22, 27, 37, 50	0
3	C	22/32 (68%)	0.41	1 (4%) 33 27	34, 40, 55, 59	0
3	F	22/32 (68%)	0.76	1 (4%) 33 27	33, 43, 53, 61	0
All	All	1286/1334 (96%)	-0.22	19 (1%) 73 70	19, 28, 43, 61	0

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	366	GLY	6.4
1	D	365	ALA	5.7
1	A	365	ALA	4.6
1	A	366	GLY	4.6
1	A	5	THR	4.3
1	D	360	GLN	3.8
1	D	245	GLY	3.3
3	C	433	SER	3.3
3	F	454	GLU	3.2
1	A	360	GLN	3.2
1	D	5	THR	2.7
1	A	247	VAL	2.6
1	A	245	GLY	2.5
1	A	364	GLU	2.2
1	D	246	GLN	2.2
1	A	363	ASP	2.1
1	A	244	ASP	2.1
1	A	246	GLN	2.1
1	D	324	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	HIC	A	73	11/12	0.96	0.09	23,30,34,35	0
1	HIC	D	73	11/12	0.96	0.09	26,34,38,39	0

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NAG	G	2	14/15	0.75	0.37	40,43,45,45	0
4	NAG	H	2	14/15	0.85	0.31	42,44,46,47	0
4	NAG	G	1	14/15	0.89	0.29	34,35,42,43	0
4	NAG	H	1	14/15	0.92	0.27	33,37,43,44	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain G:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
7	GOL	A	1383	6/6	0.67	0.21	52,58,59,60	0
7	GOL	B	1384	6/6	0.93	0.25	29,32,35,37	0
6	ATP	A	1379	31/31	0.98	0.05	18,21,25,25	0
6	ATP	D	1380	31/31	0.98	0.06	21,24,26,27	0
5	CA	A	1381	1/1	0.99	0.06	20,20,20,20	0
5	CA	B	1273	1/1	0.99	0.07	24,24,24,24	0
8	MG	B	1274	1/1	0.99	0.12	32,32,32,32	0
8	MG	E	1275	1/1	0.99	0.07	27,27,27,27	0
5	CA	D	1382	1/1	1.00	0.06	23,23,23,23	0
5	CA	E	1272	1/1	1.00	0.05	23,23,23,23	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around ATP A 1379:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around ATP D 1380:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.