



Full wwPDB X-ray Structure Validation Report ⓘ

May 16, 2020 – 03:12 pm BST

PDB ID : 3A43
Title : Crystal structure of HypA
Authors : Watanabe, S.; Arai, T.; Matsumi, R.; Aromi, H.; Imanaka, T.; Miki, K.
Deposited on : 2009-06-30
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	139	
1	B	139	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2110 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hydrogenase nickel incorporation protein hypA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	139	1083	691	182	204	6	0	0	0
1	B	118	900	572	153	169	6	0	0	0

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Zn	0	0
			1	1		
2	A	1	Total	Zn	0	0
			1	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	85	Total	O	0	0
			85	85		
3	B	40	Total	O	0	0
			40	40		

4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	83.93Å 83.93Å 365.01Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	41.97 – 2.30 41.97 – 2.30	Depositor EDS
% Data completeness (in resolution range)	94.6 (41.97-2.30) 94.7 (41.97-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.55 (at 2.32Å)	Xtrriage
Refinement program	CNS 1.2	Depositor
R, R_{free}	0.215 , 0.239 0.214 , 0.243	Depositor DCC
R_{free} test set	1104 reflections (5.17%)	wwPDB-VP
Wilson B-factor (Å ²)	47.3	Xtrriage
Anisotropy	0.666	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 66.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2110	wwPDB-VP
Average B, all atoms (Å ²)	80.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.64% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, FME

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.42	0/1093	0.63	0/1473
1	B	0.38	0/903	0.70	0/1214
All	All	0.40	0/1996	0.66	0/2687

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1083	0	1042	35	0
1	B	900	0	875	46	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	85	0	0	2	0
3	B	40	0	0	0	0
All	All	2110	0	1917	81	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 21.

All (81) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:72:LYS:HG2	1:B:79:GLU:HB3	1.36	1.05
1:B:81:LYS:O	1:B:83:LYS:HG2	1.82	0.80
1:A:80:TRP:HB2	1:A:84:GLU:HG3	1.65	0.78
1:A:93:ILE:O	1:A:96:ASP:HB2	1.83	0.77
1:B:76:CYS:SG	1:B:112:LYS:HB3	2.26	0.75
1:A:81:LYS:H	1:A:84:GLU:CG	1.99	0.74
1:B:36:LEU:HD21	1:B:69:ALA:HB2	1.69	0.73
1:B:70:VAL:HG23	1:B:121:VAL:HB	1.70	0.73
1:A:78:TYR:HE2	1:A:111:PRO:HG2	1.55	0.72
1:B:78:TYR:CG	1:B:112:LYS:HB2	2.25	0.71
1:B:72:LYS:HG2	1:B:79:GLU:CB	2.17	0.71
1:A:113:CYS:SG	1:A:114:GLY:N	2.64	0.70
1:A:81:LYS:O	1:A:84:GLU:HG2	1.92	0.70
1:A:74:ARG:HG2	1:A:117:ASP:O	1.93	0.69
1:B:71:PHE:O	1:B:79:GLU:HA	1.93	0.69
1:A:25:ARG:HE	1:A:27:LYS:HD3	1.57	0.68
1:A:81:LYS:H	1:A:84:GLU:HG2	1.58	0.68
1:B:136:GLU:HG2	1:B:136:GLU:O	1.94	0.66
1:B:36:LEU:CD2	1:B:69:ALA:HB2	2.25	0.66
1:B:120:VAL:HG11	1:B:124:ARG:NH2	2.11	0.65
1:A:83:LYS:O	1:A:94:LYS:HD2	1.96	0.65
1:B:78:TYR:CE2	1:B:111:PRO:HD2	2.31	0.65
1:A:70:VAL:HG22	1:A:81:LYS:HG2	1.82	0.61
1:B:113:CYS:SG	1:B:114:GLY:N	2.72	0.61
1:B:30:ARG:HB2	1:B:30:ARG:NH1	2.17	0.59
1:B:78:TYR:CD2	1:B:111:PRO:HD2	2.38	0.58
1:A:78:TYR:CE2	1:A:111:PRO:HG2	2.39	0.57
1:B:70:VAL:HG22	1:B:122:LYS:O	2.04	0.57
1:A:85:VAL:CB	1:A:88:LYS:HA	2.35	0.57
1:B:115:SER:OG	1:B:116:HIS:N	2.38	0.57
1:B:76:CYS:C	1:B:78:TYR:H	2.09	0.56
1:A:25:ARG:CZ	1:A:27:LYS:HZ3	2.19	0.56
1:B:120:VAL:HG11	1:B:124:ARG:CZ	2.38	0.54
1:A:26:VAL:HB	1:A:57:ALA:HA	1.90	0.52
1:A:85:VAL:CB	1:A:94:LYS:HD2	2.40	0.52
1:A:85:VAL:C	1:A:87:ASP:N	2.60	0.52
1:B:72:LYS:HA	1:B:78:TYR:O	2.10	0.52
1:B:30:ARG:CB	1:B:30:ARG:HH11	2.23	0.51
1:B:26:VAL:HB	1:B:57:ALA:HA	1.93	0.51
1:B:82:LEU:HD23	1:B:83:LYS:N	2.25	0.51
1:B:70:VAL:O	1:B:70:VAL:HG23	2.11	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:81:LYS:HG2	1:B:81:LYS:O	2.10	0.50
1:A:45:LYS:O	1:A:49:GLU:HG3	2.11	0.50
1:A:25:ARG:NE	1:A:27:LYS:HD3	2.25	0.49
1:A:81:LYS:N	1:A:84:GLU:HG2	2.25	0.49
1:B:76:CYS:C	1:B:78:TYR:N	2.66	0.48
1:A:25:ARG:CZ	1:A:27:LYS:NZ	2.76	0.48
1:A:81:LYS:O	1:A:84:GLU:CG	2.61	0.48
1:A:113:CYS:O	1:A:114:GLY:O	2.32	0.48
1:B:115:SER:HG	1:B:116:HIS:H	1.61	0.48
1:B:73:CYS:HA	1:B:118:PHE:HA	1.96	0.48
1:B:81:LYS:O	1:B:82:LEU:C	2.52	0.47
1:A:138:GLY:O	1:A:139:SER:HB2	2.14	0.47
1:B:11:VAL:HG13	1:B:52:PHE:CE2	2.50	0.47
1:A:109:ALA:HB1	1:A:115:SER:HB3	1.97	0.46
1:B:78:TYR:HE2	1:B:111:PRO:HD2	1.77	0.46
1:B:30:ARG:HH11	1:B:30:ARG:HB2	1.79	0.45
1:B:80:TRP:CD1	1:B:80:TRP:N	2.84	0.45
1:A:93:ILE:HG13	1:A:93:ILE:H	1.60	0.45
1:B:71:PHE:O	1:B:79:GLU:CA	2.63	0.45
1:B:110:CYS:HA	1:B:111:PRO:HD3	1.76	0.45
1:A:27:LYS:HE2	1:A:134:GLU:OE2	2.17	0.45
1:A:94:LYS:CB	1:A:94:LYS:NZ	2.80	0.44
1:B:36:LEU:HD21	1:B:69:ALA:CB	2.41	0.44
1:B:115:SER:OG	1:B:116:HIS:ND1	2.47	0.44
1:B:70:VAL:CG2	1:B:122:LYS:H	2.30	0.44
1:B:76:CYS:HB2	1:B:112:LYS:HG2	2.00	0.43
1:A:81:LYS:N	1:A:84:GLU:CG	2.76	0.43
1:B:27:LYS:HD3	1:B:134:GLU:OE1	2.18	0.43
1:A:85:VAL:O	1:A:86:LYS:C	2.56	0.43
1:B:76:CYS:O	1:B:78:TYR:N	2.51	0.43
1:A:50:GLN:NE2	3:A:205:HOH:O	2.46	0.42
1:B:120:VAL:O	1:B:120:VAL:HG13	2.19	0.42
1:B:45:LYS:O	1:B:49:GLU:HG3	2.19	0.42
1:B:78:TYR:CD1	1:B:112:LYS:HD2	2.54	0.42
1:A:30:ARG:NE	3:A:218:HOH:O	2.53	0.42
1:A:69:ALA:HA	1:A:123:GLY:HA3	2.00	0.41
1:A:73:CYS:SG	1:A:75:ASN:HB2	2.60	0.41
1:B:30:ARG:CB	1:B:30:ARG:NH1	2.83	0.41
1:B:76:CYS:SG	1:B:78:TYR:HB3	2.61	0.41
1:A:78:TYR:CE2	1:A:112:LYS:HG2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	137/139 (99%)	126 (92%)	9 (7%)	2 (2%)	10	10
1	B	114/139 (82%)	100 (88%)	9 (8%)	5 (4%)	2	1
All	All	251/278 (90%)	226 (90%)	18 (7%)	7 (3%)	5	3

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	83	LYS
1	B	111	PRO
1	A	114	GLY
1	B	77	ASN
1	B	117	ASP
1	A	89	PHE
1	B	123	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	107/113 (95%)	103 (96%)	4 (4%)	34	48
1	B	87/113 (77%)	81 (93%)	6 (7%)	15	20
All	All	194/226 (86%)	184 (95%)	10 (5%)	23	32

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	93	ILE
1	A	94	LYS
1	A	96	ASP
1	A	113	CYS
1	B	6	LEU
1	B	38	ASP
1	B	61	GLU
1	B	79	GLU
1	B	80	TRP
1	B	136	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	19	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	FME	A	1	1	8,9,10	0.68	0	7,9,11	1.45	1 (14%)
1	FME	B	1	1	8,9,10	0.61	0	7,9,11	1.32	1 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	FME	A	1	1	-	3/7/9/11	-
1	FME	B	1	1	-	0/7/9/11	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1	FME	C-CA-N	2.65	114.52	109.73
1	B	1	FME	C-CA-N	2.16	113.64	109.73

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	1	FME	CB-CA-N-CN
1	A	1	FME	O-C-CA-CB
1	A	1	FME	CB-CG-SD-CE

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	138/139 (99%)	1.30	41 (29%) 0 0	37, 58, 161, 180	0
1	B	117/139 (84%)	2.05	43 (36%) 0 0	38, 76, 161, 171	0
All	All	255/278 (91%)	1.64	84 (32%) 0 0	37, 65, 161, 180	0

All (84) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	111	PRO	11.3
1	B	71	PHE	10.4
1	B	118	PHE	9.4
1	B	75	ASN	7.8
1	A	112	LYS	7.6
1	B	109	ALA	7.4
1	B	138	GLY	7.3
1	A	91	GLU	7.1
1	B	80	TRP	6.9
1	B	82	LEU	6.6
1	B	70	VAL	6.6
1	A	86	LYS	6.3
1	B	113	CYS	5.8
1	B	112	LYS	5.7
1	B	85	VAL	5.6
1	B	115	SER	5.5
1	A	139	SER	5.2
1	B	139	SER	5.2
1	B	108	LEU	5.2
1	A	89	PHE	5.2
1	B	116	HIS	5.2
1	A	4	TRP	5.1
1	B	84	GLU	5.0
1	A	88	LYS	4.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	137	GLY	4.9
1	B	78	TYR	4.6
1	A	43	ILE	4.5
1	B	74	ARG	4.5
1	B	69	ALA	4.4
1	A	87	ASP	4.4
1	B	83	LYS	4.3
1	A	93	ILE	4.2
1	A	85	VAL	4.1
1	A	75	ASN	4.1
1	B	86	LYS	4.0
1	B	76	CYS	3.8
1	A	90	ASP	3.7
1	B	72	LYS	3.6
1	A	84	GLU	3.6
1	B	117	ASP	3.6
1	B	110	CYS	3.5
1	A	94	LYS	3.5
1	B	121	VAL	3.5
1	A	40	ALA	3.4
1	B	73	CYS	3.4
1	B	79	GLU	3.4
1	A	39	VAL	3.3
1	A	2	HIS	3.2
1	B	8	ASP	3.2
1	B	36	LEU	3.1
1	B	5	ALA	3.1
1	B	7	ALA	3.0
1	B	9	ALA	3.0
1	B	119	GLU	3.0
1	B	4	TRP	3.0
1	A	7	ALA	2.8
1	A	92	ARG	2.8
1	A	80	TRP	2.8
1	B	11	VAL	2.8
1	A	44	VAL	2.7
1	B	120	VAL	2.7
1	A	6	LEU	2.7
1	B	6	LEU	2.7
1	A	5	ALA	2.7
1	A	47	ALA	2.7
1	A	117	ASP	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	82	LEU	2.6
1	A	46	PHE	2.5
1	B	51	LEU	2.5
1	B	12	ARG	2.4
1	A	115	SER	2.4
1	A	38	ASP	2.3
1	B	129	ALA	2.3
1	A	111	PRO	2.3
1	A	116	HIS	2.3
1	A	113	CYS	2.2
1	A	74	ARG	2.2
1	A	137	GLY	2.2
1	A	101	PRO	2.2
1	A	3	GLU	2.1
1	A	100	ILE	2.1
1	A	77	ASN	2.1
1	A	78	TYR	2.1
1	A	114	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	FME	A	1	10/11	0.94	0.31	50,65,74,77	0
1	FME	B	1	10/11	0.95	0.27	56,58,61,63	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	ZN	B	702	1/1	0.47	0.07	170,170,170,170	0
2	ZN	A	701	1/1	0.93	0.04	118,118,118,118	0

6.5 Other polymers [i](#)

There are no such residues in this entry.