

Full wwPDB X-ray Structure Validation Report (i)

Nov 14, 2023 – 06:20 PM JST

PDB ID	:	6A4J
Title	:	Crystal structure of Thioredoxin reductase 2 from Staphylococcus aureus
Authors	:	Bose, M.; Bhattacharyya, S.; Ghosh, A.K.; Das, A.K.
Deposited on		
Resolution	:	3.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


Xtriage (Phenix) EDS buster-report Percentile statistics	: : :	20191225.v01 (using entries in the PDB archive December 25th 2019)
-	:	
CCP4 Ideal geometry (proteins)		7.0.044 (Gargrove) Engh & Huber (2001)
Ideal geometry (DNA, RNA) Validation Pipeline (wwPDB-VP)		

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 3.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \textbf{Whole archive} \\ (\#\textbf{Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R_{free}	130704	1026 (3.48-3.32)
Clashscore	141614	1055(3.48-3.32)
Ramachandran outliers	138981	1038 (3.48-3.32)
Sidechain outliers	138945	1038 (3.48-3.32)
RSRZ outliers	127900	2173 (3.50-3.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	А	344	82%	16%	·
1	В	344	5%	23%	•••



2 Entry composition (i)


There are 2 unique types of molecules in this entry. The entry contains 5243 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Ferredoxin–NADP reductase.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
1	Δ	340	Total	С	Ν	0	S	0	0	0
	A	340	2566	1626	440	490	10	0	0	0
1	р	338	Total	С	Ν	0	S	0	0	0
	D	000	2571	1629	448	484	10	0	0	0

• Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf		
2	٨	1	Total	С	Ν	Ο	Р	0	0
		1	53	27	9	15	2	0	0
0	D	1	Total	С	Ν	Ο	Р	0	0
	2 B	1	53	27	9	15	2	0	U



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Ferredoxin–NADP reductase



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	78.61Å 90.93Å 141.46Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.86 - 3.40	Depositor
Resolution (A)	19.86 - 3.40	EDS
% Data completeness	99.9 (19.86-3.40)	Depositor
(in resolution range)	99.9 (19.86-3.40)	EDS
R _{merge}	0.12	Depositor
R _{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$2.07 (at 3.44 \text{\AA})$	Xtriage
Refinement program	PHENIX (1.11.1_2575: ???)	Depositor
D D.	0.241 , 0.267	Depositor
R, R_{free}	0.241 , 0.267	DCC
R_{free} test set	733 reflections (5.08%)	wwPDB-VP
Wilson B-factor $(Å^2)$	64.7	Xtriage
Anisotropy	0.454	Xtriage
Bulk solvent $k_{sol}(e/A^3)$, $B_{sol}(A^2)$	0.38, 36.3	EDS
L-test for twinning ²	$ \langle L \rangle = 0.45, \langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.87	EDS
Total number of atoms	5243	wwPDB-VP
Average B, all atoms $(Å^2)$	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.09% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain		lengths	Bond angles		
		RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	0.27	0/2617	0.53	0/3557	
1	В	0.28	0/2620	0.57	0/3553	
All	All	0.27	0/5237	0.55	0/7110	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	2566	0	2466	36	0
1	В	2571	0	2508	68	0
2	А	53	0	31	7	0
2	В	53	0	31	10	0
All	All	5243	0	5036	99	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (99) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:206:ILE:HD11	1:B:209:LEU:HD21	1.54	0.90
1:B:179:ARG:NE	1:B:205:ARG:HB2	1.90	0.86
1:B:179:ARG:HE	1:B:205:ARG:HB2	1.47	0.80
1:B:179:ARG:HD3	1:B:226:VAL:HG22	1.66	0.76
1:B:247:ARG:HH11	1:B:266:ARG:HD2	1.52	0.74
1:B:179:ARG:HH21	1:B:205:ARG:HD2	1.58	0.69
1:A:50:ASP:OD1	2:A:401:FAD:HM71	1.95	0.67
1:A:91:GLU:OE1	1:A:313:TYR:OH	2.12	0.66
1:B:247:ARG:NH1	1:B:266:ARG:HD2	2.11	0.65
1:A:208:TYR:HB2	1:A:222:VAL:HB	1.79	0.65
1:B:136:LEU:HD11	1:B:241:ILE:HD11	1.79	0.63
1:B:179:ARG:HD3	1:B:226:VAL:CG2	2.30	0.60
1:A:130:ARG:NH1	1:A:138:TYR:OH	2.35	0.60
1:A:82:ARG:NH2	1:A:99:GLU:OE1	2.35	0.59
1:B:145:ARG:NH2	1:B:238:ASP:OD1	2.35	0.59
1:B:220:GLU:HG2	1:B:235:LYS:HG2	1.84	0.58
1:B:4:VAL:HG22	1:B:108:ALA:HB3	1.85	0.58
1:B:50:ASP:OD1	2:B:401:FAD:N3	2.33	0.57
1:B:255:THR:O	1:B:256:SER:OG	2.17	0.57
1:A:87:ARG:HG2	1:A:256:SER:OG	2.04	0.57
1:B:87:ARG:NH1	1:B:97:GLU:OE2	2.36	0.55
1:B:118:ILE:HD12	2:B:401:FAD:H2B	1.88	0.55
1:A:262:HIS:N	1:A:266:ARG:O	2.33	0.54
1:A:88:LYS:NZ	1:A:278:PRO:O	2.33	0.54
1:B:179:ARG:HH21	1:B:205:ARG:CD	2.21	0.54
2:B:401:FAD:H9	2:B:401:FAD:H2'	1.89	0.54
1:B:186:HIS:CD2	1:B:190:LYS:HE3	2.43	0.53
1:B:294:HIS:O	1:B:294:HIS:ND1	2.41	0.53
1:B:285:ASP:OD2	2:B:401:FAD:O2'	2.27	0.53
1:B:203:LYS:O	1:B:226:VAL:HG23	2.08	0.52
1:A:37:GLY:HA2	1:A:63:LEU:HD11	1.92	0.52
1:A:156:GLY:HA3	1:A:183:VAL:HG21	1.90	0.52
1:B:163:TRP:CZ3	1:B:242:SER:HB2	2.45	0.52
1:B:43:TYR:HE2	1:B:48:ILE:HD11	1.73	0.52
2:A:401:FAD:H6	1:B:326:SER:OG	2.09	0.52
1:A:39:LYS:HD2	2:A:401:FAD:H3'	1.90	0.52
1:A:225:HIS:ND1	1:A:226:VAL:O	2.42	0.52
1:B:137:HIS:HB2	1:B:240:ILE:HG12	1.92	0.51
1:B:147:LYS:HA	1:B:170:ILE:HG12	1.92	0.51
1:B:264:ASP:OD2	1:B:266:ARG:NH1	2.43	0.51
1:A:48:ILE:HG23	2:A:401:FAD:HM73	1.92	0.51
1:A:296:ILE:H	2:A:401:FAD:HM81	1.77	0.49



Continued from prev		Interatomic	Clash	
Atom-1	Atom-2	distance (Å)	overlap (Å)	
1:B:45:GLU:HB2	1:B:131:TYR:HA	1.93	0.49	
1:B:251:LEU:HD23	2:B:401:FAD:H62A	1.78	0.49	
1:A:209:LEU:HB3	1:A:218:ILE:HD12	1.96	0.48	
1:A:290:ASP:HB3	1:B:272:ASN:ND2	2.27	0.48	
1:B:39:LYS:HB2	2:B:401:FAD:O4'	2.13	0.48	
1:A:251:LEU:O	1:A:255:THR:HG23	2.13	0.48	
1:B:136:LEU:CD1	1:B:241:ILE:HD11	2.43	0.48	
1:B:251:LEU:HD23	2:B:401:FAD:N6A	2.29	0.48	
1:A:27:VAL:HB	1:A:75:PRO:HB3	1.94	0.48	
1:A:57:LYS:HE2	1:A:61:GLU:HB3	1.96	0.47	
1:B:3:ASP:N	1:B:26:SER:OG	2.38	0.47	
2:B:401:FAD:H2'	2:B:401:FAD:C9	2.43	0.47	
1:B:33:GLN:OE1	1:B:41:ARG:NH1	2.42	0.47	
1:A:339:VAL:HG11	1:B:132:GLN:HA	1.96	0.47	
1:B:42:ILE:HG12	1:B:43:TYR:HD1	1.79	0.46	
1:B:144:ARG:HD3	1:B:144:ARG:O	2.15	0.46	
1:A:154:SER:HA	1:A:177:VAL:HG23	1.97	0.46	
1:B:32:VAL:HG11	1:B:117:ILE:HG23	1.98	0.46	
1:B:275:THR:HG23	1:B:281:TYR:HA	1.98	0.46	
1:B:133:LEU:HD13	1:B:218:ILE:HD11	1.97	0.45	
1:A:135:ASN:CG	1:A:217:HIS:HB2	2.36	0.45	
2:A:401:FAD:H6	1:B:326:SER:CB	2.47	0.45	
1:B:61:GLU:OE2	1:B:64:LYS:NZ	2.48	0.44	
1:B:183:VAL:HG21	1:B:189:MET:HE2	2.00	0.44	
1:B:266:ARG:HH21	1:B:288:TYR:HD1	1.65	0.44	
1:B:158:ASN:OD1	1:B:185:GLY:HA3	2.18	0.43	
1:B:2:LYS:HD3	1:B:2:LYS:HA	1.81	0.43	
1:B:136:LEU:HD13	1:B:218:ILE:CD1	2.48	0.43	
1:B:116:GLY:O	2:B:401:FAD:H8A	2.18	0.43	
1:B:179:ARG:HH11	1:B:226:VAL:HG22	1.83	0.43	
1:B:42:ILE:CG1	1:B:43:TYR:HD1	2.32	0.43	
1:B:48:ILE:HG23	2:B:401:FAD:O4	2.19	0.43	
1:A:33:GLN:OE1	1:A:41:ARG:HD3	2.19	0.42	
1:B:161:LEU:HD11	1:B:183:VAL:HB	2.01	0.42	
1:A:339:VAL:CG1	1:B:132:GLN:HA	2.50	0.42	
1:B:51:ILE:HG21	1:B:54:ILE:HD12	2.01	0.42	
1:B:120:PRO:HA	1:B:243:HIS:NE2	2.35	0.42	
1:A:137:HIS:HB2	1:A:240:ILE:HG12	2.02	0.42	
1:A:325:SER:HB2	1:B:49:TRP:HB2	2.01	0.42	
1:A:339:VAL:HG12	1:B:132:GLN:HG3	2.01	0.42	
1:B:137:HIS:O	1:B:240:ILE:HA	2.19	0.42	

Continued from previous page...



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:226:VAL:O	1:A:228:SER:N	2.49	0.42
1:B:268:LYS:HE2	1:B:270:PHE:CE1	2.55	0.42
1:A:2:LYS:HD3	1:A:2:LYS:HA	1.85	0.41
1:A:35:GLU:OE1	1:A:41:ARG:NH2	2.53	0.41
1:A:323:TYR:CD1	1:B:141:GLN:HB3	2.54	0.41
1:B:292:LYS:HD2	1:B:292:LYS:HA	1.94	0.41
1:A:20:ALA:O	1:A:25:MET:HB2	2.21	0.41
1:A:5:THR:OG1	1:A:106:SER:HB3	2.21	0.41
1:A:300:PHE:CD2	1:B:52:GLY:HA3	2.56	0.41
1:B:27:VAL:HB	1:B:75:PRO:HB3	2.02	0.41
1:B:120:PRO:HA	1:B:243:HIS:CD2	2.56	0.41
1:A:35:GLU:HB2	1:A:41:ARG:CZ	2.51	0.40
2:A:401:FAD:C6	1:B:324:VAL:HB	2.51	0.40
1:A:328:HIS:ND1	1:A:330:VAL:HG22	2.36	0.40
1:B:51:ILE:HD12	1:B:62:ILE:HD13	2.02	0.40
1:B:266:ARG:NH2	1:B:288:TYR:HD1	2.18	0.40

Continued from previous page...

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	А	336/344~(98%)	324 (96%)	9~(3%)	3~(1%)	17	49
1	В	332/344~(96%)	317 (96%)	13 (4%)	2(1%)	25	57
All	All	668/688~(97%)	641 (96%)	22 (3%)	5(1%)	22	55

All (5) Ramachandran outliers are listed below:

Mol	Chain	\mathbf{Res}	Type
1	В	216	THR
1	А	184	SER



 $Continued \ from \ previous \ page...$

Mol	Chain	Res	Type
1	В	248	CYS
1	А	227	GLU
1	А	183	VAL

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	266/293~(91%)	265~(100%)	1 (0%)	91 95
1	В	269/293~(92%)	266~(99%)	3~(1%)	73 86
All	All	535/586~(91%)	531 (99%)	4 (1%)	84 92

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	1	MET
1	В	93	HIS
1	В	145	ARG
1	В	242	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Turne	e Chain Res Link		Link	Bond lengths			Bond angles		
	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z >2
2	FAD	В	401	-	$53,\!58,\!58$	0.47	0	68,89,89	0.65	2 (2%)
2	FAD	А	401	-	$53,\!58,\!58$	0.48	0	68,89,89	0.59	2 (2%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	В	401	-	-	11/30/50/50	0/6/6/6
2	FAD	А	401	-	-	7/30/50/50	0/6/6/6

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
2	А	401	FAD	P-O3P-PA	-3.02	122.47	132.83
2	В	401	FAD	P-O3P-PA	-2.66	123.68	132.83
2	А	401	FAD	C5A-C6A-N6A	2.26	123.78	120.35
2	В	401	FAD	C5A-C6A-N6A	2.24	123.75	120.35

There are no chirality outliers.

All (18) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
2	А	401	FAD	C5B-O5B-PA-O1A
2	А	401	FAD	C5B-O5B-PA-O2A
2	А	401	FAD	O4'-C4'-C5'-O5'
2	В	401	FAD	C2'-C1'-N10-C10
2	В	401	FAD	O4'-C4'-C5'-O5'
2	В	401	FAD	C5'-O5'-P-O2P
2	В	401	FAD	C5'-O5'-P-O3P
2	А	401	FAD	O4B-C4B-C5B-O5B
2	А	401	FAD	C3B-C4B-C5B-O5B
2	А	401	FAD	C3'-C4'-C5'-O5'
2	В	401	FAD	C3'-C4'-C5'-O5'
2	В	401	FAD	PA-O3P-P-O1P
2	В	401	FAD	C2'-C3'-C4'-C5'
2	В	401	FAD	PA-O3P-P-O5'
2	А	401	FAD	C5B-O5B-PA-O3P
2	В	401	FAD	O3'-C3'-C4'-C5'
2	В	401	FAD	N10-C1'-C2'-C3'
2	В	401	FAD	O4B-C4B-C5B-O5B


There are no ring outliers.

2 monomers are involved in 17 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	В	401	FAD	10	0
2	А	401	FAD	7	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and sufficient the outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle RSRZ \rangle$	# RSRZ >	2	$\mathbf{OWAB}(\mathbf{A}^2)$	Q<0.9
1	А	340/344~(98%)	0.05	16 (4%) 31	31	33, 49, 71, 85	0
1	В	338/344~(98%)	0.12	17 (5%) 28	29	32, 51, 72, 85	0
All	All	678/688~(98%)	0.09	33 (4%) 29	29	32, 50, 72, 85	0

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	А	228	SER	3.8
1	В	318	ALA	3.6
1	В	248	CYS	3.6
1	В	319	ASN	3.6
1	А	186	HIS	3.5
1	А	227	GLU	3.4
1	В	263	ASP	3.3
1	В	186	HIS	3.2
1	А	229	GLY	3.1
1	В	128	VAL	3.0
1	А	246	ASP	3.0
1	А	182	ASP	3.0
1	В	254	GLU	2.9
1	В	127	GLY	2.9
1	В	232	HIS	2.9
1	А	230	ASP	2.8
1	В	316	PRO	2.8
1	В	184	SER	2.8
1	А	316	PRO	2.7
1	В	219	SER	2.6
1	А	181	GLU	2.6
1	А	127	GLY	2.5
1	А	80	ASN	2.5
1	В	1	MET	2.4



Mol	Chain	Res	Type	RSRZ	
1	В	329	GLU	2.3	
1	А	263	ASP	2.3	
1	В	264	ASP	2.3	
1	А	249	ASN	2.3	
1	А	317	ASP	2.2	
1	В	317	ASP	2.2	
1	В	262	HIS	2.2	
1	А	318	ALA	2.1	
1	A	148	ASP	2.1	

Continued from previous page...

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.


6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mo	l Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathrm{\AA}^2)$	Q<0.9
2	FAD	A	401	53/53	0.90	0.24	$36,\!43,\!54,\!56$	0
2	FAD	В	401	53/53	0.93	0.22	37,42,57,59	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







6.5 Other polymers (i)

There are no such residues in this entry.

