



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 5, 2022 – 09:08 pm GMT

PDB ID : 7AB3
Title : Crystal structure of the Escherichia coli toxin-antitoxin system HipBST (HipT S57A)
Authors : Baerentsen, R.L.; Brodersen, D.E.
Deposited on : 2020-09-06
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.24
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0267
CCP4 : 7.1.010 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.24


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	107	
1	D	107	
2	B	103	
2	E	103	
3	C	341	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	341	 <p>3% 86% 11% ..</p>

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 8668 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Predicted transcriptional regulator, XRE family.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	70	Total	C	N	O	S	0	2	0
			566	364	98	103	1			
1	D	69	Total	C	N	O	S	0	1	0
			547	350	96	100	1			

- Molecule 2 is a protein called Couple_hipA domain-containing protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	100	Total	C	N	O	S	0	1	0
			814	524	148	141	1			
2	E	101	Total	C	N	O	S	0	0	0
			793	510	138	144	1			

- Molecule 3 is a protein called HipA_C domain-containing protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	P	S			
3	C	338	Total	C	N	O	P	S	0	0	0
			2739	1743	477	508	1	10			
3	F	337	Total	C	N	O	P	S	0	1	0
			2735	1741	475	508	1	10			

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	57	ALA	SER	engineered mutation	UNP B7UL96
C	336	HIS	-	expression tag	UNP B7UL96
C	337	HIS	-	expression tag	UNP B7UL96
C	338	HIS	-	expression tag	UNP B7UL96
C	339	HIS	-	expression tag	UNP B7UL96
C	340	HIS	-	expression tag	UNP B7UL96
C	341	HIS	-	expression tag	UNP B7UL96
F	57	ALA	SER	engineered mutation	UNP B7UL96

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	336	HIS	-	expression tag	UNP B7UL96
F	337	HIS	-	expression tag	UNP B7UL96
F	338	HIS	-	expression tag	UNP B7UL96
F	339	HIS	-	expression tag	UNP B7UL96
F	340	HIS	-	expression tag	UNP B7UL96
F	341	HIS	-	expression tag	UNP B7UL96


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	48	Total O 48 48	0	0
4	B	22	Total O 22 22	0	0
4	C	150	Total O 150 150	0	0
4	D	56	Total O 56 56	0	0
4	E	9	Total O 9 9	0	0
4	F	189	Total O 189 189	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Predicted transcriptional regulator, XRE family




- Molecule 1: Predicted transcriptional regulator, XRE family




- Molecule 2: Couple_hipA domain-containing protein




- Molecule 2: Couple_hipA domain-containing protein






- Molecule 3: HipA_C domain-containing protein





• Molecule 3: HipA_C domain-containing protein



4 Data and refinement statistics i

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	281.68Å 106.07Å 57.56Å 90.00° 90.65° 90.00°	Depositor
Resolution (Å)	49.75 – 2.40 49.75 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.9 (49.75-2.40) 99.9 (49.75-2.40)	Depositor EDS
R_{merge}	0.26	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.38 (at 2.39Å)	Xtrriage
Refinement program	BUSTER	Depositor
R, R_{free}	0.199 , 0.236 0.211 , 0.246	Depositor DCC
R_{free} test set	3364 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	69.9	Xtrriage
Anisotropy	0.297	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	(Not available) , (Not available)	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.017 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8668	wwPDB-VP
Average B, all atoms (Å ²)	88.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.37% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SEP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.42	0/577	0.62	0/777
1	D	0.43	0/556	0.58	0/748
2	B	0.37	0/838	0.53	0/1139
2	E	0.32	0/816	0.49	0/1113
3	C	0.43	0/2795	0.63	0/3783
3	F	0.45	0/2793	0.63	0/3780
All	All	0.42	0/8375	0.60	0/11340

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	566	0	598	3	0
1	D	547	0	582	2	0
2	B	814	0	791	12	0
2	E	793	0	745	7	0
3	C	2739	0	2663	26	0
3	F	2735	0	2664	23	0
4	A	48	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	22	0	0	0	0
4	C	150	0	0	1	0
4	D	56	0	0	0	0
4	E	9	0	0	0	0
4	F	189	0	0	1	0
All	All	8668	0	8043	64	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (64) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:F:277:GLN:HB2	3:F:338:HIS:HB2	1.53	0.90
3:F:301:ARG:HG3	3:F:301:ARG:HH11	1.38	0.89
2:B:91:ASN:HD21	3:C:66:GLN:HE22	1.25	0.84
3:C:70:ASP:O	3:C:71:GLU:HG3	1.87	0.75
3:C:52:GLN:HE21	3:C:52:GLN:H	1.38	0.69
1:A:42:ILE:HD13	1:A:48:LEU:HB2	1.75	0.68
3:F:318[A]:GLN:H	3:F:318[A]:GLN:CD	1.96	0.66
3:C:173:VAL:HG11	3:C:213:LEU:HD11	1.77	0.65
3:C:163:GLY:HA3	3:C:173:VAL:HG12	1.80	0.64
3:F:41:THR:CG2	3:F:44:GLN:HB2	2.33	0.59
3:F:188:ASP:OD2	3:F:193:LYS:HE3	2.03	0.59
2:B:5:VAL:HG11	2:B:86:ILE:HG12	1.86	0.58
3:F:41:THR:HG23	3:F:44:GLN:HB2	1.86	0.58
3:F:141:ASP:OD1	3:F:145:LEU:HB2	2.04	0.57
3:F:80:GLN:NE2	3:F:140:ARG:HH22	2.03	0.57
2:B:50:SER:OG	2:B:52:THR:O	2.23	0.57
2:B:91:ASN:HD21	3:C:66:GLN:NE2	2.01	0.56
2:E:8:LEU:HB3	2:E:13:VAL:HA	1.87	0.56
3:C:41:THR:HG22	3:C:44:GLN:HB2	1.87	0.56
2:B:9:LEU:HB3	2:B:14:VAL:HG21	1.89	0.54
3:C:188:ASP:O	3:C:189:ASN:O	2.26	0.53
3:C:198:ARG:NH2	3:C:303:GLU:OE1	2.41	0.53
2:B:4:ARG:HG3	2:B:18:SER:HB3	1.92	0.52
2:E:3:ARG:NH1	2:E:80:ASP:OD2	2.42	0.51
2:B:95:ALA:H	3:C:148:HIS:HD2	1.57	0.51
3:F:237:VAL:HG11	3:F:243:TYR:HB3	1.93	0.50
1:A:75:SER:O	1:A:79[A]:ARG:HG3	2.11	0.49
2:E:5:VAL:HG11	2:E:98:ILE:HD12	1.94	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:3:ARG:HD2	2:B:82:LEU:HG	1.96	0.48
3:F:301:ARG:HG3	3:F:301:ARG:NH1	2.18	0.48
2:B:91:ASN:ND2	3:C:66:GLN:HE22	2.03	0.47
1:D:80:LEU:HD12	1:D:87:VAL:HG11	1.97	0.47
2:B:37:ILE:O	3:C:148:HIS:HE1	1.98	0.46
3:F:245:TYR:HB2	4:F:571:HOH:O	2.16	0.46
1:A:80:LEU:HD12	1:A:87:VAL:HG11	1.97	0.46
3:C:304:ASN:O	3:C:308:MET:HG2	2.16	0.45
3:C:31:THR:HG21	3:C:33:LYS:HE2	1.98	0.45
3:C:70:ASP:C	3:C:71:GLU:HG3	2.37	0.45
3:F:127:GLU:H	3:F:127:GLU:HG3	1.57	0.44
3:F:304:ASN:O	3:F:308:MET:HG2	2.17	0.44
3:C:143:LYS:HG3	3:C:145:LEU:HD22	1.98	0.44
3:F:204:TRP:HB2	3:F:249:LEU:HD21	1.99	0.44
2:B:39:ILE:HD11	3:C:155:ALA:HB1	2.00	0.44
2:E:8:LEU:HD22	2:E:13:VAL:HG22	1.99	0.43
3:C:23:GLN:HB2	4:C:474:HOH:O	2.19	0.43
3:C:335:LEU:O	3:C:338:HIS:HD2	2.02	0.43
2:B:70:TYR:CE1	2:B:74:GLN:HG3	2.54	0.43
3:C:204:TRP:HB2	3:C:249:LEU:HD21	2.01	0.43
2:E:65:TRP:NE1	3:F:49:LEU:HD21	2.35	0.42
3:F:220:TYR:CE1	3:F:226:PRO:HB3	2.54	0.42
3:F:62:GLN:HA	3:F:63:PRO:HD3	1.94	0.42
3:F:224:LEU:HD23	3:F:224:LEU:H	1.84	0.42
3:F:212:HIS:CE1	3:F:214:ARG:HB2	2.54	0.42
3:C:212:HIS:CE1	3:C:214:ARG:HB2	2.55	0.42
1:D:46[A]:ARG:HG3	1:D:82:LYS:O	2.19	0.42
2:E:39:ILE:HD11	3:F:155:ALA:HB1	2.01	0.42
3:C:68:VAL:HG12	3:C:75:ARG:HB3	2.01	0.41
3:C:143:LYS:H	3:C:143:LYS:HG2	1.73	0.41
3:F:41:THR:HG22	3:F:44:GLN:HB2	2.02	0.41
2:E:63:GLU:OE2	3:F:59:SEP:O1P	2.38	0.41
3:C:138:TYR:OH	3:C:149:GLN:NE2	2.54	0.40
3:C:141:ASP:OD1	3:C:145:LEU:HB2	2.21	0.40
3:F:53:GLN:O	3:F:61:TYR:OH	2.34	0.40
3:C:62:GLN:HA	3:C:63:PRO:HD3	1.93	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	70/107 (65%)	68 (97%)	2 (3%)	0	100	100
1	D	68/107 (64%)	66 (97%)	2 (3%)	0	100	100
2	B	99/103 (96%)	92 (93%)	6 (6%)	1 (1%)	15	23
2	E	99/103 (96%)	94 (95%)	4 (4%)	1 (1%)	15	23
3	C	335/341 (98%)	322 (96%)	10 (3%)	3 (1%)	17	25
3	F	335/341 (98%)	323 (96%)	11 (3%)	1 (0%)	41	55
All	All	1006/1102 (91%)	965 (96%)	35 (4%)	6 (1%)	25	36

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	C	188	ASP
3	C	189	ASN
3	F	143	LYS
2	E	64	GLY
2	B	77	ASP
3	C	52	GLN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	63/93 (68%)	59 (94%)	4 (6%)	18	28
1	D	61/93 (66%)	57 (93%)	4 (7%)	16	26

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	85/89 (96%)	75 (88%)	10 (12%)	5	7
2	E	81/89 (91%)	71 (88%)	10 (12%)	4	6
3	C	290/293 (99%)	269 (93%)	21 (7%)	14	23
3	F	290/293 (99%)	273 (94%)	17 (6%)	19	32
All	All	870/950 (92%)	804 (92%)	66 (8%)	13	20

All (66) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	42	ILE
1	A	97	ASN
1	A	103	LEU
1	A	107	GLU
2	B	2	HIS
2	B	25	LEU
2	B	30[A]	HIS
2	B	30[B]	HIS
2	B	31	ASP
2	B	73	ILE
2	B	82	LEU
2	B	84	MET
2	B	88	ASN
2	B	91	ASN
3	C	13	GLU
3	C	14	ARG
3	C	16	GLU
3	C	17	GLN
3	C	23	GLN
3	C	26	LYS
3	C	27	ARG
3	C	52	GLN
3	C	54	LYS
3	C	56	MET
3	C	68	VAL
3	C	71	GLU
3	C	80	GLN
3	C	142	ASN
3	C	149	GLN
3	C	183	VAL
3	C	188	ASP
3	C	224	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	C	234	PHE
3	C	242	GLU
3	C	320	HIS
1	D	78	LYS
1	D	97	ASN
1	D	103	LEU
1	D	107	GLU
2	E	8	LEU
2	E	25	LEU
2	E	31	ASP
2	E	43	VAL
2	E	84	MET
2	E	85	LEU
2	E	86	ILE
2	E	98	ILE
2	E	101	TRP
2	E	102	GLU
3	F	13	GLU
3	F	23	GLN
3	F	26	LYS
3	F	27	ARG
3	F	49	LEU
3	F	51	ARG
3	F	52	GLN
3	F	53	GLN
3	F	56	MET
3	F	127	GLU
3	F	136	ARG
3	F	224	LEU
3	F	234	PHE
3	F	301	ARG
3	F	318[A]	GLN
3	F	318[B]	GLN
3	F	322	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	91	ASN
3	C	52	GLN
3	C	79	HIS
3	C	148	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	C	149	GLN
3	C	151	GLN
3	C	298	GLN
3	C	304	ASN
3	C	338	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SEP	C	59	3	8,9,10	0.69	0	8,12,14	2.72	4 (50%)
3	SEP	F	59	3	8,9,10	0.79	0	8,12,14	1.64	2 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SEP	C	59	3	-	0/5/8/10	-
3	SEP	F	59	3	-	4/5/8/10	-

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	59	SEP	OG-CB-CA	6.58	114.54	108.14

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	59	SEP	OG-CB-CA	3.09	111.15	108.14
3	F	59	SEP	OG-P-O1P	2.67	113.97	106.47
3	C	59	SEP	O2P-P-OG	2.30	112.84	106.73
3	C	59	SEP	O3P-P-OG	2.22	112.64	106.73
3	C	59	SEP	P-OG-CB	-2.17	112.33	118.30

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	F	59	SEP	N-CA-CB-OG
3	F	59	SEP	CB-OG-P-O1P
3	F	59	SEP	CB-OG-P-O2P
3	F	59	SEP	CB-OG-P-O3P

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	F	59	SEP	1	0

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	70/107 (65%)	0.14	2 (2%) 51 50	61, 70, 84, 100	0
1	D	69/107 (64%)	-0.12	0 100 100	59, 70, 82, 107	0
2	B	100/103 (97%)	0.69	11 (11%) 5 5	91, 113, 145, 149	0
2	E	101/103 (98%)	1.71	42 (41%) 0 0	112, 146, 211, 217	0
3	C	337/341 (98%)	0.17	9 (2%) 54 52	59, 79, 107, 127	0
3	F	336/341 (98%)	0.12	11 (3%) 46 45	54, 75, 109, 149	0
All	All	1013/1102 (91%)	0.34	75 (7%) 14 13	54, 80, 148, 217	0

All (75) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	101	TRP	5.5
3	F	143	LYS	5.3
2	E	77	ASP	5.1
2	E	79	ASN	4.9
2	E	64	GLY	4.5
3	C	54	LYS	4.5
2	E	2	HIS	4.4
2	B	75	HIS	4.3
2	E	99	LEU	4.3
2	E	73	ILE	4.2
2	E	75	HIS	4.1
3	F	55	GLY	4.1
2	E	5	VAL	4.0
2	E	44	ALA	4.0
2	E	53	LEU	3.8
2	E	54	HIS	3.7
2	E	82	LEU	3.7
3	F	224	LEU	3.7
2	B	81	LEU	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	E	78	GLU	3.6
2	E	70	TYR	3.6
3	C	52	GLN	3.5
2	E	22	SER	3.5
2	B	30[A]	HIS	3.5
2	B	79	ASN	3.5
2	E	12	GLN	3.2
2	E	28	TYR	3.2
2	E	100	PRO	3.0
2	E	74	GLN	3.0
2	E	17	LEU	3.0
3	F	33	LYS	3.0
2	E	98	ILE	2.9
1	A	107	GLU	2.8
2	B	77	ASP	2.8
2	E	3	ARG	2.8
3	C	339	HIS	2.8
3	C	55	GLY	2.7
2	E	51	GLU	2.7
2	E	30	HIS	2.6
2	E	7	VAL	2.5
2	E	26	PHE	2.5
3	F	52	GLN	2.5
3	C	61	TYR	2.5
3	F	245	TYR	2.5
2	E	80	ASP	2.5
3	C	13	GLU	2.5
2	E	21	ASP	2.4
3	F	53	GLN	2.4
2	E	81	LEU	2.4
2	E	4	ARG	2.4
2	B	22	SER	2.4
2	E	83	GLY	2.4
3	F	54	LYS	2.4
1	A	64[A]	TYR	2.4
3	C	224	LEU	2.4
3	C	143	LYS	2.3
2	E	14	VAL	2.3
2	B	6	LYS	2.3
3	F	142	ASN	2.3
2	E	101	TRP	2.3
2	E	90	LYS	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	84	MET	2.2
2	E	57	PHE	2.2
2	B	2	HIS	2.2
2	E	86	ILE	2.2
2	B	47	GLN	2.2
2	E	102	GLU	2.2
3	F	13	GLU	2.2
3	F	56	MET	2.1
2	E	43	VAL	2.1
2	E	76	ARG	2.1
2	E	52	THR	2.1
2	E	27	GLN	2.1
3	C	126	GLU	2.0
2	E	85	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	SEP	F	59	10/11	0.89	0.17	105,109,124,126	10
3	SEP	C	59	10/11	0.91	0.20	89,91,106,110	10

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.