

Full wwPDB X-ray Structure Validation Report (i)

Dec 3, 2020 – 09:04 AM GMT

PDB ID : 7ACC

> Title : human GTP cyclohydrolase I feedback regulatory protein (GFRP)

Authors Ebenhoch, R.; Nar, H.

2020-09-10 Deposited on

2.04 Å(reported) Resolution

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity 4.02b-467Xtriage (Phenix) 1.13

EDS 2.14.6buster-report 1.1.7(2018)

20191225.v01 (using entries in the PDB archive December 25th 2019) Percentile statistics

> Refmac 5.8.0158

CCP4 7.0.044 (Gargrove) Engh & Huber (2001)

Ideal geometry (proteins) Ideal geometry (DNA, RNA) Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) 2.14.6

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.04 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \text{Whole archive} \\ (\#\text{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar \; resolution} \\ (\#{\rm Entries, \; resolution \; range(\AA)}) \end{array}$
R_{free}	130704	1692 (2.04-2.04)
Clashscore	141614	1773 (2.04-2.04)
Ramachandran outliers	138981	1752 (2.04-2.04)
Sidechain outliers	138945	1752 (2.04-2.04)
RSRZ outliers	127900	1672 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	86	90%	7%	-
1	В	86	87%	9%	-
1	С	86	97%		•
1	D	86	92%	5%	-
1	Е	86	97%		



Continued from previous page...

Mol	Chain	Length	Quality of chain	
1	F	86	95%	
1	G	86	97%	•
1	Н	86	92%	7% •
1	I	86	90%	6% • •
1	J	86	90%	7% •



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 7408 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called GTP cyclohydrolase 1 feedback regulatory protein.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	A	83	Total	С	N	О	S	0	0	0
1	Λ	0.0	670	424	118	123	5	0	0	U
1	В	83	Total	С	N	О	S	0	0	0
1	D	00	670	424	118	123	5		U	U
1	$^{\circ}$ C	86	Total	С	N	Ο	S	0	0	0
1		00	694	438	123	127	6	0	U	U
1	D	83	Total	С	N	Ο	S	0	0	0
1	D	00	670	424	118	123	5		0	U
1	E	86	Total	С	N	Ο	S	0	0	0
1	П	00	694	438	123	127	6	0	O	U
1	F	83	Total	С	N	Ο	S	0	3	0
1	I	00	702	447	121	129	5	0	3	0
1	G	86	Total	С	N	Ο	S	0	0	0
1	G G	00	694	438	123	127	6	0	Ů	U
1	Н	85	Total	С	N	Ο	S	0	0	0
1	11	00	688	435	122	125	6	0	Ů	U
1	I	83	Total	С	N	О	S	0	0	0
1	1	00	670	424	118	123	5		U	U
1	J	83	Total	С	N	Ο	S	0	0	0
1	J	00	670	424	118	123	5			0

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	SER	_	expression tag	UNP P30047
A	0	HIS	-	expression tag	UNP P30047
В	-1	SER	_	expression tag	UNP P30047
В	0	HIS	-	expression tag	UNP P30047
С	-1	SER	_	expression tag	UNP P30047
С	0	HIS	_	expression tag	UNP P30047
D	-1	SER	-	expression tag	UNP P30047
D	0	HIS	_	expression tag	UNP P30047
Е	-1	SER	-	expression tag	UNP P30047



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
Е	0	HIS	-	expression tag	UNP P30047
F	-1	SER	-	expression tag	UNP P30047
F	0	HIS	-	expression tag	UNP P30047
G	-1	SER	-	expression tag	UNP P30047
G	0	HIS	-	expression tag	UNP P30047
Н	-1	SER	ı	expression tag	UNP P30047
Н	0	HIS	-	expression tag	UNP P30047
I	-1	SER	-	expression tag	UNP P30047
I	0	HIS	ı	expression tag	UNP P30047
J	-1	SER	_	expression tag	UNP P30047
J	0	HIS	_	expression tag	UNP P30047

• Molecule 2 is POTASSIUM ION (three-letter code: K) (formula: K) (labeled as "Ligand of Interest" by author).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	G	1	Total K 1 1	0	0
2	J	1	Total K 1 1	0	0
2	D	1	Total K 1 1	0	0
2	Е	1	Total K 1 1	0	0
2	Н	1	Total K 1 1	0	0
2	В	1	Total K 1 1	0	0
2	I	1	Total K 1 1	0	0
2	С	1	Total K 1 1	0	0
2	A	1	Total K 1 1	0	0
2	F	1	Total K 1 1	0	0

• Molecule 3 is water.

Mol	Chain	Residues	${f Atoms}$	ZeroOcc	AltConf
3	A	38	Total O 38 38	0	0



Continued from previous page...


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	В	52	Total O 52 52	0	0
3	С	65	Total O 65 65	0	0
3	D	56	Total O 56 56	0	0
3	E	69	Total O 69 69	0	0
3	F	67	Total O 67 67	0	0
3	G	53	Total O 53 53	0	0
3	Н	68	Total O 68 68	0	0
3	I	55	Total O 55 55	0	0
3	J	53	Total O 53 53	0	0



3 Residue-property plots (i)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: GTP cyclohydrolase 1 feedback regulatory protein



• Molecule 1: GTP cyclohydrolase 1 feedback regulatory protein







4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants	107.27Å 107.27Å 142.98Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	85.81 - 2.04	Depositor
Resolution (A)	85.81 - 2.04	EDS
% Data completeness	100.0 (85.81-2.04)	Depositor
(in resolution range)	100.0 (85.81-2.04)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	5.40 (at 2.05Å)	Xtriage
Refinement program	BUSTER 2.11.7	Depositor
D D.	0.167 , 0.191	Depositor
R, R_{free}	0.165 , 0.191	DCC
R_{free} test set	2743 reflections (5.14%)	wwPDB-VP
Wilson B-factor (Å ²)	26.6	Xtriage
Anisotropy	0.590	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.37, 55.9	EDS
L-test for twinning ²	$ < L > = 0.47, < L^2> = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7408	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 27.82 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.0542e-03.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section:

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Mol Chain		lengths	Bond angles	
MIOI	Chain	RMSZ	# Z >5	RMSZ	# Z > 5
1	A	0.37	0/684	0.56	0/923
1	В	0.43	0/684	0.58	0/923
1	С	0.44	0/709	0.57	0/957
1	D	0.43	0/684	0.57	0/923
1	E	0.42	0/709	0.58	0/957
1	F	0.45	0/718	0.60	0/969
1	G	0.43	0/709	0.58	0/957
1	Н	0.43	0/703	0.59	0/949
1	I	0.41	0/684	0.59	0/923
1	J	0.41	0/684	0.58	0/923
All	All	0.42	0/6968	0.58	0/9404

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	670	0	668	5	0
1	В	670	0	668	8	0
1	С	694	0	691	2	0



 $Continued\ from\ previous\ page...$

Mol	Chain	Non-H		H(added)	Clashes	Symm-Clashes
1	D	670	0	668	3	0
1	Ε	694	0	691	3	0
1	F	702	0	691	1	0
1	G	694	0	691	3	0
1	Н	688	0	686	4	0
1	I	670	0	668	4	0
1	J	670	0	668	5	0
2	A	1	0	0	0	0
2	В	1	0	0	0	0
2	С	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
2	G	1	0	0	0	0
2	Н	1	0	0	0	0
2	I	1	0	0	0	0
2	J	1	0	0	0	0
3	A	38	0	0	0	0
3	В	52	0	0	0	0
3	С	65	0	0	0	0
3	D	56	0	0	0	0
3	Ε	69	0	0	0	0
3	F	67	0	0	0	0
3	G	53	0	0	0	0
3	Н	68	0	0	0	0
3	I	55	0	0	0	0
3	J	53	0	0	1	0
All	All	7408	0	6790	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} \textbf{Interatomic} \\ \textbf{distance (Å)} \end{array}$	Clash overlap (Å)
1:H:2:PRO:HG3	1:I:61:GLU:HG2	1.82	0.61
1:B:10:ILE:HD11	1:H:10:ILE:HD11	1.84	0.59
1:A:14:VAL:HG21	1:H:41:GLY:HA3	1.86	0.57
1:D:10:ILE:HD11	1:F:10:ILE:HD11	1.87	0.56
1:A:82:HIS:ND1	1:A:84:GLU:OE1	2.29	0.56
1:A:82:HIS:NE2	1:B:61:GLU:OE1	2.39	0.54



Continued from previous page...

Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	${f distance}\;({ m \AA})$	overlap (Å)
1:C:27:GLU:OE1	1:C:63:ARG:NH1	2.34	0.53
1:A:10:ILE:HD11	1:I:10:ILE:HD11	1.90	0.53
1:J:22:GLU:HG3	1:J:45:TYR:HB2	1.90	0.52
1:E:10:ILE:HD11	1:J:10:ILE:HD11	1.91	0.51
1:J:61:GLU:HG2	3:J:201:HOH:O	2.11	0.51
1:E:41:GLY:HA3	1:I:14:VAL:HG21	1.92	0.50
1:A:54:ARG:HH22	1:E:1:MET:HG2	1.76	0.50
1:B:36:LYS:HZ2	1:B:45:TYR:HB3	1.77	0.49
1:D:14:VAL:HG21	1:J:41:GLY:HA3	1.97	0.47
1:B:36:LYS:NZ	1:B:45:TYR:CB	2.78	0.47
1:C:10:ILE:HD11	1:G:10:ILE:HD11	1.98	0.46
1:B:38:ARG:HD2	1:B:42:ASN:O	2.16	0.45
1:D:41:GLY:HA3	1:J:14:VAL:HG21	1.99	0.45
1:H:8:THR:HG21	1:H:53:PRO:HG3	2.00	0.43
1:I:22:GLU:HA	1:I:45:TYR:CE1	2.53	0.43
1:B:41:GLY:HA3	1:G:14:VAL:HG21	2.02	0.41
1:B:41:GLY:CA	1:G:14:VAL:HG21	2.51	0.41
1:B:36:LYS:HZ1	1:B:45:TYR:CB	2.34	0.41

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	A	81/86 (94%)	80 (99%)	1 (1%)	0	100	100
1	В	81/86 (94%)	80 (99%)	1 (1%)	0	100	100
1	С	84/86 (98%)	83 (99%)	1 (1%)	0	100	100
1	D	81/86 (94%)	80 (99%)	1 (1%)	0	100	100
1	Е	84/86 (98%)	83 (99%)	1 (1%)	0	100	100
1	F	84/86 (98%)	83 (99%)	1 (1%)	0	100	100



I'antimuad	trom	mromanne	naaa
Continued	110116	DICUIUU	Duuc

Mol	Chain	Analysed	Favoured	Allowed	Outliers Pero		\mathbf{ntiles}
1	G	84/86 (98%)	83 (99%)	1 (1%)	0	100	100
1	Н	83/86 (96%)	82 (99%)	1 (1%)	0	100	100
1	I	81/86 (94%)	81 (100%)	0	0	100	100
1	J	81/86 (94%)	81 (100%)	0	0	100	100
All	All	824/860 (96%)	816 (99%)	8 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percen	tiles
1	A	74/77~(96%)	73 (99%)	1 (1%)	67	65
1	В	74/77 (96%)	73 (99%)	1 (1%)	67	65
1	С	77/77 (100%)	77 (100%)	0	100	100
1	D	74/77~(96%)	73 (99%)	1 (1%)	67	65
1	E	77/77 (100%)	76 (99%)	1 (1%)	69	67
1	F	77/77 (100%)	77 (100%)	0	100	100
1	G	77/77~(100%)	76 (99%)	1 (1%)	69	67
1	Н	76/77~(99%)	75 (99%)	1 (1%)	69	67
1	I	74/77~(96%)	72 (97%)	2 (3%)	44	38
1	J	74/77~(96%)	74 (100%)	0	100	100
All	All	$754/770 \ (98\%)$	746 (99%)	8 (1%)	73	73

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	${f Res}$	\mathbf{Type}
1	A	61	GLU
1	В	84	GLU
1	D	61	GLU
1	Е	1	MET



Continued from previous page...

Mol	Chain	Res	Type
1	G	58	ASP
1	Н	61	GLU
1	I	38	ARG
1	I	61	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 10 ligands modelled in this entry, 10 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.



5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\#\mathrm{RSRZ}{>}2$	$\mathbf{OWAB}(\mathrm{\AA}^2)$	Q < 0.9
1	A	83/86 (96%)	-0.49	0 100 100	18, 39, 57, 62	0
1	В	83/86 (96%)	-0.56	1 (1%) 79 81	17, 31, 43, 52	0
1	С	86/86 (100%)	-0.67	0 100 100	17, 27, 43, 48	0
1	D	83/86 (96%)	-0.54	2 (2%) 59 63	18, 29, 45, 57	0
1	E	86/86 (100%)	-0.46	3 (3%) 44 48	19, 31, 48, 59	0
1	F	83/86 (96%)	-0.66	0 100 100	17, 26, 37, 46	0
1	G	86/86 (100%)	-0.52	0 100 100	17, 30, 46, 55	0
1	Н	85/86 (98%)	-0.53	2 (2%) 59 63	18, 29, 47, 62	0
1	I	83/86 (96%)	-0.46	1 (1%) 79 81	19, 34, 53, 59	0
1	J	83/86 (96%)	-0.60	0 100 100	18, 31, 49, 54	0
All	All	841/860 (97%)	-0.55	9 (1%) 80 82	17, 30, 51, 62	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	Н	0	HIS	5.1
1	I	84	GLU	4.3
1	Ε	1	MET	2.9
1	Ε	45	TYR	2.6
1	Н	1	MET	2.5
1	Ε	0	HIS	2.3
1	D	30	GLN	2.1
1	D	84	GLU	2.1
1	В	30	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



6.3 Carbohydrates (i)

There are no monosaccharides in this entry.


6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
2	K	Н	101	1/1	0.98	0.04	22,22,22,22	1
2	K	I	101	1/1	0.98	0.08	18,18,18,18	1
2	K	В	101	1/1	0.98	0.04	26,26,26,26	1
2	K	Е	101	1/1	0.98	0.07	21,21,21,21	1
2	K	С	101	1/1	0.99	0.07	20,20,20,20	1
2	K	A	101	1/1	0.99	0.04	24,24,24,24	1
2	K	D	101	1/1	0.99	0.05	15,15,15,15	1
2	K	G	101	1/1	0.99	0.07	16,16,16,16	1
2	K	F	101	1/1	0.99	0.06	15,15,15,15	1
2	K	J	101	1/1	0.99	0.07	15,15,15,15	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.











Electron density around K B 101: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)





Electron density around K E 101: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)





Electron density around K C 101:

 $2 \text{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\text{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)














Electron density around K D 101:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around K G 101:

 $2 \text{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\text{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around K F 101: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)





6.5 Other polymers (i)

There are no such residues in this entry.

