



Full wwPDB X-ray Structure Validation Report ⓘ

May 16, 2020 – 05:18 am BST

PDB ID : 2AE3
Title : Glutaryl 7-Aminocephalosporanic Acid Acylase: mutational study of activation mechanism
Authors : Kim, J.K.; Yang, I.S.; Shin, H.J.; Cho, K.J.; Ryu, E.K.; Kim, S.H.; Park, S.S.; Kim, K.H.
Deposited on : 2005-07-21
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	166	
2	B	528	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5587 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glutaryl 7-Aminocephalosporanic Acid Acylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	159	1243	790	218	234	1	0	0	0

- Molecule 2 is a protein called Glutaryl 7-Aminocephalosporanic Acid Acylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	522	4115	2605	722	777	11	0	0	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
3	A	1	6	3	3	0	0

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	68	Total 68	O 68	0	0
4	B	155	Total 155	O 155	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


Note EDS was not executed.


- Molecule 1: Glutaryl 7-Aminocephalosporanic Acid Acylase

Chain A:  84% 9%



- Molecule 2: Glutaryl 7-Aminocephalosporanic Acid Acylase

Chain B:  88% 10%



4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	73.39Å 73.39Å 379.68Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.22 – 2.40	Depositor
% Data completeness (in resolution range)	90.2 (48.22-2.40)	Depositor
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	CNS, REFMAC 5.1.24	Depositor
R, R_{free}	0.183 , 0.226	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	5587	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.64	0/1284	0.87	6/1759 (0.3%)
2	B	0.64	0/4231	0.83	12/5778 (0.2%)
All	All	0.64	0/5515	0.84	18/7537 (0.2%)

There are no bond length outliers.

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	214	ASP	CB-CG-OD2	8.94	126.35	118.30
1	A	143	ARG	NE-CZ-NH2	-8.82	115.89	120.30
1	A	143	ARG	NE-CZ-NH1	8.35	124.47	120.30
1	A	23	ASP	CB-CG-OD2	6.97	124.57	118.30
2	B	390	ASP	CB-CG-OD2	6.26	123.94	118.30
2	B	164	ASP	CB-CG-OD2	6.25	123.92	118.30
2	B	203	ASP	CB-CG-OD2	5.94	123.64	118.30
1	A	121	ASP	CB-CG-OD2	5.59	123.33	118.30
1	A	52	ASP	CB-CG-OD2	5.54	123.29	118.30
2	B	91	ASP	CB-CG-OD2	5.53	123.27	118.30
2	B	420	ASP	CB-CG-OD2	5.43	123.19	118.30
1	A	99	ASP	CB-CG-OD2	5.39	123.15	118.30
2	B	287	ASP	CB-CG-OD2	5.37	123.14	118.30
2	B	497	ASP	CB-CG-OD2	5.22	123.00	118.30
2	B	286	ASP	CB-CG-OD2	5.18	122.96	118.30
2	B	495	ARG	NE-CZ-NH1	5.17	122.89	120.30
2	B	384	ASP	CB-CG-OD2	5.10	122.89	118.30
2	B	133	ASP	CB-CG-OD2	5.09	122.88	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1243	0	1154	7	0
2	B	4115	0	3950	27	0
3	A	6	0	8	0	0
4	A	68	0	0	1	0
4	B	155	0	0	3	0
All	All	5587	0	5112	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:117:GLN:HG3	4:A:339:HOH:O	1.83	0.78
2:B:100:PRO:HD2	2:B:120:ILE:O	1.83	0.78
2:B:271:HIS:ND1	2:B:370:TRP:CZ2	2.62	0.68
2:B:229:LEU:O	2:B:231:ARG:HG3	1.97	0.65
2:B:245:ASP:HB2	2:B:271:HIS:HD2	1.60	0.65
2:B:245:ASP:HB2	2:B:271:HIS:CD2	2.36	0.59
1:A:92:TRP:CH2	1:A:142:HIS:ND1	2.72	0.55
2:B:453:HIS:HD2	2:B:454:GLY:N	2.10	0.50
2:B:100:PRO:CD	2:B:120:ILE:O	2.55	0.49
2:B:57:LYS:HE2	2:B:68:ASN:HD22	1.77	0.49
2:B:79:TYR:CG	2:B:125:HIS:HD2	2.31	0.48
1:A:161:ASP:HB3	1:A:162:PRO:HD2	1.94	0.48
2:B:453:HIS:CD2	2:B:454:GLY:N	2.81	0.48
2:B:49:THR:HB	2:B:56:ILE:HA	1.95	0.47
2:B:79:TYR:CG	2:B:125:HIS:CD2	3.04	0.45
2:B:199:ARG:HG2	2:B:207:TRP:CZ2	2.52	0.45
1:A:97:SER:HB3	4:B:673:HOH:O	2.16	0.45
1:A:52:ASP:H	2:B:480:GLN:HE22	1.67	0.43
2:B:445:GLU:H	2:B:445:GLU:CD	2.21	0.43
2:B:73:MET:SD	2:B:196:ALA:HB2	2.59	0.42
1:A:148:LEU:HD13	2:B:30:TYR:CE1	2.54	0.42
2:B:37:HIS:CD2	2:B:46:TYR:HD1	2.36	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:300:HIS:N	2:B:300:HIS:CD2	2.88	0.41
2:B:502:LEU:HD22	2:B:507:GLN:HB3	2.02	0.41
2:B:72:GLY:O	2:B:73:MET:C	2.58	0.41
2:B:257:THR:HB	2:B:258:PRO:HD2	2.02	0.41
2:B:100:PRO:CG	4:B:681:HOH:O	2.69	0.41
2:B:245:ASP:CB	2:B:271:HIS:HD2	2.32	0.41
2:B:40:THR:HB	2:B:41:PRO:CD	2.50	0.41
1:A:88:ARG:HD2	1:A:142:HIS:CE1	2.56	0.41
2:B:343:ARG:O	2:B:407:ASP:HA	2.20	0.40
2:B:453:HIS:CE1	4:B:604:HOH:O	2.74	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	157/166 (95%)	151 (96%)	3 (2%)	3 (2%)	8	10
2	B	520/528 (98%)	503 (97%)	15 (3%)	2 (0%)	34	48
All	All	677/694 (98%)	654 (97%)	18 (3%)	5 (1%)	22	32

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	164	ASP
1	A	162	PRO
1	A	163	PRO
2	B	177	PHE
2	B	100	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	122/128 (95%)	116 (95%)	6 (5%)	25	40
2	B	431/437 (99%)	422 (98%)	9 (2%)	53	72
All	All	553/565 (98%)	538 (97%)	15 (3%)	44	65

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	16	ARG
1	A	75	GLN
1	A	116	GLN
1	A	117	GLN
1	A	143	ARG
1	A	165	LEU
2	B	21	ASN
2	B	300	HIS
2	B	323	GLU
2	B	386	LYS
2	B	452	VAL
2	B	460	MET
2	B	468	ARG
2	B	475	TYR
2	B	521	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	50	HIS
1	A	53	ASN
1	A	83	ASN
2	B	50	GLN
2	B	68	ASN
2	B	189	ASN
2	B	453	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	480	GLN
2	B	485	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	A	301	-	5,5,5	0.30	0	5,5,5	0.26	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	301	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.