



# Full wwPDB X-ray Structure Validation Report ⓘ

May 21, 2020 – 02:58 am BST

PDB ID : 5AED  
Title : A bacterial protein structure in glycoside hydrolase family 31  
Authors : Jin, Y.; Speciale, G.; Davies, G.J.; Williams, S.J.; Goddard-Borger, E.D.  
Deposited on : 2015-08-28  
Resolution : 1.91 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.91 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	7937 (1.94-1.90)
Clashscore	141614	8644 (1.94-1.90)
Ramachandran outliers	138981	8530 (1.94-1.90)
Sidechain outliers	138945	8530 (1.94-1.90)
RSRZ outliers	127900	7793 (1.94-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	686	 3% 93% 5%
1	B	686	 2% 89% 8%

## 2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 11533 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ALPHA-GLUCOSIDASE YIHQ.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	674	Total	C	N	O	S	0	1	0
			5437	3483	911	1015	28			
1	B	671	Total	C	N	O	S	0	2	0
			5414	3469	906	1011	28			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	679	LEU	-	expression tag	UNP P32138
A	680	GLU	-	expression tag	UNP P32138
A	681	HIS	-	expression tag	UNP P32138
A	682	HIS	-	expression tag	UNP P32138
A	683	HIS	-	expression tag	UNP P32138
A	684	HIS	-	expression tag	UNP P32138
A	685	HIS	-	expression tag	UNP P32138
A	686	HIS	-	expression tag	UNP P32138
B	679	LEU	-	expression tag	UNP P32138
B	680	GLU	-	expression tag	UNP P32138
B	681	HIS	-	expression tag	UNP P32138
B	682	HIS	-	expression tag	UNP P32138
B	683	HIS	-	expression tag	UNP P32138
B	684	HIS	-	expression tag	UNP P32138
B	685	HIS	-	expression tag	UNP P32138
B	686	HIS	-	expression tag	UNP P32138

- Molecule 2 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula: C<sub>6</sub>H<sub>14</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 8 6 2	0	0
2	B	1	Total C O 8 6 2	0	0
2	B	1	Total C O 8 6 2	0	0

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	2	Total Ca 2 2	0	0
3	A	1	Total Ca 1 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	253	Total O 253 253	0	2
4	B	402	Total O 402 402	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: ALPHA-GLUCOSIDASE YIHQ



- Molecule 1: ALPHA-GLUCOSIDASE YIHQ



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	78.71Å 113.18Å 111.72Å 90.00° 109.23° 90.00°	Depositor
Resolution (Å)	105.49 – 1.91 47.81 – 1.91	Depositor EDS
% Data completeness (in resolution range)	96.6 (105.49-1.91) 96.7 (47.81-1.91)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.22 (at 1.91Å)	Xtrriage
Refinement program	REFMAC 5.8.0124	Depositor
R, $R_{free}$	0.168 , 0.201 0.182 , 0.211	Depositor DCC
$R_{free}$ test set	6763 reflections (4.89%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	27.2	Xtrriage
Anisotropy	0.143	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 54.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.019 for h,-k,-h-l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	11533	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	33.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.65% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, MPD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.84	2/5600 (0.0%)	0.87	9/7612 (0.1%)
1	B	1.10	6/5581 (0.1%)	1.02	28/7585 (0.4%)
All	All	0.98	8/11181 (0.1%)	0.95	37/15197 (0.2%)

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	472	ASP	CG-OD2	5.93	1.39	1.25
1	B	546	TRP	CE3-CZ3	5.57	1.48	1.38
1	B	538	GLU	C-O	5.57	1.33	1.23
1	B	126	GLU	CD-OE1	5.54	1.31	1.25
1	A	472	ASP	CG-OD2	5.42	1.37	1.25
1	A	472	ASP	CB-CG	5.35	1.62	1.51
1	B	472	ASP	CB-CG	5.17	1.62	1.51
1	B	193	TYR	CE1-CZ	-5.16	1.31	1.38

All (37) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	276	ARG	NE-CZ-NH2	-9.74	115.43	120.30
1	A	455	ARG	NE-CZ-NH2	-8.69	115.96	120.30
1	B	140	ASP	CB-CG-OD1	8.40	125.86	118.30
1	B	357	ASP	CB-CG-OD2	-8.19	110.93	118.30
1	B	611	ARG	NE-CZ-NH1	7.99	124.29	120.30
1	A	455	ARG	NE-CZ-NH1	7.26	123.93	120.30
1	B	663	ARG	NE-CZ-NH2	-6.68	116.96	120.30
1	B	663	ARG	NE-CZ-NH1	6.62	123.61	120.30
1	B	198	ASP	CB-CG-OD1	6.62	124.26	118.30
1	A	611	ARG	NE-CZ-NH1	6.53	123.57	120.30
1	B	476	ASP	CB-CG-OD1	6.30	123.97	118.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	611	ARG	NE-CZ-NH2	-6.27	117.17	120.30
1	B	472	ASP	CB-CG-OD1	-6.18	112.73	118.30
1	B	276	ARG	NE-CZ-NH1	6.17	123.39	120.30
1	B	357	ASP	CB-CG-OD1	6.17	123.85	118.30
1	B	198	ASP	CB-CG-OD2	-6.16	112.75	118.30
1	B	88	ASP	CB-CG-OD2	6.14	123.83	118.30
1	B	319	ARG	NE-CZ-NH1	6.08	123.34	120.30
1	B	623	ARG	NE-CZ-NH1	5.94	123.27	120.30
1	B	549	ASP	CB-CG-OD1	5.76	123.49	118.30
1	A	476	ASP	CB-CG-OD1	5.62	123.36	118.30
1	B	319	ARG	NE-CZ-NH2	-5.55	117.53	120.30
1	B	560	ARG	NE-CZ-NH1	5.53	123.06	120.30
1	A	663	ARG	NE-CZ-NH1	5.46	123.03	120.30
1	B	176	ASP	CB-CG-OD1	5.39	123.16	118.30
1	B	317	ASP	CB-CG-OD1	5.39	123.15	118.30
1	B	394	MET	CG-SD-CE	-5.32	91.69	100.20
1	A	541	ARG	NE-CZ-NH1	5.29	122.95	120.30
1	B	220	ASP	CB-CG-OD2	5.27	123.04	118.30
1	B	481	ASP	CB-CG-OD1	5.25	123.02	118.30
1	B	303	MET	CG-SD-CE	-5.17	91.92	100.20
1	A	611	ARG	NE-CZ-NH2	-5.07	117.76	120.30
1	B	107	LEU	CA-CB-CG	5.07	126.96	115.30
1	A	394	MET	CG-SD-CE	-5.07	92.09	100.20
1	B	110	ASP	CB-CG-OD1	5.05	122.84	118.30
1	B	541	ARG	NE-CZ-NH1	5.04	122.82	120.30
1	A	623	ARG	NE-CZ-NH1	5.03	122.82	120.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5437	0	5142	14	0
1	B	5414	0	5131	14	0
2	A	8	0	14	0	0

*Continued on next page...*



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	16	0	28	0	0
3	A	1	0	0	0	0
3	B	2	0	0	0	0
4	A	253	0	0	1	0
4	B	402	0	0	4	0
All	All	11533	0	10315	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:13:PHE:CG	1:A:72:VAL:HG21	2.32	0.65
1:A:74:GLN:OE1	1:A:79:TRP:NE1	2.29	0.65
1:A:13:PHE:CD1	1:A:72:VAL:CG2	2.82	0.62
1:B:14:HIS:O	1:B:17:ASN:O	2.18	0.61
1:A:13:PHE:CG	1:A:72:VAL:CG2	2.92	0.53
1:A:13:PHE:CD1	1:A:72:VAL:HG21	2.44	0.52
1:A:224:LEU:HD12	1:A:224:LEU:N	2.28	0.49
1:B:494:ALA:HB1	1:B:585:PRO:HG3	1.96	0.48
1:A:468:MET:HG2	1:A:498:HIS:CE1	2.50	0.46
1:B:369:GLU:HG3	4:B:2248:HOH:O	2.14	0.46
1:A:9:LEU:HD22	1:A:79:TRP:HZ3	1.80	0.46
1:A:531:THR:HB	1:A:532:PRO:HD2	1.97	0.46
1:A:72:VAL:HG12	1:A:73:SER:N	2.32	0.45
1:B:404:ALA:HB3	1:B:454:MET:HG2	1.98	0.45
1:B:654:PRO:HG2	1:B:657:LYS:HD2	2.01	0.43
1:B:599:HIS:HE1	4:B:2351:HOH:O	2.02	0.43
1:A:512:PHE:O	1:A:513:GLU:HB2	2.18	0.43
1:B:19:SER:HA	1:B:32:HIS:O	2.18	0.42
1:A:163:VAL:HG12	4:A:2086:HOH:O	2.20	0.42
1:B:9:LEU:HD21	1:B:103:LEU:HD13	2.02	0.42
1:B:483:LEU:HD23	1:B:483:LEU:C	2.40	0.41
1:B:240:THR:HB	1:B:245:ARG:HG2	2.02	0.41
1:B:599:HIS:CE1	4:B:2351:HOH:O	2.73	0.41
1:B:23:HIS:HA	1:B:27:ARG:O	2.21	0.41
1:A:23:HIS:HA	1:A:27:ARG:O	2.21	0.40
1:A:304:TRP:HA	1:A:306:TRP:CH2	2.57	0.40
1:B:124:GLN:HG2	4:B:2020:HOH:O	2.20	0.40
1:B:276:ARG:NH2	1:B:326:GLU:O	2.54	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	673/686 (98%)	653 (97%)	20 (3%)	0	100	100
1	B	671/686 (98%)	646 (96%)	24 (4%)	1 (0%)	51	42
All	All	1344/1372 (98%)	1299 (97%)	44 (3%)	1 (0%)	51	42

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	76	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	569/580 (98%)	559 (98%)	10 (2%)	59	53
1	B	567/580 (98%)	555 (98%)	12 (2%)	53	46
All	All	1136/1160 (98%)	1114 (98%)	22 (2%)	57	51

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	17	ASN
1	A	34	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	60	LYS
1	A	75	SER
1	A	181	PHE
1	A	182	PHE
1	A	224	LEU
1	A	276	ARG
1	A	469	TRP
1	A	628	LEU
1	B	14	HIS
1	B	74	GLN
1	B	144	LYS
1	B	181	PHE
1	B	182	PHE
1	B	221	LYS
1	B	270	LYS
1	B	276	ARG
1	B	297	SER
1	B	336	PRO
1	B	469	TRP
1	B	628	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	17	ASN
1	A	312	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 3 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	MPD	B	1679	-	7,7,7	0.37	0	9,10,10	1.02	1 (11%)
2	MPD	A	1682	-	7,7,7	0.36	0	9,10,10	0.53	0
2	MPD	B	1680	-	7,7,7	0.36	0	9,10,10	0.64	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MPD	B	1679	-	-	2/5/5/5	-
2	MPD	A	1682	-	-	3/5/5/5	-
2	MPD	B	1680	-	-	0/5/5/5	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
2	B	1679	MPD	O2-C2-C3	-2.03	102.18	109.80

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1682	MPD	O2-C2-C3-C4
2	A	1682	MPD	C1-C2-C3-C4
2	A	1682	MPD	CM-C2-C3-C4
2	B	1679	MPD	C2-C3-C4-C5
2	B	1679	MPD	C2-C3-C4-O4

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	674/686 (98%)	-0.09	20 (2%) 50 53	23, 37, 60, 91	2 (0%)
1	B	671/686 (97%)	-0.32	11 (1%) 72 74	14, 25, 50, 100	2 (0%)
All	All	1345/1372 (98%)	-0.20	31 (2%) 60 63	14, 31, 58, 100	4 (0%)

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	76	PRO	5.4
1	A	75	SER	4.1
1	B	13	PHE	3.9
1	A	79	TRP	3.7
1	B	16	ASN	3.6
1	A	74	GLN	3.5
1	A	13	PHE	3.5
1	B	15	GLN	3.3
1	A	77	ASP	3.3
1	B	76	PRO	3.2
1	A	298	PHE	3.1
1	B	99	ASP	3.1
1	A	11	PHE	3.0
1	B	298	PHE	3.0
1	B	77	ASP	3.0
1	A	358	ALA	3.0
1	A	99	ASP	2.9
1	A	470	ALA	2.7
1	A	73	SER	2.6
1	A	681	HIS	2.6
1	A	645	ARG	2.5
1	A	8	LEU	2.4
1	A	505	ILE	2.4
1	A	80	LEU	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	14	HIS	2.4
1	A	315	GLN	2.3
1	B	74	GLN	2.3
1	A	78	GLY	2.1
1	B	79	TRP	2.0
1	A	493	LEU	2.0
1	B	17	ASN	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	MPD	A	1682	8/8	0.89	0.16	47,54,56,57	0
2	MPD	B	1680	8/8	0.92	0.20	53,57,59,59	0
2	MPD	B	1679	8/8	0.93	0.11	45,53,56,56	0
3	CA	B	1682	1/1	0.99	0.05	32,32,32,32	0
3	CA	A	1683	1/1	0.99	0.09	26,26,26,26	0
3	CA	B	1681	1/1	1.00	0.09	18,18,18,18	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.