



# Full wwPDB X-ray Structure Validation Report ⓘ

Dec 17, 2023 – 06:47 pm GMT

PDB ID : 4AIK  
Title : Crystal structure of RovA from Yersinia in complex with an invasin promoter fragment  
Authors : Quade, N.; Mendonca, C.; Herbst, K.; Heroven, A.K.; Heinz, D.W.; Dersch, P.  
Deposited on : 2012-02-10  
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



| Metric                | Whole archive<br>(#Entries) | Similar resolution<br>(#Entries, resolution range(Å)) |
|-----------------------|-----------------------------|---|
| $R_{free}$            | 130704                      | 2469 (1.86-1.86)                                      |
| Clashscore            | 141614                      | 2625 (1.86-1.86)                                      |
| Ramachandran outliers | 138981                      | 2592 (1.86-1.86)                                      |
| Sidechain outliers    | 138945                      | 2592 (1.86-1.86)                                      |

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

| Mol | Chain | Length | Quality of chain   |
|-----|-------|--------|--|
| 1   | A     | 151    | 66% (green), 25% (yellow), 8% (orange), 1% (red), 0% (grey)  |
| 1   | B     | 151    | 72% (green), 22% (yellow), 6% (orange), 0% (red), 0% (grey)  |
| 2   | C     | 21     | 43% (green), 33% (yellow), 24% (orange), 0% (red), 0% (grey) |
| 3   | D     | 21     | 81% (green), 19% (yellow), 0% (orange), 0% (red), 0% (grey)  |

## 2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 3281 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TRANSCRIPTIONAL REGULATOR SLYA.

| Mol | Chain | Residues | Atoms |     |     |     | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---------|---------|-------|
|     |       |          | Total | C   | N   | O   |         |         |       |
| 1   | A     | 139      | 1106  | 695 | 202 | 209 | 0       | 1       | 0     |
| 1   | B     | 142      | 1136  | 713 | 206 | 217 | 0       | 2       | 0     |

There are 20 discrepancies between the modelled and reference sequences:

| Chain | Residue | Modelled | Actual | Comment             | Reference  |
|-------|---------|----------|--------|---------------------|------------|
| A     | 144     | LEU      | -      | expression tag      | UNP B1JJ73 |
| A     | 145     | GLU      | -      | expression tag      | UNP B1JJ73 |
| A     | 146     | HIS      | -      | expression tag      | UNP B1JJ73 |
| A     | 147     | HIS      | -      | expression tag      | UNP B1JJ73 |
| A     | 148     | HIS      | -      | expression tag      | UNP B1JJ73 |
| A     | 149     | HIS      | -      | expression tag      | UNP B1JJ73 |
| A     | 150     | HIS      | -      | expression tag      | UNP B1JJ73 |
| A     | 151     | HIS      | -      | expression tag      | UNP B1JJ73 |
| A     | 81      | SER      | CYS    | engineered mutation | UNP B1JJ73 |
| A     | 108     | SER      | CYS    | engineered mutation | UNP B1JJ73 |
| B     | 144     | LEU      | -      | expression tag      | UNP B1JJ73 |
| B     | 145     | GLU      | -      | expression tag      | UNP B1JJ73 |
| B     | 146     | HIS      | -      | expression tag      | UNP B1JJ73 |
| B     | 147     | HIS      | -      | expression tag      | UNP B1JJ73 |
| B     | 148     | HIS      | -      | expression tag      | UNP B1JJ73 |
| B     | 149     | HIS      | -      | expression tag      | UNP B1JJ73 |
| B     | 150     | HIS      | -      | expression tag      | UNP B1JJ73 |
| B     | 151     | HIS      | -      | expression tag      | UNP B1JJ73 |
| B     | 81      | SER      | CYS    | engineered mutation | UNP B1JJ73 |
| B     | 108     | SER      | CYS    | engineered mutation | UNP B1JJ73 |

- Molecule 2 is a DNA chain called ROVA PROMOTER FRAGMENT, 5'-D(\*AP\*TP\*GP\*A P\*TP\*AP\*TP\*TP \*AP\*TP\*\*TP\*TP\*AP\*TP\*AP\*TP\*GP\*AP\*TP\*AP\*A)-3'.

| Mol | Chain | Residues | Atoms |     |    |     |    | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|-----|----|-----|----|---------|---------|-------|
| 2   | C     | 21       | Total | C   | N  | O   | P  | 0       | 0       | 0     |
|     |       |          | 430   | 210 | 75 | 125 | 20 |         |         |       |

- Molecule 3 is a DNA chain called ROVA PROMOTER FRAGMENT, 5'-D(\*TP\*TP\*TP\*A P\*TP\*CP\*AP\*TP \*AP\*TP\*AP\*AP\*AP\*TP\*AP\*AP\*TP\*AP\*TP\*CP\*A)-3'.

| Mol | Chain | Residues | Atoms |     |    |     |    | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|-----|----|-----|----|---------|---------|-------|
| 3   | D     | 21       | Total | C   | N  | O   | P  | 0       | 0       | 0     |
|     |       |          | 425   | 208 | 74 | 123 | 20 |         |         |       |

- Molecule 4 is water.


| Mol | Chain | Residues | Atoms |    | ZeroOcc | AltConf |
|-----|-------|----------|-------|----|---------|---------|
| 4   | A     | 93       | Total | O  | 0       | 0       |
|     |       |          | 93    | 93 |         |         |
| 4   | B     | 66       | Total | O  | 0       | 0       |
|     |       |          | 66    | 66 |         |         |
| 4   | C     | 12       | Total | O  | 0       | 0       |
|     |       |          | 12    | 12 |         |         |
| 4   | D     | 13       | Total | O  | 0       | 0       |
|     |       |          | 13    | 13 |         |         |

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: TRANSCRIPTIONAL REGULATOR SLYA

Chain A: 



- Molecule 1: TRANSCRIPTIONAL REGULATOR SLYA

Chain B: 




- Molecule 2: ROVA PROMOTER FRAGMENT, 5'-D(\*AP\*TP\*GP\*AP\*TP\*AP\*TP\*TP \*AP\*TP\*\*TP\*TP\*AP\*TP\*AP\*TP\*GP\*AP\*TP\*AP\*A)-3'

Chain C: 



- Molecule 3: ROVA PROMOTER FRAGMENT, 5'-D(\*TP\*TP\*TP\*AP\*TP\*CP\*AP\*TP \*AP\*TP\*AP\*AP\*AP\*TP\*AP\*AP\*TP\*AP\*TP\*CP\*A)-3'

Chain D: 



## 4 Data and refinement statistics i

| Property  | Value  | Source           |
|---|--|------------------|
| Space group   | P 3  | Depositor        |
| Cell constants<br>a, b, c, $\alpha$ , $\beta$ , $\gamma$                | 88.68Å 88.68Å 67.50Å<br>90.00° 90.00° 120.00°                  | Depositor        |
| Resolution (Å)  | 76.81 – 1.85<br>26.67 – 1.85                                   | Depositor<br>EDS |
| % Data completeness<br>(in resolution range)                            | 98.0 (76.81-1.85)<br>98.5 (26.67-1.85)                         | Depositor<br>EDS |
| $R_{merge}$   | 0.03   | Depositor        |
| $R_{sym}$   | (Not available)  | Depositor        |
| $\langle I/\sigma(I) \rangle$ <sup>1</sup>                              | 2.62 (at 1.85Å)  | Xtrriage         |
| Refinement program  | REFMAC 5.5.0109  | Depositor        |
| R, $R_{free}$   | 0.198 , 0.230<br>0.255 , 0.287                                 | Depositor<br>DCC |
| $R_{free}$ test set   | 2510 reflections (5.02%)                                       | wwPDB-VP         |
| Wilson B-factor (Å <sup>2</sup> )                                       | 32.4   | Xtrriage         |
| Anisotropy  | 0.607  | Xtrriage         |
| Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> ) | 0.33 , 14.6  | EDS              |
| L-test for twinning <sup>2</sup>  | $\langle  L  \rangle = 0.53$ , $\langle L^2 \rangle = 0.37$    | Xtrriage         |
| Estimated twinning fraction   | 0.008 for -h,-k,l<br>0.196 for h,-h-k,-l<br>0.017 for -k,-h,-l | Xtrriage         |
| Reported twinning fraction  | 0.502 for H, K, L<br>0.498 for h,-h-k,-l                       | Depositor        |
| Outliers  | 0 of 49952 reflections   | Xtrriage         |
| $F_o, F_c$ correlation  | 0.94   | EDS              |
| Total number of atoms   | 3281   | wwPDB-VP         |
| Average B, all atoms (Å <sup>2</sup> )                                  | 50.0   | wwPDB-VP         |

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.87% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Chain | Bond lengths |         | Bond angles |               |
|-----|-------|--------------|---------|-------------|---------------|
|     |       | RMSZ         | # Z  >5 | RMSZ        | # Z  >5       |
| 1   | A     | 0.46         | 0/1121  | 0.64        | 0/1516        |
| 1   | B     | 0.46         | 0/1154  | 0.60        | 0/1560        |
| 2   | C     | 0.64         | 0/482   | 1.42        | 6/743 (0.8%)  |
| 3   | D     | 0.66         | 0/476   | 1.27        | 1/732 (0.1%)  |
| All | All   | 0.52         | 0/3233  | 0.92        | 7/4551 (0.2%) |

There are no bond length outliers.

All (7) bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms      | Z    | Observed(°) | Ideal(°) |
|-----|-------|-----|------|------------|------|-------------|----------|
| 2   | C     | 9   | DA   | P-O3'-C3'  | 8.23 | 129.58      | 119.70   |
| 3   | D     | 11  | DA   | P-O3'-C3'  | 7.54 | 128.75      | 119.70   |
| 2   | C     | 18  | DA   | P-O3'-C3'  | 7.36 | 128.53      | 119.70   |
| 2   | C     | 3   | DG   | P-O3'-C3'  | 6.66 | 127.70      | 119.70   |
| 2   | C     | 6   | DA   | P-O3'-C3'  | 5.60 | 126.42      | 119.70   |
| 2   | C     | 5   | DT   | O4'-C1'-N1 | 5.54 | 111.88      | 108.00   |
| 2   | C     | 20  | DA   | P-O3'-C3'  | 5.02 | 125.72      | 119.70   |

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 1   | A     | 1106  | 0        | 1172     | 31      | 0            |
| 1   | B     | 1136  | 0        | 1202     | 33      | 0            |

*Continued on next page...*

*Continued from previous page...*

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 2   | C     | 430   | 0        | 243      | 9       | 0            |
| 3   | D     | 425   | 0        | 242      | 2       | 0            |
| 4   | A     | 93    | 0        | 0        | 3       | 0            |
| 4   | B     | 66    | 0        | 0        | 4       | 0            |
| 4   | C     | 12    | 0        | 0        | 0       | 0            |
| 4   | D     | 13    | 0        | 0        | 0       | 0            |
| All | All   | 3281  | 0        | 2859     | 61      | 0            |

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

| Atom-1             | Atom-2           | Interatomic distance (Å) | Clash overlap (Å) |
|--------------------|------------------|--------------------------|-------------------|
| 1:A:131:ASP:O      | 1:A:135:LYS:HG2  | 1.66                     | 0.95              |
| 1:B:65:ARG:HG2     | 1:B:65:ARG:HH11  | 1.43                     | 0.84              |
| 1:A:140:LEU:O      | 1:A:141:GLN:HB3  | 1.76                     | 0.84              |
| 1:A:102:GLN:HG2    | 4:A:2072:HOH:O   | 1.79                     | 0.82              |
| 1:A:129:LEU:HD21   | 1:B:122:GLU:HB3  | 1.62                     | 0.81              |
| 1:B:27:LEU:HB3     | 1:B:29:LEU:HD13  | 1.68                     | 0.74              |
| 2:C:18:DA:H2''     | 2:C:19:DT:H5''   | 1.75                     | 0.68              |
| 1:B:27:LEU:HD21    | 4:B:2052:HOH:O   | 1.93                     | 0.68              |
| 2:C:16:DT:H2''     | 2:C:17:DG:C8     | 2.29                     | 0.67              |
| 1:B:20[B]:ILE:HG22 | 1:B:110:THR:HG21 | 1.77                     | 0.66              |
| 1:A:118:ILE:HD13   | 1:B:133:LEU:HD11 | 1.77                     | 0.66              |
| 1:A:53:ALA:HB1     | 1:A:58:ILE:O     | 1.95                     | 0.66              |
| 1:B:65:ARG:HH11    | 1:B:65:ARG:CG    | 2.08                     | 0.65              |
| 1:A:15:VAL:HG12    | 1:B:137:ILE:CG2  | 2.29                     | 0.63              |
| 1:A:31:GLN:O       | 1:A:35:VAL:HG23  | 2.03                     | 0.59              |
| 1:A:60:GLN:HG3     | 1:A:61:PRO:HD3   | 1.84                     | 0.58              |
| 1:A:102:GLN:NE2    | 4:A:2071:HOH:O   | 2.37                     | 0.58              |
| 1:A:15:VAL:HG12    | 1:B:137:ILE:HG22 | 1.86                     | 0.57              |
| 1:B:35:VAL:HG12    | 1:B:56:ILE:HG23  | 1.87                     | 0.57              |
| 1:A:96:SER:HA      | 1:A:99:ILE:HD12  | 1.86                     | 0.56              |
| 1:A:125:VAL:HG13   | 1:A:129:LEU:HD13 | 1.87                     | 0.56              |
| 1:B:22:HIS:O       | 1:B:25:LYS:HB2   | 2.06                     | 0.56              |
| 1:A:118:ILE:HG22   | 1:B:132:LYS:HD2  | 1.88                     | 0.55              |
| 1:A:118:ILE:HG22   | 1:B:132:LYS:CD   | 2.38                     | 0.54              |
| 2:C:20:DA:H61      | 3:D:3:DT:H3      | 1.55                     | 0.53              |
| 1:A:111:ARG:NH2    | 4:A:2041:HOH:O   | 2.22                     | 0.53              |
| 1:B:31:GLN:NE2     | 4:B:2010:HOH:O   | 2.42                     | 0.53              |

*Continued on next page...*



*Continued from previous page...*

| Atom-1             | Atom-2              | Interatomic distance (Å) | Clash overlap (Å) |
|--------------------|---------------------|--------------------------|-------------------|
| 2:C:9:DA:H1'       | 2:C:10:DT:C5'       | 2.40                     | 0.52              |
| 2:C:9:DA:H1'       | 2:C:10:DT:H5'       | 1.92                     | 0.52              |
| 1:B:92:LEU:HD22    | 1:B:100:ILE:HD11    | 1.91                     | 0.50              |
| 1:B:20[B]:ILE:HD11 | 1:B:34:TRP:CZ3      | 2.47                     | 0.49              |
| 1:A:9:LEU:HG       | 1:B:130:ILE:HD11    | 1.95                     | 0.48              |
| 3:D:20:DC:H2''     | 3:D:21:DA:N7        | 2.29                     | 0.48              |
| 1:A:122:GLU:HB3    | 1:B:129:LEU:HD21    | 1.96                     | 0.47              |
| 1:A:15:VAL:HG12    | 1:B:137:ILE:HG21    | 1.96                     | 0.47              |
| 1:A:69:GLN:O       | 1:A:72:GLU:HB2      | 2.14                     | 0.47              |
| 1:A:119:SER:O      | 1:A:123[B]:ILE:HG23 | 2.14                     | 0.47              |
| 1:A:140:LEU:O      | 1:A:141:GLN:CB      | 2.53                     | 0.46              |
| 2:C:2:DT:H2''      | 2:C:3:DG:C8         | 2.50                     | 0.46              |
| 1:B:17:ARG:HD2     | 1:B:31:GLN:OE1      | 2.15                     | 0.46              |
| 1:B:94:GLU:H       | 1:B:94:GLU:CD       | 2.19                     | 0.46              |
| 1:B:20[B]:ILE:HD13 | 1:B:107:ILE:CG1     | 2.46                     | 0.45              |
| 1:A:63:LEU:O       | 1:A:67:LEU:HG       | 2.17                     | 0.45              |
| 1:B:20[B]:ILE:HD13 | 1:B:107:ILE:HG12    | 1.98                     | 0.44              |
| 1:A:69:GLN:O       | 1:A:73:LYS:HG2      | 2.17                     | 0.44              |
| 1:B:140:LEU:HD23   | 1:B:143:LYS:HE3     | 2.00                     | 0.44              |
| 2:C:9:DA:H2''      | 2:C:10:DT:OP2       | 2.17                     | 0.44              |
| 1:B:47:GLN:HG2     | 1:B:51:GLN:HB2      | 1.99                     | 0.44              |
| 1:A:42:ARG:NH1     | 1:B:4:THR:HG21      | 2.34                     | 0.43              |
| 2:C:6:DA:H2'       | 2:C:7:DT:H71        | 2.00                     | 0.43              |
| 1:B:27:LEU:CD2     | 4:B:2052:HOH:O      | 2.57                     | 0.43              |
| 1:A:59:GLU:HB2     | 1:A:61:PRO:HD2      | 2.01                     | 0.42              |
| 1:A:73:LYS:HB2     | 1:A:75:LEU:HG       | 2.02                     | 0.42              |
| 1:A:130:ILE:HG21   | 1:B:8:ASP:HB3       | 2.01                     | 0.42              |
| 1:B:52:LEU:HD23    | 1:B:63:LEU:HD21     | 2.02                     | 0.41              |
| 1:A:126:LEU:HD11   | 1:B:9:LEU:HD21      | 2.02                     | 0.41              |
| 1:B:65:ARG:CG      | 1:B:65:ARG:NH1      | 2.74                     | 0.41              |
| 1:A:8:ASP:HB3      | 1:B:130:ILE:HG21    | 2.03                     | 0.40              |
| 2:C:19:DT:H2''     | 2:C:20:DA:C8        | 2.56                     | 0.40              |
| 1:B:102:GLN:NE2    | 4:B:2052:HOH:O      | 2.38                     | 0.40              |

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed      | Favoured  | Allowed | Outliers | Percentiles |     |
|-----|-------|---------------|-----------|---------|----------|-------------|-----|
| 1   | A     | 138/151 (91%) | 136 (99%) | 2 (1%)  | 0        | 100         | 100 |
| 1   | B     | 142/151 (94%) | 140 (99%) | 2 (1%)  | 0        | 100         | 100 |
| All | All   | 280/302 (93%) | 276 (99%) | 4 (1%)  | 0        | 100         | 100 |

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed      | Rotameric | Outliers | Percentiles |    |
|-----|-------|---------------|-----------|----------|-------------|----|
| 1   | A     | 126/137 (92%) | 116 (92%) | 10 (8%)  | 12          | 2  |
| 1   | B     | 130/137 (95%) | 127 (98%) | 3 (2%)   | 50          | 34 |
| All | All   | 256/274 (93%) | 243 (95%) | 13 (5%)  | 24          | 9  |

All (13) residues with a non-rotameric sidechain are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1   | A     | 5   | LEU  |
| 1   | A     | 29  | LEU  |
| 1   | A     | 59  | GLU  |
| 1   | A     | 60  | GLN  |
| 1   | A     | 65  | ARG  |
| 1   | A     | 83  | ASN  |
| 1   | A     | 86  | ARG  |
| 1   | A     | 97  | SER  |

*Continued on next page...*

*Continued from previous page...*

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1   | A     | 121 | ASP  |
| 1   | A     | 141 | GLN  |
| 1   | B     | 65  | ARG  |
| 1   | B     | 85  | ARG  |
| 1   | B     | 97  | SER  |

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1   | A     | 60  | GLN  |
| 1   | A     | 102 | GLN  |
| 1   | A     | 136 | ASN  |
| 1   | B     | 31  | GLN  |
| 1   | B     | 41  | ASN  |

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.5 Other polymers

Unable to reproduce the depositors R factor - this section is therefore empty.