



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 2, 2024 – 01:48 pm GMT

PDB ID : 8AI2
Title : Crystal structure of radical SAM epimerase EpeE from *Bacillus subtilis* with [4Fe-4S] clusters, S-adenosyl-L-homocysteine and RiPP peptide 5 bound
Authors : Polsinelli, I.; Fyfe, C.D.; Legrand, P.; Kubiak, X.; Chavas, L.M.G.; Berteau, O.; Benjdia, A.
Deposited on : 2022-07-25
Resolution : 2.39 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.39 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	344	 2% 84% 8% 8%
1	B	344	 5% 85% 9% 6%
2	C	11	 36% 18% 45% 36%
2	D	11	 18% 18% 36% 45%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	EDO	B	414	-	-	-	X
5	EDO	B	416	-	-	-	X
5	EDO	B	419	-	-	-	X
6	PEG	A	412	-	-	-	X

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 5858 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative peptide biosynthesis protein YydG.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	317	2635	1694	425	499	17	0	4	0
1	B	323	2681	1727	431	506	17	0	3	0

There are 50 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-24	MET	-	initiating methionine	UNP Q45595
A	-23	ALA	-	expression tag	UNP Q45595
A	-22	SER	-	expression tag	UNP Q45595
A	-21	TRP	-	expression tag	UNP Q45595
A	-20	SER	-	expression tag	UNP Q45595
A	-19	HIS	-	expression tag	UNP Q45595
A	-18	PRO	-	expression tag	UNP Q45595
A	-17	GLN	-	expression tag	UNP Q45595
A	-16	PHE	-	expression tag	UNP Q45595
A	-15	GLU	-	expression tag	UNP Q45595
A	-14	LYS	-	expression tag	UNP Q45595
A	-13	SER	-	expression tag	UNP Q45595
A	-12	GLY	-	expression tag	UNP Q45595
A	-11	GLY	-	expression tag	UNP Q45595
A	-10	GLY	-	expression tag	UNP Q45595
A	-9	GLY	-	expression tag	UNP Q45595
A	-8	GLY	-	expression tag	UNP Q45595
A	-7	GLU	-	expression tag	UNP Q45595
A	-6	ASN	-	expression tag	UNP Q45595
A	-5	LEU	-	expression tag	UNP Q45595
A	-4	TYR	-	expression tag	UNP Q45595
A	-3	PHE	-	expression tag	UNP Q45595
A	-2	GLN	-	expression tag	UNP Q45595
A	-1	GLY	-	expression tag	UNP Q45595
A	0	HIS	-	expression tag	UNP Q45595

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-24	MET	-	initiating methionine	UNP Q45595
B	-23	ALA	-	expression tag	UNP Q45595
B	-22	SER	-	expression tag	UNP Q45595
B	-21	TRP	-	expression tag	UNP Q45595
B	-20	SER	-	expression tag	UNP Q45595
B	-19	HIS	-	expression tag	UNP Q45595
B	-18	PRO	-	expression tag	UNP Q45595
B	-17	GLN	-	expression tag	UNP Q45595
B	-16	PHE	-	expression tag	UNP Q45595
B	-15	GLU	-	expression tag	UNP Q45595
B	-14	LYS	-	expression tag	UNP Q45595
B	-13	SER	-	expression tag	UNP Q45595
B	-12	GLY	-	expression tag	UNP Q45595
B	-11	GLY	-	expression tag	UNP Q45595
B	-10	GLY	-	expression tag	UNP Q45595
B	-9	GLY	-	expression tag	UNP Q45595
B	-8	GLY	-	expression tag	UNP Q45595
B	-7	GLU	-	expression tag	UNP Q45595
B	-6	ASN	-	expression tag	UNP Q45595
B	-5	LEU	-	expression tag	UNP Q45595
B	-4	TYR	-	expression tag	UNP Q45595
B	-3	PHE	-	expression tag	UNP Q45595
B	-2	GLN	-	expression tag	UNP Q45595
B	-1	GLY	-	expression tag	UNP Q45595
B	0	HIS	-	expression tag	UNP Q45595

- Molecule 2 is a protein called Putative exported peptide YydF.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	7	Total	C	N	O	0	0	0
			68	44	13	11			
2	D	6	Total	C	N	O	0	0	0
			59	38	11	10			

- Molecule 3 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe₄S₄) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Fe S 8 4 4	0	0
3	A	1	Total Fe S 8 4 4	0	0
3	B	1	Total Fe S 8 4 4	0	0
3	B	1	Total Fe S 8 4 4	0	0

- Molecule 4 is S-ADENOSYL-L-HOMOCYSTEINE (three-letter code: SAH) (formula: $C_{14}H_{20}N_6O_5S$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S	0	0
			26	14	6	5	1		
4	B	1	Total	C	N	O	S	0	0
			26	14	6	5	1		

- Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		
5	D	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 7 4 3	0	0
6	A	1	Total C O 7 4 3	0	0
6	A	1	Total C O 7 4 3	0	0
6	B	1	Total C O 7 4 3	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	117	Total O 117 117	0	0
7	B	83	Total O 84 84	0	1
7	C	4	Total O 4 4	0	0
7	D	6	Total O 6 6	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Putative peptide biosynthesis protein YydG




- Molecule 1: Putative peptide biosynthesis protein YydG



- Molecule 2: Putative exported peptide YydF



- Molecule 2: Putative exported peptide YydF



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	79.73Å 92.37Å 126.72Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.19 – 2.39 46.19 – 2.39	Depositor EDS
% Data completeness (in resolution range)	99.9 (46.19-2.39) 99.9 (46.19-2.39)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.98 (at 2.39Å)	Xtrriage
Refinement program	BUSTER 2.10.4	Depositor
R, R_{free}	0.186 , 0.222 0.181 , 0.222	Depositor DCC
R_{free} test set	1881 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	54.6	Xtrriage
Anisotropy	0.457	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 47.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5858	wwPDB-VP
Average B, all atoms (Å ²)	63.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.32% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, SAH, SF4, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.43	0/2700	0.57	0/3638
1	B	0.42	0/2745	0.57	0/3699
2	C	0.56	0/69	0.72	0/90
2	D	0.52	0/60	0.67	0/79
All	All	0.43	0/5574	0.57	0/7506

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2635	0	2581	18	0
1	B	2681	0	2624	19	0
2	C	68	0	69	5	0
2	D	59	0	56	4	0
3	A	16	0	0	0	0
3	B	16	0	0	0	0
4	A	26	0	19	1	0
4	B	26	0	19	0	0
5	A	28	0	42	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	60	0	90	5	0
5	D	4	0	6	0	0
6	A	21	0	30	1	0
6	B	7	0	10	1	0
7	A	117	0	0	3	0
7	B	84	0	0	1	0
7	C	4	0	0	0	0
7	D	6	0	0	0	0
All	All	5858	0	5546	39	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:240:GLN:HA	5:B:406:EDO:H21	1.60	0.84
1:B:212[B]:VAL:HG22	1:B:223:CYS:SG	2.20	0.80
1:B:157:ASN:HA	5:B:412:EDO:H21	1.70	0.73
1:B:140:PRO:HA	5:B:419:EDO:H12	1.73	0.69
1:A:230:THR:HA	1:A:297:ASN:HA	1.81	0.61
1:B:231:LYS:HB2	1:B:302:ILE:HD12	1.81	0.61
1:B:52:GLN:HA	6:B:418:PEG:H42	1.83	0.60
1:A:252:ASN:HD22	1:A:255:LEU:H	1.52	0.58
1:B:207:PRO:HG2	2:D:6:GLU:HB2	1.87	0.57
1:A:238:TYR:HE1	6:A:412:PEG:H12	1.71	0.56
1:A:44:GLU:O	5:A:404:EDO:H21	2.08	0.53
1:B:244:ARG:HB2	5:B:406:EDO:H11	1.91	0.52
1:A:214:HIS:HE1	7:A:611:HOH:O	1.93	0.50
1:A:180:VAL:HA	1:A:184:LYS:HD3	1.95	0.49
1:B:223:CYS:SG	2:D:11:LEU:HD11	2.53	0.49
5:B:410:EDO:H11	7:B:567:HOH:O	2.12	0.49
1:B:208:GLY:O	2:D:7:ASN:HB3	2.12	0.48
1:A:201:GLU:HB2	7:A:588:HOH:O	2.14	0.48
1:A:207:PRO:HG2	2:C:6:GLU:HB2	1.94	0.48
1:B:180:VAL:HA	1:B:184:LYS:HD3	1.96	0.47
1:A:231:LYS:HB2	1:A:302:ILE:HD12	1.94	0.47
1:B:212[B]:VAL:CG2	1:B:223:CYS:SG	2.98	0.46
1:B:230:THR:HA	1:B:297:ASN:HA	1.97	0.46
1:B:39:ARG:HD3	1:B:70:GLU:OE2	2.17	0.44
1:A:125:LYS:HD3	7:A:604:HOH:O	2.16	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:223:CYS:SG	2:C:11:LEU:HD11	2.59	0.43
4:A:403:SAH:H5'2	4:A:403:SAH:HB2	1.79	0.43
1:A:44:GLU:HG3	5:A:404:EDO:H12	2.01	0.42
1:B:276:ILE:HG23	1:B:281:ILE:HB	2.02	0.42
1:A:207:PRO:HB3	2:C:5:LYS:HB2	2.01	0.42
1:A:225:PRO:HD3	2:C:9:TRP:HB3	2.02	0.42
1:A:252:ASN:ND2	1:A:255:LEU:H	2.15	0.42
1:A:91:GLY:HA2	1:A:97:VAL:CG2	2.50	0.42
1:A:208:GLY:O	2:C:7:ASN:HB3	2.20	0.42
1:B:293:GLY:O	1:B:297:ASN:HB3	2.20	0.42
1:A:293:GLY:O	1:A:297:ASN:HB3	2.20	0.41
1:B:58:GLY:HA3	1:B:59:GLY:C	2.41	0.41
1:B:91:GLY:HA2	1:B:97:VAL:CG2	2.51	0.40
1:B:223:CYS:O	2:D:10:ILE:HG12	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	319/344 (93%)	314 (98%)	5 (2%)	0	100	100
1	B	324/344 (94%)	318 (98%)	6 (2%)	0	100	100
2	C	5/11 (46%)	3 (60%)	2 (40%)	0	100	100
2	D	4/11 (36%)	3 (75%)	1 (25%)	0	100	100
All	All	652/710 (92%)	638 (98%)	14 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	300/316 (95%)	291 (97%)	9 (3%)	41	61
1	B	304/316 (96%)	297 (98%)	7 (2%)	50	70
2	C	7/11 (64%)	7 (100%)	0	100	100
2	D	6/11 (54%)	6 (100%)	0	100	100
All	All	617/654 (94%)	601 (97%)	16 (3%)	46	66

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	73	GLU
1	A	96	LYS
1	A	144	ILE
1	A	199	GLU
1	A	210	ASP
1	A	252	ASN
1	A	280	ASP
1	A	317	PHE
1	B	-5	LEU
1	B	66	LYS
1	B	73	GLU
1	B	199	GLU
1	B	210	ASP
1	B	280	ASP
1	B	311	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	252	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

33 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	EDO	B	415	-	3,3,3	0.57	0	2,2,2	0.31	0
5	EDO	B	405	-	3,3,3	0.56	0	2,2,2	0.22	0
6	PEG	A	408	-	6,6,6	0.35	0	5,5,5	0.23	0
5	EDO	A	405	-	3,3,3	0.68	0	2,2,2	0.13	0
5	EDO	A	404	-	3,3,3	0.45	0	2,2,2	0.26	0
3	SF4	A	402	1	0,12,12	-	-	-		
5	EDO	B	406	-	3,3,3	1.03	0	2,2,2	0.47	0
3	SF4	B	401	4,1	0,12,12	-	-	-		
5	EDO	B	412	-	3,3,3	0.50	0	2,2,2	0.46	0
6	PEG	A	412	-	6,6,6	0.31	0	5,5,5	0.28	0
5	EDO	A	406	-	3,3,3	0.76	0	2,2,2	0.03	0
5	EDO	B	411	-	3,3,3	0.52	0	2,2,2	0.38	0
5	EDO	A	409	-	3,3,3	0.67	0	2,2,2	0.15	0
5	EDO	B	407	-	3,3,3	0.72	0	2,2,2	0.13	0
5	EDO	B	410	-	3,3,3	0.69	0	2,2,2	0.05	0
3	SF4	B	402	1	0,12,12	-	-	-		
5	EDO	B	419	-	3,3,3	0.66	0	2,2,2	0.34	0
5	EDO	B	408	-	3,3,3	0.61	0	2,2,2	0.29	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	PEG	B	418	-	6,6,6	0.22	0	5,5,5	0.13	0
4	SAH	A	403	3	24,28,28	0.80	1 (4%)	25,40,40	1.06	2 (8%)
5	EDO	A	407	-	3,3,3	0.76	0	2,2,2	0.15	0
5	EDO	D	101	-	3,3,3	0.62	0	2,2,2	0.35	0
5	EDO	A	413	-	3,3,3	0.67	0	2,2,2	0.17	0
5	EDO	B	409	-	3,3,3	0.68	0	2,2,2	0.12	0
5	EDO	B	404	-	3,3,3	0.78	0	2,2,2	0.21	0
3	SF4	A	401	4,1	0,12,12	-	-	-	-	-
5	EDO	B	416	-	3,3,3	0.69	0	2,2,2	0.06	0
6	PEG	A	410	-	6,6,6	0.24	0	5,5,5	0.12	0
5	EDO	A	411	-	3,3,3	0.64	0	2,2,2	0.22	0
5	EDO	B	413	-	3,3,3	0.78	0	2,2,2	0.10	0
5	EDO	B	417	-	3,3,3	0.59	0	2,2,2	0.11	0
5	EDO	B	414	-	3,3,3	0.54	0	2,2,2	0.40	0
4	SAH	B	403	3	24,28,28	0.90	1 (4%)	25,40,40	1.04	2 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	B	415	-	-	1/1/1/1	-
5	EDO	B	405	-	-	0/1/1/1	-
6	PEG	A	408	-	-	2/4/4/4	-
5	EDO	A	405	-	-	1/1/1/1	-
5	EDO	A	404	-	-	0/1/1/1	-
3	SF4	A	402	1	-	-	0/6/5/5
5	EDO	B	406	-	-	0/1/1/1	-
3	SF4	B	401	4,1	-	-	0/6/5/5
5	EDO	B	412	-	-	0/1/1/1	-
6	PEG	A	412	-	-	2/4/4/4	-
5	EDO	A	406	-	-	0/1/1/1	-
5	EDO	B	411	-	-	0/1/1/1	-
5	EDO	A	409	-	-	1/1/1/1	-
5	EDO	B	407	-	-	0/1/1/1	-
5	EDO	B	410	-	-	1/1/1/1	-
5	EDO	B	419	-	-	0/1/1/1	-
3	SF4	B	402	1	-	-	0/6/5/5
5	EDO	B	408	-	-	0/1/1/1	-
6	PEG	B	418	-	-	2/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	SAH	A	403	3	-	4/11/31/31	0/3/3/3
5	EDO	A	407	-	-	0/1/1/1	-
5	EDO	D	101	-	-	1/1/1/1	-
5	EDO	A	413	-	-	1/1/1/1	-
5	EDO	B	409	-	-	0/1/1/1	-
5	EDO	B	404	-	-	1/1/1/1	-
3	SF4	A	401	4,1	-	-	0/6/5/5
5	EDO	B	416	-	-	0/1/1/1	-
6	PEG	A	410	-	-	2/4/4/4	-
5	EDO	A	411	-	-	0/1/1/1	-
5	EDO	B	413	-	-	1/1/1/1	-
5	EDO	B	417	-	-	0/1/1/1	-
5	EDO	B	414	-	-	1/1/1/1	-
4	SAH	B	403	3	-	2/11/31/31	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	403	SAH	CB-CA	3.49	1.60	1.53
4	A	403	SAH	CB-CA	2.61	1.58	1.53

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	403	SAH	CB-CA-N	3.41	119.12	110.17
4	B	403	SAH	CB-CA-N	2.85	117.63	110.17
4	B	403	SAH	C5-C6-N6	2.49	124.14	120.35
4	A	403	SAH	C5-C6-N6	2.20	123.70	120.35

There are no chirality outliers.

All (23) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	B	418	PEG	O2-C3-C4-O4
6	A	412	PEG	O2-C3-C4-O4
5	A	405	EDO	O1-C1-C2-O2
5	A	409	EDO	O1-C1-C2-O2
5	D	101	EDO	O1-C1-C2-O2
4	A	403	SAH	C-CA-CB-CG
4	B	403	SAH	C-CA-CB-CG

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
5	B	415	EDO	O1-C1-C2-O2
6	B	418	PEG	C4-C3-O2-C2
4	A	403	SAH	N-CA-CB-CG
6	A	408	PEG	C1-C2-O2-C3
6	A	410	PEG	C4-C3-O2-C2
6	A	412	PEG	C1-C2-O2-C3
6	A	410	PEG	C1-C2-O2-C3
6	A	408	PEG	O1-C1-C2-O2
4	A	403	SAH	CB-CG-SD-C5'
5	A	413	EDO	O1-C1-C2-O2
5	B	404	EDO	O1-C1-C2-O2
5	B	413	EDO	O1-C1-C2-O2
5	B	414	EDO	O1-C1-C2-O2
4	A	403	SAH	CA-CB-CG-SD
4	B	403	SAH	N-CA-CB-CG
5	B	410	EDO	O1-C1-C2-O2



There are no ring outliers.


8 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	404	EDO	2	0
5	B	406	EDO	2	0
5	B	412	EDO	1	0
6	A	412	PEG	1	0
5	B	410	EDO	1	0
5	B	419	EDO	1	0
6	B	418	PEG	1	0
4	A	403	SAH	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	317/344 (92%)	-0.04	6 (1%) 66 64	44, 57, 77, 106	3 (0%)
1	B	323/344 (93%)	0.14	17 (5%) 26 25	43, 62, 104, 119	3 (0%)
2	C	7/11 (63%)	2.34	4 (57%) 0 0	67, 70, 74, 76	7 (100%)
2	D	6/11 (54%)	2.19	2 (33%) 0 0	76, 79, 82, 85	6 (100%)
All	All	653/710 (91%)	0.10	29 (4%) 34 33	43, 60, 93, 119	19 (2%)

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	317	PHE	5.6
2	D	11	LEU	5.3
1	B	306	TYR	5.0
1	A	283	TYR	4.4
2	C	8	ARG	4.2
1	A	317	PHE	4.0
1	A	1	MET	3.5
1	A	313	TYR	3.1
1	B	203	SER	2.9
2	C	5	LYS	2.9
1	B	312	TYR	2.9
1	B	253	LEU	2.9
2	C	9	TRP	2.8
1	B	305	PHE	2.7
1	B	313	TYR	2.5
2	C	6	GLU	2.4
1	B	311	LYS	2.3
2	D	9	TRP	2.3
1	A	201	GLU	2.3
1	B	256	PHE	2.3
1	B	309	MET	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	316	ASN	2.3
1	B	308	TYR	2.2
1	B	310	GLU	2.2
1	B	202	ASP	2.2
1	A	306	TYR	2.1
1	B	257	ILE	2.1
1	B	315	GLU	2.1
1	B	-3	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	EDO	B	413	4/4	0.49	0.32	91,91,92,92	0
5	EDO	B	419	4/4	0.54	0.45	90,91,91,92	0
5	EDO	B	405	4/4	0.61	0.40	89,90,90,90	0
5	EDO	B	406	4/4	0.74	0.38	66,66,66,67	0
5	EDO	A	409	4/4	0.75	0.26	93,93,94,94	0
6	PEG	A	408	7/7	0.75	0.24	81,82,84,84	0
5	EDO	A	406	4/4	0.77	0.27	85,85,86,86	0
5	EDO	A	413	4/4	0.77	0.33	94,94,94,94	0
5	EDO	B	409	4/4	0.77	0.37	99,99,99,100	0
5	EDO	B	414	4/4	0.78	0.68	87,88,88,88	0
5	EDO	B	416	4/4	0.79	0.41	96,96,96,96	0
6	PEG	A	410	7/7	0.79	0.36	82,82,83,83	0
5	EDO	B	408	4/4	0.80	0.33	97,97,97,97	0
6	PEG	A	412	7/7	0.80	0.41	79,79,79,79	7
5	EDO	B	404	4/4	0.81	0.36	82,82,82,82	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	EDO	B	411	4/4	0.81	0.19	92,92,92,93	0
5	EDO	A	411	4/4	0.81	0.17	90,90,90,90	0
5	EDO	B	407	4/4	0.82	0.42	83,83,83,83	0
6	PEG	B	418	7/7	0.82	0.28	90,90,90,90	0
5	EDO	D	101	4/4	0.85	0.21	86,86,87,87	0
5	EDO	B	415	4/4	0.87	0.45	86,86,86,87	0
5	EDO	B	410	4/4	0.88	0.20	86,86,86,86	0
5	EDO	A	407	4/4	0.88	0.33	81,81,81,81	0
5	EDO	A	404	4/4	0.89	0.64	64,64,65,66	0
5	EDO	B	412	4/4	0.89	0.70	100,100,101,101	0
5	EDO	A	405	4/4	0.90	0.34	78,78,78,78	0
5	EDO	B	417	4/4	0.91	0.23	78,78,78,78	0
3	SF4	B	401	8/8	0.96	0.14	51,52,52,55	8
4	SAH	B	403	26/26	0.98	0.12	54,55,57,58	0
3	SF4	A	401	8/8	0.98	0.18	48,49,50,53	0
3	SF4	B	402	8/8	0.98	0.16	56,56,57,57	8
4	SAH	A	403	26/26	0.98	0.13	48,48,51,51	0
3	SF4	A	402	8/8	0.99	0.17	53,53,54,55	8

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around SF4 B 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around SAH B 403:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around SF4 A 401:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




Electron density around SF4 B 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







6.5 Other polymers [i](#)

There are no such residues in this entry.