



Full wwPDB X-ray Structure Validation Report ⓘ

May 17, 2021 – 02:05 pm BST

PDB ID : 7AKE
Title : Structure of DYRK1A in complex with compound 58
Authors : Dokurno, P.; Surgenor, A.E.; Kotschy, A.
Deposited on : 2020-09-30
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.18
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.18


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	359	<div style="display: flex; align-items: center;"> <div style="width: 6%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 72%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 18%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 7%; height: 10px; background-color: grey; margin-right: 5px;"></div> </div> <p style="text-align: center;">6% 72% 18% • 7%</p>
1	B	359	<div style="display: flex; align-items: center;"> <div style="width: 5%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 66%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 24%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 7%; height: 10px; background-color: grey; margin-right: 5px;"></div> </div> <p style="text-align: center;">5% 66% 24% • 7%</p>

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5722 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dual specificity tyrosine-phosphorylation-regulated kinase 1A.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	P	S			
1	A	333	Total	C	N	O	P	S	0	0	0
			2702	1740	467	475	2	18			
1	B	335	Total	C	N	O	P	S	0	0	0
			2716	1746	468	482	2	18			

There are 42 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	127	MET	-	initiating methionine	UNP Q13627
A	128	GLY	-	expression tag	UNP Q13627
A	129	SER	-	expression tag	UNP Q13627
A	130	SER	-	expression tag	UNP Q13627
A	131	HIS	-	expression tag	UNP Q13627
A	132	HIS	-	expression tag	UNP Q13627
A	133	HIS	-	expression tag	UNP Q13627
A	134	HIS	-	expression tag	UNP Q13627
A	135	HIS	-	expression tag	UNP Q13627
A	136	HIS	-	expression tag	UNP Q13627
A	137	SER	-	expression tag	UNP Q13627
A	138	SER	-	expression tag	UNP Q13627
A	139	GLY	-	expression tag	UNP Q13627
A	140	LEU	-	expression tag	UNP Q13627
A	141	VAL	-	expression tag	UNP Q13627
A	142	PRO	-	expression tag	UNP Q13627
A	143	ARG	-	expression tag	UNP Q13627
A	144	GLY	-	expression tag	UNP Q13627
A	145	SEP	-	expression tag	UNP Q13627
A	146	HIS	-	expression tag	UNP Q13627
A	147	MET	-	expression tag	UNP Q13627
B	127	MET	-	initiating methionine	UNP Q13627
B	128	GLY	-	expression tag	UNP Q13627
B	129	SER	-	expression tag	UNP Q13627
B	130	SER	-	expression tag	UNP Q13627

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	131	HIS	-	expression tag	UNP Q13627
B	132	HIS	-	expression tag	UNP Q13627
B	133	HIS	-	expression tag	UNP Q13627
B	134	HIS	-	expression tag	UNP Q13627
B	135	HIS	-	expression tag	UNP Q13627
B	136	HIS	-	expression tag	UNP Q13627
B	137	SER	-	expression tag	UNP Q13627
B	138	SER	-	expression tag	UNP Q13627
B	139	GLY	-	expression tag	UNP Q13627
B	140	LEU	-	expression tag	UNP Q13627
B	141	VAL	-	expression tag	UNP Q13627
B	142	PRO	-	expression tag	UNP Q13627
B	143	ARG	-	expression tag	UNP Q13627
B	144	GLY	-	expression tag	UNP Q13627
B	145	SEP	-	expression tag	UNP Q13627
B	146	HIS	-	expression tag	UNP Q13627
B	147	MET	-	expression tag	UNP Q13627

- Molecule 2 is 4-[3-[(2 {S})-2-(6-bromanylpyridin-2-yl)oxypropyl]-2-methyl-imidazo[4,5-b]pyridin-5-yl]pyridine-2,6-diamine (three-letter code: RK8) (formula: C₂₀H₂₀BrN₇O) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	Br	C	N	O		
2	A	1	29	1	20	7	1	0	0
2	B	1	29	1	20	7	1	0	0

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total Cl 1 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	106	Total O 106 106	0	0
4	B	139	Total O 139 139	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Dual specificity tyrosine-phosphorylation-regulated kinase 1A



- Molecule 1: Dual specificity tyrosine-phosphorylation-regulated kinase 1A



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	58.25Å 84.95Å 84.91Å 90.00° 107.66° 90.00°	Depositor
Resolution (Å)	25.00 – 2.30 24.10 – 2.30	Depositor EDS
% Data completeness (in resolution range)	96.6 (25.00-2.30) 96.7 (24.10-2.30)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.92 (at 2.31Å)	Xtrriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.228 , 0.284 0.233 , 0.288	Depositor DCC
R_{free} test set	1714 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	33.2	Xtrriage
Anisotropy	0.128	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 32.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.029 for h,-k,-h-l	Xtrriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	5722	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.48% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PTR, RK8, SEP, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.64	0/2736	0.82	1/3685 (0.0%)
1	B	0.63	0/2751	0.82	0/3707
All	All	0.63	0/5487	0.82	1/7392 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	219	TYR	CB-CA-C	5.45	121.31	110.40

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	219	TYR	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2702	0	2718	53	0
1	B	2716	0	2720	92	1
2	A	29	0	0	1	0
2	B	29	0	0	2	0
3	B	1	0	0	1	0
4	A	106	0	0	7	0
4	B	139	0	0	6	0
All	All	5722	0	5438	143	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (143) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:150:LYS:CB	1:B:153:GLU:HG3	1.74	1.15
1:B:150:LYS:HB3	1:B:153:GLU:HG3	1.17	1.12
1:A:366:GLU:OE1	1:A:393:LYS:NZ	1.88	1.05
1:A:451:LYS:HD3	1:A:477:SER:OG	1.57	1.05
1:A:392:ARG:O	1:A:406:LYS:NZ	1.99	0.96
1:B:150:LYS:HB3	1:B:153:GLU:CG	1.96	0.95
1:B:148:ILE:H	1:B:148:ILE:HD12	1.45	0.82
1:B:396:GLU:HB2	4:B:707:HOH:O	1.81	0.80
1:A:246:TYR:CE1	1:B:143:ARG:HG2	2.16	0.80
1:B:297:ASN:HB2	1:B:298:PRO:HD2	1.65	0.79
1:A:156:MET:HE3	1:A:230:PHE:CD1	2.18	0.79
1:B:242:SER:HB2	1:B:297:ASN:HA	1.67	0.77
1:A:246:TYR:CZ	1:B:143:ARG:HG2	2.20	0.76
1:A:199:GLN:HG3	1:A:309:GLY:O	1.88	0.74
1:B:150:LYS:CB	1:B:153:GLU:CG	2.58	0.73
1:A:181:GLU:OE2	1:A:181:GLU:HA	1.91	0.71
1:B:392:ARG:NH2	1:B:397:LYS:H	1.88	0.70
1:A:321:PTR:CD2	1:B:194:LYS:HE2	2.21	0.70
1:A:451:LYS:CD	1:A:477:SER:OG	2.36	0.70
1:A:446:VAL:O	1:A:450:LEU:HG	1.93	0.68
1:A:400:ASP:OD1	1:A:400:ASP:N	2.17	0.68
1:B:151:ASN:N	1:B:151:ASN:OD1	2.27	0.67
1:B:344:MET:HE3	1:B:347:LEU:HB3	1.75	0.67
1:B:307:ASP:OD2	4:B:601:HOH:O	2.13	0.66
1:A:156:MET:CE	1:A:230:PHE:CD1	2.78	0.66

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:150:LYS:CA	1:B:153:GLU:HG3	2.26	0.66
1:B:334:LEU:O	1:B:390:LYS:HB2	1.96	0.64
1:A:393:LYS:O	1:A:406:LYS:HE2	1.96	0.64
1:B:344:MET:HE1	1:B:347:LEU:HD23	1.80	0.64
1:A:397:LYS:HE2	1:A:403:TRP:CZ2	2.33	0.63
1:B:392:ARG:O	1:B:406:LYS:HE3	1.99	0.62
1:A:475:GLN:NE2	4:A:602:HOH:O	2.33	0.61
1:B:158:ARG:HA	1:B:179:ARG:HD2	1.82	0.61
1:A:284:ILE:HG12	1:A:340:LEU:HD12	1.83	0.60
1:A:352:VAL:HG11	1:A:360:LEU:HD13	1.82	0.59
1:A:156:MET:HE3	1:A:230:PHE:CG	2.37	0.59
1:B:367:VAL:HG23	3:B:502:CL:CL	2.39	0.59
1:B:381:PRO:HG2	1:B:384:ILE:HG13	1.85	0.58
1:A:156:MET:HE3	1:A:230:PHE:HB2	1.85	0.58
1:B:150:LYS:H	1:B:153:GLU:CD	2.07	0.58
1:B:159:TYR:OH	1:B:226:ARG:HD2	2.03	0.58
1:B:294:LEU:HD21	2:B:501:RK8:O1	2.05	0.57
1:B:344:MET:CE	1:B:347:LEU:HD23	2.35	0.57
1:B:381:PRO:HG2	1:B:384:ILE:CG1	2.35	0.56
1:B:148:ILE:HD12	1:B:148:ILE:N	2.18	0.56
1:A:321:PTR:O2P	1:A:390:LYS:NZ	2.39	0.55
1:A:322:ILE:HD13	1:A:338:TYR:CZ	2.41	0.55
1:B:145:SEP:O1P	1:B:191:LYS:HE2	2.07	0.55
1:B:181:GLU:HA	1:B:181:GLU:OE1	2.07	0.55
1:B:392:ARG:HH22	1:B:397:LYS:H	1.55	0.55
1:B:145:SEP:O	1:B:233:HIS:HE1	1.89	0.54
1:B:398:LEU:HD22	1:B:399:PRO:HD2	1.88	0.54
1:B:218:LYS:HB2	1:B:275:PHE:CE2	2.42	0.54
1:B:397:LYS:HG3	1:B:403:TRP:CE2	2.42	0.54
1:B:344:MET:HE1	1:B:459:MET:CE	2.38	0.53
2:B:501:RK8:C1	4:B:721:HOH:O	2.56	0.53
1:B:334:LEU:C	1:B:390:LYS:HB2	2.29	0.53
1:A:219:TYR:HA	4:A:630:HOH:O	2.09	0.53
1:B:179:ARG:HD3	1:B:179:ARG:C	2.29	0.53
1:B:344:MET:CE	1:B:459:MET:SD	2.97	0.53
1:B:397:LYS:HG3	1:B:403:TRP:CZ2	2.44	0.52
1:B:230:PHE:CE1	1:B:231:ARG:HG3	2.44	0.52
1:A:197:LEU:O	1:A:201:GLN:HG2	2.09	0.52
1:B:218:LYS:HB2	1:B:275:PHE:CZ	2.44	0.52
1:A:145:SEP:O1P	1:A:193:LYS:NZ	2.43	0.52
1:A:218:LYS:HD3	1:A:223:HIS:HB2	1.92	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:156:MET:HE3	1:A:230:PHE:CB	2.41	0.51
1:B:155:TRP:HH2	1:B:189:ILE:HD11	1.75	0.51
1:B:148:ILE:H	1:B:148:ILE:CD1	2.21	0.51
1:B:429:VAL:HG22	1:B:449:TYR:HB3	1.93	0.51
1:B:398:LEU:HB2	1:B:402:THR:O	2.11	0.50
1:B:407:LYS:H	1:B:407:LYS:CE	2.24	0.50
1:A:243:TYR:CE1	1:A:295:LEU:HD12	2.47	0.50
1:B:284:ILE:HG12	1:B:340:LEU:HD12	1.92	0.49
1:B:417:PRO:HB2	1:B:420:THR:HG21	1.93	0.49
1:A:421:ARG:HG2	1:A:426:ILE:HD11	1.94	0.49
1:B:406:LYS:HB3	4:B:707:HOH:O	2.12	0.49
1:B:374:VAL:HG11	1:B:405:LEU:HD11	1.94	0.49
1:A:386:ASP:OD1	1:A:397:LYS:NZ	2.40	0.49
1:A:154:LYS:HA	1:A:159:TYR:O	2.11	0.49
1:B:155:TRP:HH2	1:B:189:ILE:CD1	2.26	0.49
1:B:438:ARG:HG3	1:B:444:HIS:CD2	2.48	0.48
1:A:393:LYS:C	1:A:406:LYS:HE2	2.32	0.48
1:B:344:MET:HE2	1:B:459:MET:SD	2.53	0.48
1:B:392:ARG:NH2	1:B:397:LYS:N	2.60	0.48
1:B:222:VAL:HB	1:B:306:VAL:HG12	1.95	0.48
1:B:396:GLU:CB	4:B:707:HOH:O	2.48	0.47
1:B:154:LYS:CG	1:B:160:GLU:HG3	2.44	0.47
1:B:388:ALA:HB3	1:B:391:ALA:HB2	1.96	0.47
1:A:193:LYS:HG3	4:A:666:HOH:O	2.14	0.47
1:A:257:VAL:HG22	1:A:354:MET:O	2.14	0.47
1:B:164:LEU:HD21	1:B:172:GLN:HB3	1.96	0.47
1:B:206:LEU:HD22	1:B:283:ILE:HG12	1.97	0.46
1:B:213:HIS:O	1:B:218:LYS:HE3	2.15	0.46
1:B:407:LYS:N	1:B:407:LYS:CD	2.78	0.46
1:B:179:ARG:C	1:B:179:ARG:CD	2.84	0.46
1:B:344:MET:HE1	1:B:459:MET:SD	2.55	0.46
1:B:196:PHE:HD1	1:B:199:GLN:NE2	2.13	0.46
1:A:181:GLU:HB3	1:A:183:GLU:HG2	1.98	0.45
1:A:322:ILE:CD1	1:A:338:TYR:CZ	2.99	0.45
1:B:179:ARG:HH11	1:B:179:ARG:HG2	1.82	0.45
1:B:261:LEU:HD11	1:B:265:PHE:CZ	2.52	0.45
1:B:373:ILE:HG23	1:B:377:LEU:HD12	1.99	0.45
1:B:196:PHE:CD1	1:B:199:GLN:NE2	2.85	0.45
1:B:345:TRP:C	1:B:345:TRP:CD1	2.90	0.45
1:B:395:PHE:CE1	1:B:405:LEU:HD23	2.52	0.44
1:A:334:LEU:HB3	1:A:388:ALA:HB1	2.00	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:218:LYS:HD3	1:B:219:TYR:CE1	2.53	0.44
1:B:395:PHE:CD1	1:B:405:LEU:HD23	2.53	0.44
1:A:262:THR:HA	4:A:698:HOH:O	2.18	0.44
1:A:392:ARG:C	1:A:406:LYS:NZ	2.69	0.44
1:B:425:ASN:ND2	4:B:616:HOH:O	2.51	0.43
1:A:226:ARG:NH1	4:A:603:HOH:O	2.50	0.43
1:A:397:LYS:HE2	1:A:403:TRP:CE2	2.53	0.43
1:B:154:LYS:HG3	1:B:160:GLU:HA	2.01	0.43
1:A:404:ASN:HB2	4:A:661:HOH:O	2.17	0.43
1:B:179:ARG:HH11	1:B:179:ARG:CG	2.32	0.43
1:B:407:LYS:H	1:B:407:LYS:HE2	1.83	0.43
1:B:164:LEU:HD13	1:B:166:GLY:H	1.83	0.43
1:A:275:PHE:O	1:A:278:THR:HB	2.19	0.43
1:A:451:LYS:NZ	4:A:601:HOH:O	2.52	0.43
1:B:242:SER:OG	1:B:298:PRO:HD3	2.18	0.43
1:B:168:GLY:HA3	1:B:170:PHE:CE1	2.53	0.43
1:B:256:GLY:HA3	1:B:355:HIS:O	2.19	0.43
1:B:158:ARG:HG3	1:B:159:TYR:CD2	2.55	0.42
1:A:363:GLY:HA2	1:A:368:ASP:OD2	2.19	0.42
1:B:417:PRO:HB2	1:B:420:THR:CG2	2.49	0.42
1:B:334:LEU:HB3	1:B:388:ALA:HB1	2.02	0.42
1:A:345:TRP:CD1	1:A:345:TRP:C	2.93	0.42
1:A:275:PHE:O	1:A:281:LEU:HD12	2.20	0.42
1:A:435:GLY:HA3	1:A:437:ARG:NH1	2.35	0.42
1:B:154:LYS:HD2	1:B:160:GLU:CG	2.50	0.42
1:A:335:GLY:HA3	1:A:390:LYS:HD2	2.02	0.42
1:B:143:ARG:HD2	1:B:143:ARG:HA	1.51	0.42
1:B:288:LEU:HD23	1:B:288:LEU:HA	1.85	0.41
1:A:294:LEU:HD21	2:A:501:RK8:C2	2.50	0.41
1:B:277:ALA:HA	1:B:340:LEU:CD2	2.51	0.41
1:A:158:ARG:HG3	1:A:180:VAL:HG23	2.03	0.41
1:B:366:GLU:H	1:B:366:GLU:CD	2.24	0.41
1:A:156:MET:HE2	1:A:230:PHE:CD1	2.56	0.41
1:A:156:MET:CE	1:A:230:PHE:HD1	2.29	0.40
1:B:178:ASP:CG	1:B:181:GLU:HB2	2.41	0.40
1:B:269:MET:SD	1:B:293:ILE:HD13	2.62	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:182:GLN:O	1:B:406:LYS:NZ[1_655]	1.95	0.25

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	325/359 (90%)	309 (95%)	14 (4%)	2 (1%)	25	31
1	B	329/359 (92%)	316 (96%)	13 (4%)	0	100	100
All	All	654/718 (91%)	625 (96%)	27 (4%)	2 (0%)	41	50

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	220	TYR
1	A	323	GLN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	289/315 (92%)	271 (94%)	18 (6%)	18	25
1	B	290/315 (92%)	265 (91%)	25 (9%)	10	12
All	All	579/630 (92%)	536 (93%)	43 (7%)	13	17

All (43) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	151	ASN
1	A	153	GLU
1	A	154	LYS
1	A	169	SER
1	A	181	GLU
1	A	182	GLN
1	A	193	LYS
1	A	226	ARG
1	A	239	GLU
1	A	278	THR
1	A	310	SER
1	A	324	SER
1	A	358	GLU
1	A	394	PHE
1	A	400	ASP
1	A	408	THR
1	A	454	ASP
1	A	465	LYS
1	B	143	ARG
1	B	146	HIS
1	B	149	VAL
1	B	150	LYS
1	B	151	ASN
1	B	154	LYS
1	B	157	ASP
1	B	172	GLN
1	B	179	ARG
1	B	197	LEU
1	B	211	ASN
1	B	212	LYS
1	B	214	ASP
1	B	215	THR
1	B	218	LYS
1	B	226	ARG
1	B	239	GLU
1	B	300	ARG
1	B	310	SER
1	B	392	ARG
1	B	397	LYS
1	B	398	LEU
1	B	400	ASP
1	B	407	LYS
1	B	408	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	213	HIS
1	A	251	ASN
1	A	320	GLN
1	A	424	HIS
1	A	425	ASN
1	B	182	GLN
1	B	199	GLN
1	B	201	GLN
1	B	251	ASN
1	B	320	GLN
1	B	369	GLN
1	B	425	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	SEP	A	145	1	8,9,10	0.59	0	8,12,14	0.86	0
1	SEP	B	145	1	8,9,10	0.61	0	8,12,14	0.69	0
1	PTR	B	321	1	15,16,17	0.56	0	19,22,24	1.14	2 (10%)
1	PTR	A	321	1	15,16,17	0.51	0	19,22,24	0.93	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	SEP	A	145	1	-	1/5/8/10	-
1	SEP	B	145	1	-	1/5/8/10	-
1	PTR	B	321	1	-	0/10/11/13	0/1/1/1
1	PTR	A	321	1	-	0/10/11/13	0/1/1/1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	B	321	PTR	O3P-P-OH	2.82	114.06	105.24
1	A	321	PTR	O2P-P-OH	2.63	113.48	105.24
1	B	321	PTR	OH-CZ-CE1	2.11	125.52	119.23

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	145	SEP	CA-CB-OG-P
1	B	145	SEP	CA-CB-OG-P

There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	145	SEP	1	0
1	B	145	SEP	2	0
1	A	321	PTR	2	0

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	RK8	A	501	-	30,32,32	0.90	1 (3%)	33,46,46	3.42	8 (24%)
2	RK8	B	501	-	30,32,32	0.91	1 (3%)	33,46,46	2.31	8 (24%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	RK8	A	501	-	-	4/10/12/12	0/4/4/4
2	RK8	B	501	-	-	1/10/12/12	0/4/4/4

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	RK8	C3-N1	-2.89	1.45	1.48
2	B	501	RK8	C3-N1	-2.39	1.45	1.48

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	RK8	O1-C16-N7	-14.61	108.56	119.45
2	A	501	RK8	O1-C16-C17	10.59	129.06	115.89
2	B	501	RK8	O1-C16-N7	9.80	126.75	119.45
2	B	501	RK8	C1-C2-C3	4.09	117.86	111.36
2	B	501	RK8	O1-C16-C17	-3.86	111.08	115.89
2	A	501	RK8	BR1-C20-C19	3.31	122.03	118.64
2	A	501	RK8	C11-C15-C14	3.08	119.84	117.38
2	B	501	RK8	C11-C15-C14	2.95	119.74	117.38
2	B	501	RK8	C11-C12-C13	2.93	119.72	117.38
2	B	501	RK8	C16-O1-C2	2.91	122.48	118.19
2	B	501	RK8	O1-C2-C3	-2.78	102.80	106.08
2	A	501	RK8	BR1-C20-N7	-2.49	113.51	115.71
2	A	501	RK8	C8-C7-C6	-2.49	117.71	120.84
2	A	501	RK8	O1-C2-C3	2.39	108.90	106.08
2	B	501	RK8	C8-C7-C6	-2.25	118.01	120.84
2	A	501	RK8	C11-C12-C13	2.25	119.18	117.38

There are no chirality outliers.

All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	501	RK8	N7-C16-O1-C2
2	A	501	RK8	C17-C16-O1-C2
2	B	501	RK8	C1-C2-O1-C16
2	A	501	RK8	O1-C2-C3-N1
2	A	501	RK8	C1-C2-C3-N1


There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	RK8	1	0
2	B	501	RK8	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	331/359 (92%)	0.45	20 (6%) 21 28	20, 34, 58, 85	2 (0%)
1	B	333/359 (92%)	0.41	19 (5%) 23 30	19, 31, 71, 97	2 (0%)
All	All	664/718 (92%)	0.43	39 (5%) 22 28	19, 33, 65, 97	4 (0%)

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	214	ASP	4.8
1	B	399	PRO	4.5
1	B	398	LEU	4.4
1	B	401	GLY	4.1
1	A	148	ILE	3.8
1	A	439	ALA	3.8
1	B	219	TYR	3.4
1	A	351	LEU	3.1
1	A	407	LYS	3.1
1	A	217	MET	3.1
1	A	408	THR	3.0
1	A	477	SER	2.9
1	B	400	ASP	2.9
1	A	252	THR	2.8
1	A	347	LEU	2.7
1	B	182	GLN	2.7
1	B	215	THR	2.7
1	B	217	MET	2.6
1	B	144	GLY	2.5
1	A	219	TYR	2.5
1	A	399	PRO	2.5
1	A	144	GLY	2.5
1	B	440	GLY	2.5
1	A	254	PHE	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	177	TYR	2.3
1	B	154	LYS	2.3
1	A	440	GLY	2.3
1	B	402	THR	2.3
1	A	401	GLY	2.3
1	B	481	LYS	2.2
1	A	454	ASP	2.2
1	B	221	ILE	2.2
1	A	466	THR	2.1
1	B	142	PRO	2.1
1	A	450	LEU	2.1
1	A	428	GLY	2.1
1	B	213	HIS	2.0
1	B	157	ASP	2.0
1	A	300	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	SEP	B	145	10/11	0.87	0.15	55,59,67,68	5
1	PTR	A	321	16/17	0.89	0.14	36,41,60,61	0
1	PTR	B	321	16/17	0.90	0.15	35,39,55,57	0
1	SEP	A	145	10/11	0.93	0.19	40,47,58,66	0

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	RK8	A	501	29/29	0.87	0.17	25,31,52,68	1
2	RK8	B	501	29/29	0.91	0.15	20,22,44,63	1
3	CL	B	502	1/1	0.97	0.06	54,54,54,54	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.