



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 23, 2023 – 08:34 PM EDT

PDB ID : 3AL7
Title : Recombinant thaumatin I at 1.1 Å
Authors : Masuda, T.; Mikami, B.; Kitabatake, N.
Deposited on : 2010-07-27
Resolution : 1.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1619 (1.14-1.06)
Clashscore	141614	1671 (1.14-1.06)
Ramachandran outliers	138981	1615 (1.14-1.06)
Sidechain outliers	138945	1613 (1.14-1.06)
RSRZ outliers	127900	1588 (1.14-1.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	207	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	230	-	X	-	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	260	-	X	-	-

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 3928 atoms, of which 1633 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Thaumatin I.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	207	3386	1079	1633	319	336	19	0	21	0

- Molecule 2 is L(+)-TARTARIC ACID (three-letter code: TLA) (formula: C₄H₆O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	10	4	6	0	0
2	A	1	10	4	6	0	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	476	Total O 498 498	0	22

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Thaumatin I



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	57.70Å 57.70Å 149.78Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 1.10 18.11 – 1.10	Depositor EDS
% Data completeness (in resolution range)	94.0 (10.00-1.10) 98.6 (18.11-1.10)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.11 (at 1.10Å)	Xtrriage
Refinement program	SHELXL-97	Depositor
R, R_{free}	0.108 , 0.135 0.107 , 0.105	Depositor DCC
R_{free} test set	5113 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	8.3	Xtrriage
Anisotropy	0.265	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 87.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.99	EDS
Total number of atoms	3928	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.45% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, TLA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.87	0/1794	1.58	37/2422 (1.5%)

There are no bond length outliers.

All (37) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	76[A]	ARG	NE-CZ-NH2	12.81	126.70	120.30
1	A	76[B]	ARG	NE-CZ-NH2	12.81	126.70	120.30
1	A	76[C]	ARG	NE-CZ-NH2	12.81	126.70	120.30
1	A	119[A]	ARG	NE-CZ-NH1	11.70	126.15	120.30
1	A	119[B]	ARG	NE-CZ-NH1	11.70	126.15	120.30
1	A	79[A]	ARG	NE-CZ-NH1	-11.48	114.56	120.30
1	A	79[A]	ARG	NE-CZ-NH2	11.48	126.04	120.30
1	A	79[B]	ARG	NE-CZ-NH1	-11.48	114.56	120.30
1	A	79[B]	ARG	NE-CZ-NH2	11.48	126.04	120.30
1	A	163[A]	LYS	O-C-N	-11.29	104.64	122.70
1	A	163[B]	LYS	O-C-N	-11.29	104.64	122.70
1	A	122[A]	ARG	CD-NE-CZ	9.05	136.27	123.60
1	A	122[B]	ARG	CD-NE-CZ	9.05	136.27	123.60
1	A	76[A]	ARG	NE-CZ-NH1	-8.94	115.83	120.30
1	A	76[B]	ARG	NE-CZ-NH1	-8.94	115.83	120.30
1	A	76[C]	ARG	NE-CZ-NH1	-8.94	115.83	120.30
1	A	55[A]	ASP	CB-CG-OD2	8.14	125.62	118.30
1	A	55[B]	ASP	CB-CG-OD2	8.14	125.62	118.30
1	A	163[A]	LYS	CA-C-O	7.95	136.80	120.10
1	A	163[B]	LYS	CA-C-O	7.95	136.80	120.10
1	A	70[A]	ASP	CB-CG-OD1	7.61	125.15	118.30
1	A	70[B]	ASP	CB-CG-OD1	7.61	125.15	118.30
1	A	95	TYR	CB-CG-CD1	7.31	125.39	121.00
1	A	57[A]	TYR	CZ-CE2-CD2	-7.28	113.25	119.80

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	57[B]	TYR	CZ-CE2-CD2	-7.28	113.25	119.80
1	A	119[A]	ARG	NH1-CZ-NH2	-6.88	111.83	119.40
1	A	119[B]	ARG	NH1-CZ-NH2	-6.88	111.83	119.40
1	A	70[A]	ASP	CB-CG-OD2	-6.54	112.42	118.30
1	A	70[B]	ASP	CB-CG-OD2	-6.54	112.42	118.30
1	A	175	ARG	NE-CZ-NH1	6.38	123.49	120.30
1	A	200[A]	ARG	NE-CZ-NH1	6.18	123.39	120.30
1	A	200[B]	ARG	NE-CZ-NH1	6.18	123.39	120.30
1	A	55[A]	ASP	CB-CG-OD1	-5.25	113.57	118.30
1	A	55[B]	ASP	CB-CG-OD1	-5.25	113.57	118.30
1	A	122[A]	ARG	NE-CZ-NH1	-5.19	117.70	120.30
1	A	122[B]	ARG	NE-CZ-NH1	-5.19	117.70	120.30
1	A	162	GLY	O-C-N	-5.05	114.63	122.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1753	1633	1656	26	2
2	A	20	0	8	2	0
3	A	24	0	32	6	0
4	A	498	0	0	19	6
All	All	2295	1633	1696	32	6

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:207:ALA:CB	4:A:651:HOH:O	2.14	0.93
1:A:207:ALA:HB3	4:A:651:HOH:O	1.66	0.92
1:A:139:LYS:HE3	4:A:600:HOH:O	1.77	0.85

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:76[C]:ARG:NH2	4:A:628:HOH:O	2.09	0.84
3:A:250:GOL:H32	4:A:619:HOH:O	1.84	0.77
1:A:168[A]:GLU:HG3	4:A:656:HOH:O	1.84	0.76
1:A:168[B]:GLU:HG3	4:A:656:HOH:O	1.87	0.75
1:A:20:GLY:O	1:A:79[B]:ARG:HD2	1.95	0.66
3:A:260:GOL:H12	4:A:367:HOH:O	1.98	0.64
1:A:207:ALA:HB2	4:A:651:HOH:O	1.88	0.62
1:A:26:ALA:O	1:A:76[B]:ARG:HB3	2.00	0.62
1:A:79[B]:ARG:NH1	4:A:599:HOH:O	2.34	0.53
1:A:179[B]:ASP:OD1	4:A:666:HOH:O	2.18	0.53
3:A:240:GOL:C1	4:A:743:HOH:O	2.57	0.52
1:A:91:SER:OG	3:A:230:GOL:H32	2.09	0.52
2:A:210:TLA:O4	2:A:210:TLA:O2	2.27	0.52
1:A:206:THR:O	1:A:207:ALA:O	2.30	0.48
1:A:124[A]:VAL:HG12	1:A:179[A]:ASP:OD1	2.14	0.48
1:A:119[A]:ARG:NE	1:A:119[A]:ARG:H	2.13	0.47
3:A:240:GOL:H12	4:A:743:HOH:O	2.15	0.47
1:A:163[A]:LYS:HE3	4:A:682:HOH:O	2.14	0.47
1:A:8[B]:ARG:NH2	1:A:205:PRO:O	2.48	0.46
1:A:122[A]:ARG:NH1	4:A:750:HOH:O	2.48	0.46
1:A:163[A]:LYS:C	4:A:657:HOH:O	2.55	0.45
1:A:76[B]:ARG:NH1	4:A:535:HOH:O	2.48	0.45
1:A:159[A]:CYS:SG	1:A:184:VAL:HB	2.57	0.45
1:A:119[B]:ARG:NH1	2:A:210:TLA:O3	2.49	0.44
1:A:57[A]:TYR:O	1:A:64:GLY:HA3	2.17	0.44
1:A:119[A]:ARG:H	1:A:119[A]:ARG:CD	2.31	0.44
1:A:168[A]:GLU:CG	4:A:656:HOH:O	2.56	0.41
3:A:240:GOL:C3	4:A:743:HOH:O	2.68	0.40
1:A:119[B]:ARG:HG2	1:A:120:GLY:CA	2.51	0.40

All (6) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:57[B]:TYR:OH	4:A:761:HOH:O[8_665]	0.95	1.25
4:A:759:HOH:O	4:A:759:HOH:O[8_665]	1.53	0.67
4:A:444:HOH:O	4:A:596:HOH:O[5_545]	1.96	0.24
4:A:617:HOH:O	4:A:618:HOH:O[8_655]	2.00	0.20
4:A:596:HOH:O	4:A:626:HOH:O[5_555]	2.02	0.18
1:A:57[B]:TYR:CZ	4:A:761:HOH:O[8_665]	2.19	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	227/207 (110%)	224 (99%)	3 (1%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	189/167 (113%)	184 (97%)	5 (3%)	46 9

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	119[A]	ARG
1	A	119[B]	ARG
1	A	163[A]	LYS
1	A	163[B]	LYS
1	A	176	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	TLA	A	210	-	9,9,9	1.65	2 (22%)	12,12,12	2.30	6 (50%)
2	TLA	A	220	-	9,9,9	1.44	1 (11%)	12,12,12	0.78	0
3	GOL	A	230	-	5,5,5	0.78	0	5,5,5	3.04	4 (80%)
3	GOL	A	240	-	5,5,5	0.72	0	5,5,5	1.18	0
3	GOL	A	250	-	5,5,5	0.42	0	5,5,5	0.62	0
3	GOL	A	260	-	5,5,5	1.64	1 (20%)	5,5,5	2.92	2 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TLA	A	210	-	-	1/12/12/12	-
2	TLA	A	220	-	-	0/12/12/12	-
3	GOL	A	230	-	-	2/4/4/4	-
3	GOL	A	240	-	-	4/4/4/4	-
3	GOL	A	250	-	-	0/4/4/4	-
3	GOL	A	260	-	-	4/4/4/4	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	210	TLA	C3-C4	4.08	1.58	1.52
2	A	220	TLA	C2-C1	-3.10	1.48	1.52
3	A	260	GOL	O2-C2	3.02	1.52	1.43
2	A	210	TLA	O2-C2	2.26	1.46	1.42

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	260	GOL	O3-C3-C2	4.86	133.51	110.20
2	A	210	TLA	O3-C3-C4	-3.80	102.69	110.66
3	A	230	GOL	O2-C2-C3	-3.70	92.81	109.12
2	A	210	TLA	O2-C2-C3	-3.66	102.97	110.23
3	A	260	GOL	O1-C1-C2	3.63	127.59	110.20
3	A	230	GOL	C3-C2-C1	3.45	125.11	111.70
3	A	230	GOL	O2-C2-C1	-3.35	94.36	109.12
2	A	210	TLA	O11-C1-O1	2.99	130.87	124.09
2	A	210	TLA	C2-C3-C4	-2.88	103.45	109.87
2	A	210	TLA	O4-C4-C3	-2.81	114.24	121.63
3	A	230	GOL	O1-C1-C2	-2.72	97.17	110.20
2	A	210	TLA	O41-C4-C3	-2.50	106.53	113.27

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	230	GOL	O2-C2-C3-O3
3	A	240	GOL	O1-C1-C2-C3
3	A	240	GOL	C1-C2-C3-O3
3	A	260	GOL	O1-C1-C2-C3
3	A	260	GOL	C1-C2-C3-O3
2	A	210	TLA	O3-C3-C4-O4
3	A	240	GOL	O1-C1-C2-O2
3	A	240	GOL	O2-C2-C3-O3
3	A	260	GOL	O1-C1-C2-O2
3	A	260	GOL	O2-C2-C3-O3
3	A	230	GOL	C1-C2-C3-O3

There are no ring outliers.

5 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	210	TLA	2	0
3	A	230	GOL	1	0
3	A	240	GOL	3	0
3	A	250	GOL	1	0
3	A	260	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	207/207 (100%)	-0.55	4 (1%) 66 62	6, 9, 22, 76	4 (1%)

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	207	ALA	10.3
1	A	119[A]	ARG	3.6
1	A	57[A]	TYR	2.6
1	A	162	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GOL	A	250	6/6	0.70	0.26	44,81,87,93	0
3	GOL	A	230	6/6	0.82	0.17	29,42,51,63	0
3	GOL	A	240	6/6	0.86	0.17	28,35,53,54	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	TLA	A	210	10/10	0.88	0.23	21,25,35,59	0
3	GOL	A	260	6/6	0.94	0.18	8,20,25,32	0
2	TLA	A	220	10/10	1.00	0.03	6,7,8,9	0

6.5 Other polymers [i](#)

There are no such residues in this entry.