



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 17, 2023 – 01:05 am GMT

PDB ID : 4AMO
Title : Crystal Structure of the Acyltransferase Domain of the Iterative Polyketide Synthase in Eneidyne Biosynthesis Reveals the Molecular Basis of Substrate Specificity
Authors : Liew, C.W.; Lescar, J.
Deposited on : 2012-03-13
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	421	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	ACY	A	1877	-	-	X	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	ACY	A	1878	-	-	X	-

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2875 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called DYNE8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	371	2677	1672	509	490	6	0	6	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	682	VAL	ALA	conflict	UNP Q84HI8

- Molecule 2 is ACETIC ACID (three-letter code: ACY) (formula: C₂H₄O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	4	2	2	0	0
2	A	1	4	2	2	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	190	Total 190	O 190	0	0

4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	66.05Å 66.40Å 85.99Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.66 – 1.90 29.66 – 1.90	Depositor EDS
% Data completeness (in resolution range)	100.0 (29.66-1.90) 100.0 (29.66-1.90)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.08 (at 1.91Å)	Xtrriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.178 , 0.218 0.178 , 0.219	Depositor DCC
R_{free} test set	1538 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	26.0	Xtrriage
Anisotropy	0.084	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 49.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.025 for k,h,-l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2875	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.78% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: ACY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.11	7/2742 (0.3%)	1.10	18/3738 (0.5%)

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	715	ASN	C-N	-15.25	1.05	1.33
1	A	650	HIS	C-N	-7.90	1.15	1.34
1	A	714	ASP	C-N	-7.13	1.17	1.34
1	A	660	TRP	CB-CG	6.16	1.61	1.50
1	A	483	CYS	CB-SG	-5.32	1.73	1.81
1	A	834	GLU	CG-CD	5.08	1.59	1.51
1	A	861	PHE	CG-CD2	5.03	1.46	1.38

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	757	MET	CG-SD-CE	-13.24	79.01	100.20
1	A	715	ASN	O-C-N	-12.11	102.62	123.20
1	A	650	HIS	O-C-N	-9.97	106.75	122.70
1	A	715	ASN	CA-C-N	9.72	135.64	116.20
1	A	800	ARG	NE-CZ-NH2	-8.72	115.94	120.30
1	A	528[A]	ARG	NE-CZ-NH2	-8.48	116.06	120.30
1	A	528[B]	ARG	NE-CZ-NH2	-8.48	116.06	120.30
1	A	800	ARG	NE-CZ-NH1	7.49	124.05	120.30
1	A	696	ASP	CB-CG-OD1	6.89	124.50	118.30
1	A	648	LEU	CA-CB-CG	6.83	131.01	115.30
1	A	522	ARG	NE-CZ-NH1	6.83	123.72	120.30
1	A	701	ARG	NE-CZ-NH2	-6.72	116.94	120.30
1	A	650	HIS	CA-C-N	6.69	131.92	117.20
1	A	701	ARG	NE-CZ-NH1	5.71	123.15	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	791	ASP	CB-CG-OD1	5.64	123.38	118.30
1	A	819	LEU	CA-CB-CG	5.56	128.09	115.30
1	A	814	LEU	CB-CG-CD1	-5.53	101.60	111.00
1	A	838	ARG	NE-CZ-NH2	-5.27	117.67	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2677	0	2720	41	0
2	A	8	0	6	11	0
3	A	190	0	0	7	0
All	All	2875	0	2726	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:651[A]:SER:OG	2:A:1877:ACY:H3	1.26	1.31
1:A:639[A]:ARG:HH11	1:A:639[A]:ARG:CG	1.62	1.08
1:A:639[A]:ARG:HG3	1:A:639[A]:ARG:NH1	1.60	1.08
1:A:718[B]:ARG:HG2	3:A:2121:HOH:O	1.52	1.07
1:A:651[A]:SER:OG	2:A:1877:ACY:CH3	2.14	0.95
1:A:581:GLN:H	2:A:1877:ACY:H1	1.33	0.90
1:A:580:GLY:H	2:A:1878:ACY:H1	1.37	0.90
1:A:639[A]:ARG:HH11	1:A:639[A]:ARG:HG3	0.76	0.89
1:A:581:GLN:HG2	1:A:582:ALA:H	1.37	0.89
1:A:549:HIS:HE1	1:A:556:ARG:H	1.29	0.80
1:A:581:GLN:H	2:A:1877:ACY:CH3	1.95	0.80
1:A:581:GLN:O	1:A:582:ALA:HB3	1.81	0.79
1:A:581:GLN:HG2	1:A:582:ALA:N	2.01	0.76

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:639[A]:ARG:CG	1:A:639[A]:ARG:NH1	2.35	0.70
1:A:834:GLU:OE1	3:A:2177:HOH:O	2.09	0.70
1:A:581:GLN:O	1:A:582:ALA:CB	2.38	0.70
1:A:651[B]:SER:HB3	2:A:1877:ACY:H3	1.77	0.66
1:A:528[B]:ARG:HH22	1:A:570:GLY:HA3	1.59	0.66
1:A:580:GLY:H	2:A:1878:ACY:CH3	2.13	0.59
1:A:484:GLY:HA3	1:A:525:VAL:CG1	2.33	0.58
1:A:581:GLN:C	1:A:583:ALA:H	2.06	0.57
1:A:528[B]:ARG:NH1	3:A:2025:HOH:O	2.38	0.57
1:A:581:GLN:CG	1:A:582:ALA:H	2.16	0.55
1:A:870:THR:N	1:A:871:PRO:CD	2.72	0.52
1:A:580:GLY:N	2:A:1878:ACY:H1	2.15	0.52
1:A:555:ARG:HD3	1:A:568:VAL:HB	1.93	0.51
1:A:581:GLN:N	2:A:1877:ACY:H1	2.14	0.49
1:A:651[B]:SER:OG	1:A:753[B]:HIS:HE1	1.95	0.49
1:A:577:LEU:HB3	1:A:829:LEU:HD23	1.95	0.49
1:A:549:HIS:HE1	1:A:556:ARG:N	2.05	0.48
1:A:549:HIS:HD2	3:A:2033:HOH:O	1.96	0.47
1:A:829:LEU:HD13	2:A:1878:ACY:H2	1.96	0.47
1:A:581:GLN:C	1:A:583:ALA:N	2.70	0.46
1:A:829:LEU:CD1	2:A:1878:ACY:H2	2.47	0.45
1:A:521:HIS:HD2	3:A:2005:HOH:O	1.99	0.45
1:A:528[B]:ARG:CZ	3:A:2025:HOH:O	2.65	0.44
1:A:737:ARG:HG3	3:A:2132:HOH:O	2.18	0.44
1:A:651[A]:SER:OG	1:A:652:LEU:N	2.50	0.44
1:A:723:ALA:HB1	1:A:751:ALA:HB1	2.01	0.43
1:A:665:ASP:OD1	1:A:668:ASP:OD2	2.39	0.41
1:A:581:GLN:CG	1:A:582:ALA:N	2.74	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	373/421 (89%)	361 (97%)	12 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	249/282 (88%)	240 (96%)	9 (4%)	35	26

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	503	THR
1	A	505	LEU
1	A	617	VAL
1	A	639[A]	ARG
1	A	639[B]	ARG
1	A	648	LEU
1	A	762	GLU
1	A	802	LEU
1	A	829	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	521	HIS
1	A	549	HIS
1	A	581	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	ACY	A	1877	-	3,3,3	1.24	0	3,3,3	0.86	0
2	ACY	A	1878	-	3,3,3	1.17	0	3,3,3	0.84	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1877	ACY	6	0
2	A	1878	ACY	5	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	3

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	714:ASP	C	715:ASN	N	1.17
1	A	650:HIS	C	651:SER	N	1.15
1	A	715:ASN	C	716:GLY	N	1.05

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	371/421 (88%)	0.13	22 (5%) 22 25	15, 25, 46, 56	0

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	504	ALA	5.2
1	A	848	ALA	4.6
1	A	616	PRO	3.9
1	A	584	PRO	3.9
1	A	617	VAL	3.8
1	A	874	ALA	3.6
1	A	585	VAL	3.5
1	A	791	ASP	3.2
1	A	849	SER	3.2
1	A	583	ALA	3.2
1	A	582	ALA	2.9
1	A	762	GLU	2.8
1	A	775	ARG	2.7
1	A	846	GLY	2.6
1	A	875	GLY	2.4
1	A	682	VAL	2.4
1	A	685	GLU	2.3
1	A	847	ALA	2.3
1	A	749	SER	2.2
1	A	784	VAL	2.2
1	A	790	ALA	2.1
1	A	667	ASP	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	ACY	A	1878	4/4	0.34	0.36	57,59,62,63	0
2	ACY	A	1877	4/4	0.74	0.27	41,45,48,52	0

6.5 Other polymers [i](#)

There are no such residues in this entry.