Full wwPDB X-ray Structure Validation Report (i)

Sep 25, 2023 – 11:43 PM EDT

PDB ID	:	6APS
Title	:	Trypanosoma brucei hypoxanthine guanine phosphoribosyltransferase in complex with [(2-((Guanine-9H-yl)methyl)propane-1,3 diyl)bis(oxy)]bis(methylene))diphosphonic acid
Authors	:	Teran, D.; Guddat, L.W.
Deposited on	:	2017-08-17
Resolution	:	1.76 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.35.1
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.76 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2340 (1.76-1.76)
Clashscore	141614	2466 (1.76-1.76)
Ramachandran outliers	138981	2437 (1.76-1.76)
Sidechain outliers	138945	2437 (1.76-1.76)
RSRZ outliers	127900	2298 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	216	3% 73% 13% 14%
1	B	216	3% 75% 7% 17%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	PEG	A	303	-	-	X	-
4	PEG	B	303	-	-	X	-
5	TRS	A	304	-	X	-	-

2 Entry composition (i)

There are 7 unique types of molecules in this entry. The entry contains 3435 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hypoxanthine-guanine phosphoribosyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	185	1508	965	257	279	7	0	5	0
1	B	179	1456	936	246	270	4	0	7	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	HIS	-	expression tag	UNP Q07010
A	-4	HIS	-	expression tag	UNP Q07010
A	-3	HIS	-	expression tag	UNP Q07010
A	-2	HIS	-	expression tag	UNP Q07010
A	-1	HIS	-	expression tag	UNP Q07010
A	0	HIS	-	expression tag	UNP Q07010
B	-5	HIS	-	expression tag	UNP Q07010
B	-4	HIS	-	expression tag	UNP Q07010
B	-3	HIS	-	expression tag	UNP Q07010
B	-2	HIS	-	expression tag	UNP Q07010
B	-1	HIS	-	expression tag	UNP Q07010
B	0	HIS	-	expression tag	UNP Q07010

- Molecule 2 is [{2-[(guanine-9-yl)methyl]propane-1,3-diyl}bis(oxymethylene)]bis(phosphonic acid) (three-letter code: SV2) (formula: $C_{11}H_{19}N_5O_9P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	27	11	5	9	2	0	0
			Total	C	N	O	P		
2	B	1	27	11	5	9	2	0	0

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
3	A	1	1	1	0	0
			Total	Mg		
3	B	1	1	1	0	0

- Molecule 4 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	O			
4	A	1	7	4	3		0	0
			Total	C	O			
4	B	1	7	4	3		0	0

- Molecule 5 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula: $C_4H_{12}NO_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
5	A	1	8	4	1	3	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
5	B	1	8	4	1	3	0	0

- Molecule 6 is SULFATE ION (three-letter code: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
6	A	1	5	4	1	0	0

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
7	A	185	188	188	0	3
7	B	186	Total	O	0	6
			192	192		

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Hypoxanthine-guanine phosphoribosyltransferase

Chain A:



• Molecule 1: Hypoxanthine-guanine phosphoribosyltransferase

Chain B:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α, β, γ	116.01Å 93.89Å 44.72Å 90.00° 107.80° 90.00°	Depositor
Resolution (Å)	42.58 – 1.76 42.57 – 1.76	Depositor EDS
% Data completeness (in resolution range)	98.9 (42.58-1.76) 98.9 (42.57-1.76)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.12 (at 1.76Å)	Xtriage
Refinement program	PHENIX (1.11.1_2575: ???)	Depositor
R, R_{free}	0.158 , 0.192 0.158 , 0.192	Depositor DCC
R_{free} test set	2000 reflections (4.49%)	wwPDB-VP
Wilson B-factor (Å^2)	17.8	Xtriage
Anisotropy	0.481	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.37 , 56.5	EDS
L-test for twinning[2]	$< \|L\| > = 0.50$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.96	EDS
Total number of atoms	3435	wwPDB-VP
Average B, all atoms (Å^2)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.11% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: SV2, MG, SO4, PEG, TRS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.89	1/1540 (0.1%)	0.87	2/2087 (0.1%)
1	B	0.85	1/1490 (0.1%)	0.87	2/2019 (0.1%)
All	All	0.87	2/3030 (0.1%)	0.87	4/4106 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	180	GLU	CG-CD	5.80	1.60	1.51
1	B	178	TYR	CD2-CE2	5.26	1.47	1.39

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	114	ASP	CB-CG-OD1	7.79	125.31	118.30
1	B	114	ASP	CB-CG-OD1	7.34	124.91	118.30
1	B	179	ARG	NE-CZ-NH2	-6.35	117.12	120.30
1	A	149	ARG	NE-CZ-NH2	-5.06	117.77	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1508	0	1527	22	0
1	B	1456	0	1498	13	0
2	A	27	0	17	1	0
2	B	27	0	17	1	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	7	0	10	4	0
4	B	7	0	10	5	0
5	A	8	0	12	4	0
5	B	8	0	11	1	0
6	A	5	0	0	0	0
7	A	188	0	0	2	3
7	B	192	0	0	5	3
All	All	3435	0	3102	37	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:182:ARG:HH12	4:A:303:PEG:H32	1.34	0.90
1:A:105:ARG:NH2	1:A:133:GLU:OE2	2.07	0.87
1:A:182:ARG:HH22	4:A:303:PEG:H11	1.42	0.85
1:B:182:ARG:HH22	4:B:303:PEG:H31	1.41	0.84
1:A:46:PRO:O	7:A:401:HOH:O	2.11	0.69
1:A:182:ARG:NH1	4:A:303:PEG:H32	2.08	0.68
1:A:198:GLU:OE1	1:A:201:ARG:NH1	2.27	0.67
1:A:123:ARG:HH22	5:A:304:TRS:H32	1.60	0.66
1:A:182:ARG:HH22	4:A:303:PEG:C1	2.09	0.64
1:B:193:GLU:OE2	7:B:401:HOH:O	2.13	0.64
1:A:115:ILE:HD11	1:A:145:LYS:HD2	1.81	0.63
1:B:75[B]:ARG:NH1	7:B:403:HOH:O	2.28	0.62
1:A:176:GLN:HE22	1:A:179:ARG:HH11	1.46	0.61
1:A:66:ILE:HD11	4:B:303:PEG:H41	1.83	0.60
1:B:75[A]:ARG:NH1	7:B:406[A]:HOH:O	2.34	0.60
1:B:182:ARG:NH2	4:B:303:PEG:H31	2.14	0.60
5:A:304:TRS:H31	7:A:517:HOH:O	2.01	0.59
1:B:8:ASP:O	1:B:188:LYS:HE2	2.02	0.59
1:A:123:ARG:HH12	5:A:304:TRS:HN2	1.50	0.57
1:B:131:LYS:NZ	7:B:404:HOH:O	2.30	0.57
1:B:182:ARG:HH22	4:B:303:PEG:C3	2.15	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:123:ARG:NH2	5:A:304:TRS:H32	2.26	0.51
2:B:301:SV2:H6	7:B:410[B]:HOH:O	2.11	0.49
1:B:182:ARG:HH12	4:B:303:PEG:H31	1.78	0.49
1:A:193:GLU:O	1:A:197:LYS:HG3	2.13	0.48
1:A:117:ASP:OD2	1:A:148:GLY:HA3	2.13	0.48
1:B:75[B]:ARG:NH1	1:B:77:GLU:OE2	2.45	0.47
1:A:116:LEU:HD13	1:A:122:LEU:HD12	1.97	0.47
1:A:107:LYS:HB3	1:A:107:LYS:HE3	1.58	0.46
1:A:53:LEU:HD13	2:A:301:SV2:H10	1.98	0.46
1:B:50:VAL:HG13	1:B:79:LEU:HD13	1.98	0.45
1:A:176:GLN:NE2	1:A:179:ARG:HH11	2.13	0.44
1:B:180:GLU:HG2	5:B:304:TRS:N	2.33	0.43
1:B:162[B]:VAL:HG12	1:B:163:PRO:O	2.18	0.43
1:A:145:LYS:HG2	1:A:162:VAL:HG23	2.02	0.41
1:A:144:ASP:O	1:A:146:PRO:HD3	2.20	0.41
1:A:104:ILE:HD13	1:A:129:LEU:HD22	2.03	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:A:551:HOH:O	7:B:555:HOH:O[4_747]	2.09	0.11
7:A:427:HOH:O	7:B:543:HOH:O[4_748]	2.13	0.07
7:A:502:HOH:O	7:B:543:HOH:O[4_748]	2.13	0.07

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	186/216 (86%)	180 (97%)	6 (3%)	0	100	100
1	B	182/216 (84%)	177 (97%)	5 (3%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	368/432 (85%)	357 (97%)	11 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	169/186 (91%)	169 (100%)	0	100	100
1	B	163/186 (88%)	161 (99%)	2 (1%)	71	56
All	All	332/372 (89%)	330 (99%)	2 (1%)	92	79

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	12[A]	SER
1	B	12[B]	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	176	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 9 ligands modelled in this entry, 2 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	SV2	A	301	3	23,28,28	1.73	5 (21%)	28,41,41	1.83	7 (25%)
4	PEG	B	303	-	6,6,6	0.38	0	5,5,5	0.75	0
5	TRS	A	304	-	7,7,7	0.39	0	9,9,9	1.48	2 (22%)
6	SO4	A	305	-	4,4,4	0.15	0	6,6,6	0.15	0
4	PEG	A	303	-	6,6,6	0.39	0	5,5,5	0.72	0
2	SV2	B	301	3	23,28,28	2.03	7 (30%)	28,41,41	1.94	9 (32%)
5	TRS	B	304	-	7,7,7	0.93	0	9,9,9	1.34	1 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SV2	A	301	3	-	1/17/18/18	0/2/2/2
4	PEG	B	303	-	-	4/4/4/4	-
5	TRS	A	304	-	-	9/9/9/9	-
4	PEG	A	303	-	-	0/4/4/4	-
2	SV2	B	301	3	-	1/17/18/18	0/2/2/2
5	TRS	B	304	-	-	6/9/9/9	-

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	301	SV2	C2-N1	4.38	1.43	1.33

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	301	SV2	C2-N1	4.23	1.43	1.33
2	B	301	SV2	C2-N2	3.43	1.42	1.34
2	B	301	SV2	CAN-N9	-3.18	1.44	1.48
2	A	301	SV2	PBA-CAM	3.04	1.87	1.80
2	B	301	SV2	PAZ-CAL	2.98	1.87	1.80
2	B	301	SV2	PAZ-OAE	-2.83	1.48	1.54
2	B	301	SV2	PBA-CAM	2.80	1.87	1.80
2	A	301	SV2	C2-N2	2.61	1.40	1.34
2	A	301	SV2	PAZ-CAL	2.27	1.86	1.80
2	A	301	SV2	PAZ-OAE	-2.13	1.50	1.54
2	B	301	SV2	OAR-CAJ	-2.01	1.37	1.42

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	B	301	SV2	OAG-PBA-OAD	-4.28	101.07	112.39
2	A	301	SV2	OAH-PBA-OAD	-4.08	101.59	112.39
2	B	301	SV2	CAM-OAS-CAK	-3.58	104.31	112.84
2	A	301	SV2	CAM-OAS-CAK	-3.55	104.37	112.84
2	A	301	SV2	CAN-CAX-CAK	-3.13	105.13	111.01
2	B	301	SV2	OAG-PBA-CAM	3.09	116.16	106.68
2	B	301	SV2	OAH-PBA-CAM	-3.00	97.48	106.68
2	A	301	SV2	OAH-PBA-CAM	3.00	115.87	106.68
2	B	301	SV2	OAF-PAZ-OAE	2.90	116.54	108.08
2	A	301	SV2	OAG-PBA-CAM	-2.81	98.06	106.68
2	A	301	SV2	C8-N7-C5	2.37	107.50	102.99
2	A	301	SV2	N2-C2-N3	2.36	121.73	116.72
2	B	301	SV2	C8-N7-C5	2.35	107.46	102.99
5	B	304	TRS	O1-C1-C	-2.20	104.02	111.00
2	B	301	SV2	N2-C2-N3	2.20	121.38	116.72
2	B	301	SV2	OAH-PBA-OAD	2.17	118.12	112.39
5	A	304	TRS	O1-C1-C	-2.10	104.34	111.00
5	A	304	TRS	C2-C-N	-2.09	101.73	107.98
2	B	301	SV2	OAH-PBA-OAG	2.07	114.12	108.08

There are no chirality outliers.

All (21) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	304	TRS	C2-C-C3-O3
5	B	304	TRS	C2-C-C1-O1
5	B	304	TRS	C3-C-C1-O1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
5	B	304	TRS	N-C-C1-O1
5	B	304	TRS	C1-C-C2-O2
5	B	304	TRS	C3-C-C2-O2
5	B	304	TRS	N-C-C2-O2
4	B	303	PEG	O1-C1-C2-O2
4	B	303	PEG	O2-C3-C4-O4
5	A	304	TRS	C1-C-C3-O3
4	B	303	PEG	C1-C2-O2-C3
5	A	304	TRS	C2-C-C1-O1
5	A	304	TRS	N-C-C1-O1
5	A	304	TRS	C3-C-C2-O2
5	A	304	TRS	N-C-C3-O3
5	A	304	TRS	C1-C-C2-O2
4	B	303	PEG	C4-C3-O2-C2
5	A	304	TRS	C3-C-C1-O1
5	A	304	TRS	N-C-C2-O2
2	A	301	SV2	PAZ-CAL-OAR-CAJ
2	B	301	SV2	PAZ-CAL-OAR-CAJ

There are no ring outliers.

6 monomers are involved in 16 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	301	SV2	1	0
4	B	303	PEG	5	0
5	A	304	TRS	4	0
4	A	303	PEG	4	0
2	B	301	SV2	1	0
5	B	304	TRS	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand SV2 A 301



Ligand SV2 B 301



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	185/216 (85%)	-0.11	7 (3%)	40	47	8, 19, 54, 71	0
1	B	179/216 (82%)	-0.12	6 (3%)	45	51	9, 18, 50, 73	0
All	All	364/432 (84%)	-0.11	13 (3%)	42	49	8, 19, 54, 73	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	200	GLU	5.0
1	B	204	ALA	4.3
1	A	-2	HIS	4.0
1	B	203	LYS	3.7
1	A	101	LEU	3.5
1	A	4	ALA	3.5
1	B	193	GLU	2.8
1	B	200	GLU	2.7
1	A	3	PRO	2.5
1	A	199	LEU	2.4
1	B	8	ASP	2.2
1	A	201	ARG	2.1
1	B	197	LYS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
3	MG	A	302	1/1	0.71	0.11	65,65,65,65	0
6	SO4	A	305	5/5	0.76	0.21	108,108,109,111	0
4	PEG	B	303	7/7	0.86	0.14	42,45,48,49	0
3	MG	B	302	1/1	0.86	0.17	40,40,40,40	0
4	PEG	A	303	7/7	0.88	0.12	33,37,41,43	0
5	TRS	B	304	8/8	0.90	0.16	35,40,48,48	0
5	TRS	A	304	8/8	0.90	0.29	24,50,61,65	0
2	SV2	B	301	27/27	0.97	0.07	12,17,34,38	0
2	SV2	A	301	27/27	0.98	0.07	12,18,42,47	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around SV2 B 301:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around SV2 A 301:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.