



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 22, 2023 – 07:25 pm BST

PDB ID : 8ART
Title : ABC transporter binding protein MalE from *Streptomyces scabiei* in complex with maltose
Authors : Jadot, C.; Kerff, F.; Rigali, S.
Deposited on : 2022-08-17
Resolution : 3.17 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.17 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1467 (3.20-3.16)
Clashscore	141614	1599 (3.20-3.16)
Ramachandran outliers	138981	1574 (3.20-3.16)
Sidechain outliers	138945	1573 (3.20-3.16)
RSRZ outliers	127900	1423 (3.20-3.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	429	 8% 88% 9%
1	B	429	 10% 87% 8%
2	C	2	 50% 50%
2	D	2	 50% 50%

2 Entry composition i

There are 2 unique types of molecules in this entry. The entry contains 11866 atoms, of which 5873 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative secreted maltose-binding protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	392	Total	C	H	N	O	S	0	0	0
			5878	1899	2909	488	581	1			
1	B	393	Total	C	H	N	O	S	0	0	0
			5902	1905	2924	490	582	1			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	424	HIS	-	expression tag	UNP C9ZHD5
A	425	HIS	-	expression tag	UNP C9ZHD5
A	426	HIS	-	expression tag	UNP C9ZHD5
A	427	HIS	-	expression tag	UNP C9ZHD5
A	428	HIS	-	expression tag	UNP C9ZHD5
A	429	HIS	-	expression tag	UNP C9ZHD5
B	424	HIS	-	expression tag	UNP C9ZHD5
B	425	HIS	-	expression tag	UNP C9ZHD5
B	426	HIS	-	expression tag	UNP C9ZHD5
B	427	HIS	-	expression tag	UNP C9ZHD5
B	428	HIS	-	expression tag	UNP C9ZHD5
B	429	HIS	-	expression tag	UNP C9ZHD5

- Molecule 2 is an oligosaccharide called alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	H	O			
2	C	2	Total	C	H	O	0	0	0
			43	12	20	11			
2	D	2	Total	C	H	O	0	0	0
			43	12	20	11			


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Putative secreted maltose-binding protein



- Molecule 1: Putative secreted maltose-binding protein



- Molecule 2: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose



- Molecule 2: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	68.31Å 68.31Å 260.55Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	48.89 – 3.17 43.79 – 3.17	Depositor EDS
% Data completeness (in resolution range)	99.4 (48.89-3.17) 99.5 (43.79-3.17)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.40 (at 3.19Å)	Xtrriage
Refinement program	BUSTER 2.10.3	Depositor
R, R_{free}	0.260 , 0.284 0.288 , 0.309	Depositor DCC
R_{free} test set	631 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	126.2	Xtrriage
Anisotropy	0.038	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 58.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.41$, $\langle L^2 \rangle = 0.23$	Xtrriage
Estimated twinning fraction	0.078 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	11866	wwPDB-VP
Average B, all atoms (Å ²)	125.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.27% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.34	0/3042	0.48	0/4137
1	B	0.34	0/3052	0.47	0/4151
All	All	0.34	0/6094	0.47	0/8288

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2969	2909	2910	5	0
1	B	2978	2924	2924	7	0
2	C	23	20	21	0	0
2	D	23	20	21	0	0
All	All	5993	5873	5876	12	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (12) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:36:LEU:HD13	1:B:93:ILE:HD11	1.83	0.60
1:B:93:ILE:HD13	1:B:326:PHE:CD1	2.39	0.57
1:A:346:LEU:HD13	1:A:370:LEU:HD11	1.89	0.53
1:A:421:TYR:HB2	1:A:422:SER:HA	1.92	0.51
1:B:150:ALA:HB2	1:B:369:VAL:HG21	1.94	0.49
1:A:150:ALA:HB2	1:A:369:VAL:HG21	1.94	0.49
1:B:346:LEU:HD13	1:B:370:LEU:HD11	1.95	0.48
1:B:346:LEU:HD11	1:B:366:TYR:HB3	1.99	0.44
1:A:217:ILE:CG1	1:A:421:TYR:CD2	3.01	0.43
1:B:226:LYS:NZ	1:B:296:THR:HG21	2.35	0.41
1:A:421:TYR:CB	1:A:422:SER:HA	2.50	0.41
1:B:252:HIS:N	1:B:252:HIS:ND1	2.68	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	388/429 (90%)	375 (97%)	13 (3%)	0	100	100
1	B	391/429 (91%)	377 (96%)	13 (3%)	1 (0%)	41	73
All	All	779/858 (91%)	752 (96%)	26 (3%)	1 (0%)	51	83

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	422	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	301/325 (93%)	292 (97%)	9 (3%)	41	72
1	B	302/325 (93%)	292 (97%)	10 (3%)	38	70
All	All	603/650 (93%)	584 (97%)	19 (3%)	39	70

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	46	ASN
1	A	66	LYS
1	A	147	ASP
1	A	165	GLU
1	A	171	ASP
1	A	252	HIS
1	A	412	THR
1	A	418	VAL
1	A	420	ASP
1	B	147	ASP
1	B	165	GLU
1	B	171	ASP
1	B	208	ASP
1	B	226	LYS
1	B	252	HIS
1	B	308	ASN
1	B	412	THR
1	B	418	VAL
1	B	420	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates i

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GLC	C	1	2	12,12,12	0.24	0	17,17,17	0.63	0
2	GLC	C	2	2	11,11,12	0.26	0	15,15,17	0.74	1 (6%)
2	GLC	D	1	2	12,12,12	0.21	0	17,17,17	0.96	1 (5%)
2	GLC	D	2	2	11,11,12	0.23	0	15,15,17	0.72	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	C	1	2	-	1/2/22/22	0/1/1/1
2	GLC	C	2	2	-	0/2/19/22	0/1/1/1
2	GLC	D	1	2	-	1/2/22/22	0/1/1/1
2	GLC	D	2	2	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	2	GLC	C1-O5-C5	2.33	115.34	112.19
2	D	1	GLC	C1-O5-C5	2.29	117.98	113.66

There are no chirality outliers.


All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	D	1	GLC	O5-C5-C6-O6
2	C	1	GLC	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	392/429 (91%)	0.63	36 (9%) 9 5	55, 122, 138, 147	0
1	B	393/429 (91%)	0.73	44 (11%) 5 3	89, 129, 143, 150	0
All	All	785/858 (91%)	0.68	80 (10%) 6 3	55, 126, 141, 150	0

All (80) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	31	GLY	7.5
1	A	67	VAL	5.8
1	A	34	VAL	5.1
1	B	60	ALA	4.7
1	A	32	GLY	4.7
1	B	63	LYS	4.5
1	A	57	GLU	4.4
1	B	108	GLY	4.4
1	B	96	SER	4.1
1	B	121	ASP	4.1
1	A	58	PHE	4.1
1	A	36	LEU	4.1
1	A	65	ILE	3.9
1	B	326	PHE	3.9
1	A	311	VAL	3.9
1	A	33	PRO	3.9
1	B	144	PHE	3.9
1	A	69	TYR	3.7
1	A	320	GLN	3.7
1	B	311	VAL	3.6
1	B	421	TYR	3.6
1	B	82	ASP	3.5
1	B	34	VAL	3.5
1	B	274	ASN	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	110	PHE	3.4
1	B	219	VAL	3.4
1	A	54	LEU	3.3
1	A	422	SER	3.3
1	B	32	GLY	3.3
1	B	133	ALA	3.2
1	A	68	LYS	3.1
1	B	416	LYS	3.1
1	B	92	ASP	3.0
1	B	118	ALA	3.0
1	A	327	ILE	2.9
1	B	84	ALA	2.9
1	A	37	THR	2.9
1	A	64	ASP	2.8
1	A	88	THR	2.8
1	B	278	GLY	2.7
1	B	309	LEU	2.7
1	B	275	PHE	2.7
1	B	119	LEU	2.7
1	B	94	LEU	2.7
1	A	93	ILE	2.7
1	A	228	TYR	2.6
1	A	219	VAL	2.6
1	B	33	PRO	2.6
1	B	366	TYR	2.6
1	B	300	ALA	2.6
1	B	325	LYS	2.6
1	A	205	ASP	2.6
1	A	346	LEU	2.5
1	A	269	PRO	2.5
1	A	262	VAL	2.5
1	B	134	GLN	2.5
1	A	421	TYR	2.5
1	A	309	LEU	2.5
1	A	186	VAL	2.5
1	B	86	GLY	2.5
1	B	85	ALA	2.4
1	B	149	LEU	2.4
1	B	310	SER	2.3
1	B	287	LEU	2.2
1	B	353	TYR	2.2
1	A	423	LYS	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	377	PRO	2.2
1	B	38	TRP	2.2
1	B	312	TYR	2.1
1	B	313	ALA	2.1
1	B	156	GLN	2.1
1	A	195	ALA	2.1
1	A	38	TRP	2.1
1	A	59	GLU	2.1
1	A	66	LYS	2.1
1	A	55	VAL	2.1
1	B	281	PHE	2.0
1	B	184	THR	2.0
1	A	351	ASP	2.0
1	B	401	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GLC	C	1	12/12	0.86	0.21	106,111,114,115	0
2	GLC	C	2	11/12	0.89	0.24	96,99,103,104	0
2	GLC	D	1	12/12	0.92	0.21	86,88,95,100	0
2	GLC	D	2	11/12	0.93	0.23	80,84,86,87	0




The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain C:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain D:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.