



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 7, 2023 – 01:20 pm GMT

PDB ID : 8AS9
Title : Crystal structure of the talin-KANK1 complex
Authors : Zacharchenko, T.
Deposited on : 2022-08-18
Resolution : 3.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1026 (3.48-3.32)
Clashscore	141614	1055 (3.48-3.32)
Ramachandran outliers	138981	1038 (3.48-3.32)
Sidechain outliers	138945	1038 (3.48-3.32)
RSRZ outliers	127900	2173 (3.50-3.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	137	 2% (poor fit), 79% (0-1 outliers), 9% (2 outliers), 12% (3+ outliers)
1	B	137	 4% (poor fit), 90% (0-1 outliers), 7% (2 outliers), .. (3+ outliers)
2	D	51	 2% (poor fit), 65% (0-1 outliers), 18% (2 outliers), 18% (3+ outliers)
3	C	309	 18% (poor fit), 84% (0-1 outliers), 14% (2 outliers), .. (3+ outliers)

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4704 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called B-cell lymphoma 6 protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	121	996	630	176	182	8	0	3	0
1	B	133	1077	679	191	199	8	0	2	0

There are 32 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	GLY	-	expression tag	UNP P41182
A	-6	PRO	-	expression tag	UNP P41182
A	-5	SER	-	expression tag	UNP P41182
A	-4	SER	-	expression tag	UNP P41182
A	-3	ASP	-	expression tag	UNP P41182
A	-2	LEU	-	expression tag	UNP P41182
A	-1	TYR	-	expression tag	UNP P41182
A	0	LEU	-	expression tag	UNP P41182
A	1	ARG	-	expression tag	UNP P41182
A	2	PRO	-	expression tag	UNP P41182
A	3	GLY	-	expression tag	UNP P41182
A	4	GLY	-	expression tag	UNP P41182
A	5	GLY	-	expression tag	UNP P41182
A	8	GLN	CYS	engineered mutation	UNP P41182
A	67	ARG	CYS	engineered mutation	UNP P41182
A	84	ASN	CYS	engineered mutation	UNP P41182
B	-7	GLY	-	expression tag	UNP P41182
B	-6	PRO	-	expression tag	UNP P41182
B	-5	SER	-	expression tag	UNP P41182
B	-4	SER	-	expression tag	UNP P41182
B	-3	ASP	-	expression tag	UNP P41182
B	-2	LEU	-	expression tag	UNP P41182
B	-1	TYR	-	expression tag	UNP P41182
B	0	LEU	-	expression tag	UNP P41182
B	1	ARG	-	expression tag	UNP P41182

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	2	PRO	-	expression tag	UNP P41182
B	3	GLY	-	expression tag	UNP P41182
B	4	GLY	-	expression tag	UNP P41182
B	5	GLY	-	expression tag	UNP P41182
B	8	GLN	CYS	engineered mutation	UNP P41182
B	67	ARG	CYS	engineered mutation	UNP P41182
B	84	ASN	CYS	engineered mutation	UNP P41182

- Molecule 2 is a protein called KN-motif NCoR1 BBD fusion,Nuclear receptor corepressor 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	D	42	351	229	55	66	1	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	32	GLY	-	linker	UNP Q8BRZ8
D	33	GLY	-	linker	UNP Q8BRZ8
D	34	GLY	-	linker	UNP Q8BRZ8
D	35	GLY	-	linker	UNP Q8BRZ8

- Molecule 3 is a protein called Talin-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	306	2254	1385	406	448	15	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	1351	GLY	-	expression tag	UNP P26039
C	1352	ILE	-	expression tag	UNP P26039
C	1353	ASP	-	expression tag	UNP P26039
C	1354	PRO	-	expression tag	UNP P26039
C	1355	PHE	-	expression tag	UNP P26039
C	1356	THR	-	expression tag	UNP P26039
C	1357	LYS	-	expression tag	UNP P26039
C	1358	HIS	-	expression tag	UNP P26039

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	C	1	Total	O	S	0	0
			5	4	1		
4	C	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	C	1	Total	C	O	0	0
			6	3	3		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: B-cell lymphoma 6 protein





- Molecule 1: B-cell lymphoma 6 protein



- Molecule 2: KN-motif NCoR1 BBD fusion,Nuclear receptor corepressor 1



- Molecule 3: Talin-1



4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	207.02Å 207.02Å 151.86Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	59.76 – 3.40 61.88 – 3.40	Depositor EDS
% Data completeness (in resolution range)	99.3 (59.76-3.40) 99.8 (61.88-3.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.23 (at 3.40Å)	Xtrriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.245 , 0.285 0.246 , 0.286	Depositor DCC
R_{free} test set	872 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	99.7	Xtrriage
Anisotropy	0.242	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 77.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.27$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4704	wwPDB-VP
Average B, all atoms (Å ²)	127.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.05% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.24	0/1013	0.46	0/1366
1	B	0.24	0/1093	0.48	0/1474
2	D	0.23	0/359	0.44	0/481
3	C	0.24	0/2286	0.44	0/3100
All	All	0.24	0/4751	0.45	0/6421

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	996	0	1015	7	0
1	B	1077	0	1092	6	0
2	D	351	0	346	4	0
3	C	2254	0	2251	26	0
4	A	10	0	0	1	0
4	C	10	0	0	0	0
5	C	6	0	8	0	0
All	All	4704	0	4712	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 4.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:1361:LYS:O	3:C:1365:ASN:ND2	2.31	0.64
2:D:4:VAL:HG22	2:D:12:LEU:HB2	1.81	0.62
3:C:1487:THR:OG1	3:C:1488:GLN:N	2.34	0.59
1:A:23:ASN:OD1	1:A:26:ARG:NH2	2.39	0.56
1:A:24:ARG:NH2	3:C:1394:ASP:OD1	2.40	0.55
1:B:103:MET:HG2	3:C:1510:ARG:HD2	1.90	0.54
1:B:77:GLU:HB3	3:C:1517:ALA:HA	1.91	0.53
3:C:1376:LEU:HD22	3:C:1384:ILE:HD12	1.91	0.53
3:C:1513:SER:HB2	3:C:1522:LYS:HG3	1.91	0.53
2:D:13:ASP:HB2	3:C:1638:ARG:HG2	1.92	0.51
1:B:24:ARG:NH1	2:D:49:ILE:O	2.43	0.51
3:C:1461:LEU:HG	3:C:1516:THR:HG22	1.92	0.50
3:C:1362:GLU:OE1	3:C:1362:GLU:N	2.44	0.50
3:C:1555:ARG:NH1	3:C:1558:CYS:SG	2.84	0.50
3:C:1488:GLN:O	3:C:1490:GLN:N	2.43	0.49
3:C:1365:ASN:OD1	3:C:1368:ARG:NH2	2.45	0.49
3:C:1620:LEU:HD12	3:C:1633:LEU:HD12	1.95	0.49
3:C:1550:PHE:HD1	3:C:1550:PHE:H	1.60	0.48
3:C:1366:ALA:HB1	3:C:1430:SER:HB2	1.96	0.48
3:C:1564:PRO:HA	3:C:1567:GLU:HG2	1.95	0.47
4:A:202:SO4:O3	3:C:1652:ARG:NH1	2.48	0.47
2:D:1:PRO:O	2:D:18:LYS:HB2	2.15	0.47
3:C:1546:LEU:HD21	3:C:1555:ARG:HG2	1.97	0.47
1:A:31:LEU:HA	1:B:67:ARG:HH21	1.80	0.46
3:C:1426:ILE:HD12	3:C:1614:ILE:HG23	1.98	0.45
1:B:38:VAL:HG22	1:B:74:LEU:HD12	1.99	0.45
1:A:106:MET:HE2	1:A:121:CYS:HB3	2.00	0.44
3:C:1458:GLN:HE21	3:C:1584:VAL:HG23	1.82	0.43
3:C:1476:MET:HA	3:C:1479:GLN:HE22	1.83	0.43
3:C:1383:PRO:HD3	3:C:1445:TYR:CZ	2.54	0.42
1:A:79:ASN:OD1	1:A:82:GLY:N	2.51	0.41
1:B:19:LEU:HD11	1:B:91:TYR:HA	2.02	0.41
1:A:20:LEU:O	1:A:24:ARG:HG3	2.22	0.40
1:A:86:LEU:O	1:A:90:MET:HG3	2.22	0.40
3:C:1489:ALA:O	3:C:1491:VAL:N	2.53	0.40
3:C:1483:GLU:O	3:C:1485:GLY:N	2.54	0.40
3:C:1526:VAL:HG12	3:C:1530:LYS:HE2	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	122/137 (89%)	118 (97%)	4 (3%)	0	100	100
1	B	133/137 (97%)	127 (96%)	6 (4%)	0	100	100
2	D	38/51 (74%)	37 (97%)	0	1 (3%)	5	26
3	C	304/309 (98%)	282 (93%)	18 (6%)	4 (1%)	12	39
All	All	597/634 (94%)	564 (94%)	28 (5%)	5 (1%)	19	51

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	C	1485	GLY
3	C	1490	GLN
3	C	1488	GLN
2	D	2	TYR
3	C	1482	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	114/123 (93%)	112 (98%)	2 (2%)	59	79
1	B	122/123 (99%)	119 (98%)	3 (2%)	47	72
2	D	39/44 (89%)	37 (95%)	2 (5%)	24	54
3	C	241/243 (99%)	235 (98%)	6 (2%)	47	72
All	All	516/533 (97%)	503 (98%)	13 (2%)	47	72

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	29	ASP
1	A	40	ARG
1	B	6	ASP
1	B	65	LEU
1	B	77	GLU
2	D	25	LYS
2	D	47	HIS
3	C	1360	GLN
3	C	1463	GLU
3	C	1475	GLN
3	C	1505	LEU
3	C	1550	PHE
3	C	1555	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
3	C	1658	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	SO4	C	1703	-	4,4,4	0.13	0	6,6,6	0.05	0
4	SO4	A	201	-	4,4,4	0.14	0	6,6,6	0.05	0
4	SO4	C	1702	-	4,4,4	0.14	0	6,6,6	0.05	0
4	SO4	A	202	-	4,4,4	0.14	0	6,6,6	0.05	0
5	GOL	C	1701	-	5,5,5	0.91	0	5,5,5	1.00	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	C	1701	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	202	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	121/137 (88%)	0.36	3 (2%) 57 55	73, 99, 139, 169	0
1	B	133/137 (97%)	0.30	6 (4%) 33 33	68, 95, 153, 180	0
2	D	42/51 (82%)	0.58	1 (2%) 59 57	124, 134, 161, 177	0
3	C	306/309 (99%)	0.86	55 (17%) 1 1	75, 123, 284, 352	0
All	All	602/634 (94%)	0.62	65 (10%) 5 7	68, 111, 248, 352	0

All (65) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	C	1484	PRO	13.6
3	C	1483	GLU	6.5
3	C	1493	SER	6.1
3	C	1554	ASN	5.7
3	C	1544	LYS	5.5
3	C	1545	ALA	5.3
3	C	1478	CYS	5.2
2	D	1	PRO	5.1
3	C	1546	LEU	5.0
3	C	1481	LEU	4.8
3	C	1476	MET	4.8
3	C	1480	SER	4.7
3	C	1487	THR	4.7
3	C	1548	GLY	4.6
3	C	1543	ILE	4.4
3	C	1485	GLY	4.2
3	C	1479	GLN	4.2
3	C	1551	THR	4.2
3	C	1558	CYS	4.2
3	C	1542	THR	4.1
3	C	1491	VAL	4.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	C	1496	THR	3.9
3	C	1488	GLN	3.9
3	C	1477	ALA	3.8
3	C	1475	GLN	3.8
3	C	1486	CYS	3.8
3	C	1489	ALA	3.7
3	C	1498	VAL	3.7
3	C	1497	ILE	3.7
3	C	1495	ALA	3.6
3	C	1474	ILE	3.6
3	C	1547	ASP	3.5
3	C	1555	ARG	3.5
1	B	-4	SER	3.5
3	C	1556	ALA	3.5
3	C	1492	LEU	3.3
3	C	1490	GLN	3.2
1	B	97	LEU	3.1
3	C	1536	THR	3.0
3	C	1540	VAL	2.9
3	C	1581	PHE	2.8
3	C	1541	LYS	2.8
3	C	1559	ARG	2.7
3	C	1482	GLY	2.6
3	C	1552	GLU	2.6
3	C	1550	PHE	2.6
3	C	1500	LYS	2.5
3	C	1561	ALA	2.5
1	B	1	ARG	2.4
3	C	1560	ALA	2.4
3	C	1630	TRP	2.4
3	C	1467	PHE	2.3
1	B	-1	TYR	2.3
1	A	67	ARG	2.2
3	C	1462	VAL	2.2
3	C	1553	GLU	2.2
3	C	1557	GLN	2.2
1	A	72	ILE	2.1
3	C	1494	ALA	2.1
3	C	1572	LEU	2.1
3	C	1529	ALA	2.1
3	C	1539	LEU	2.1
1	B	-3	ASP	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	57	PHE	2.0
1	B	105	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	SO4	A	201	5/5	0.77	0.17	134,136,152,154	0
5	GOL	C	1701	6/6	0.92	0.76	120,120,125,125	6
4	SO4	C	1703	5/5	0.94	0.36	111,112,113,115	5
4	SO4	A	202	5/5	0.96	0.16	81,104,112,121	0
4	SO4	C	1702	5/5	0.98	0.18	102,103,108,109	5

6.5 Other polymers [i](#)

There are no such residues in this entry.