

Full wwPDB X-ray Structure Validation Report (i)

Jul 26, 2023 – 01:16 AM EDT

PDB ID : 1AXW

Title : E. COLI THYMIDYLATE SYNTHASE IN COMPLEX WITH

METHOTREXATE (MTX) AND 2'-DEOXYURIDINE 5'-

MONOPHOSPHATE (DUMP)

Authors: Stout, T.J.; Sage, C.R.; Stroud, R.M.

Deposited on : 1997-10-23

Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : NOT EXECUTED

EDS : NOT EXECUTED

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.34

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY\ DIFFRACTION$

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution	
Metric	$(\# \mathrm{Entries})$	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$	
Clashscore	141614	4695 (1.70-1.70)	
Ramachandran outliers	138981	4610 (1.70-1.70)	
Sidechain outliers	138945	4610 (1.70-1.70)	

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain	
1	A	265	94%	6%
1	В	265	89%	11%



2 Entry composition (i)


There are 4 unique types of molecules in this entry. The entry contains 4600 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called THYMIDYLATE SYNTHASE.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	A	265	Total 2153	C 1375	N 371	O 395	S 12	0	0	0
1	В	265	Total 2153	C 1375	N 371	O 395	S 12	0	0	0


• Molecule 2 is 2'-DEOXYURIDINE 5'-MONOPHOSPHATE (three-letter code: UMP) (formula: $C_9H_{13}N_2O_8P$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
9	Λ	1	Total	otal C N O P		0	0		
	2 A	1	20	9	2	8	1	0	0
2	D	1	Total	С	N	О	Р	0	0
	Б	1	20	9	2	8	1		_

• Molecule 3 is METHOTREXATE (three-letter code: MTX) (formula: $C_{20}H_{22}N_8O_5$).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
3	Δ	1	Total	С	N	О	0	0	
9	Λ	1	33	20	8	5	0		
2	D	1	Total	С	N	О	0	0	
)	Ъ	1	33	20	8	5	0		

• Molecule 4 is water.

Mo	ol	Chain	Residues	Atoms	ZeroOcc	AltConf
4		A	124	Total O 124 124	0	0
4		В	64	Total O 64 64	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: THYMIDYLATE SYNTHASE





4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source	
Space group	P 63	Depositor	
Cell constants	127.34Å 127.34Å 68.06Å	Depositor	
a, b, c, α , β , γ	90.00° 90.00° 120.00°	Depositor	
Resolution (Å)	50.00 - 1.70	Depositor	
% Data completeness	78.3 (50.00-1.70)	Depositor	
(in resolution range)	10.0 (00.00 1.10)	Depositor	
R_{merge}	0.09	Depositor	
R_{sym}	(Not available)	Depositor	
Refinement program	X-PLOR 3.854	Depositor	
R, R_{free}	0.217 , 0.265	Depositor	
Estimated twinning fraction	No twinning to report.	Xtriage	
Total number of atoms	4600	wwPDB-VP	
Average B, all atoms (Å ²)	28.0	wwPDB-VP	



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: FMT, MTX, UMP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.40	0/2210	0.64	1/3000 (0.0%)	
1	В	0.38	0/2210	0.64	0/3000	
All	All	0.39	0/4420	0.64	1/6000 (0.0%)	

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^{o})$	$\operatorname{Ideal}({}^{o})$
1	A	27	LEU	N-CA-C	-5.16	97.07	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2153	0	2080	6	0
1	В	2153	0	2080	11	0
2	A	20	0	11	1	0
2	В	20	0	11	1	0
3	A	33	0	20	1	0
3	В	33	0	20	0	0
4	A	124	0	0	0	0
4	В	64	0	0	0	0

Continued on next page...



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	4600	0	4222	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	${\rm distance}(\mathring{\rm A})$	overlap (Å)
1:B:147:HIS:HB2	1:B:163:LEU:HD11	1.87	0.57
1:A:146:CYS:SG	2:A:603:UMP:C6	3.01	0.54
1:A:147:HIS:HB2	1:A:163:LEU:HD11	1.90	0.54
1:B:146:CYS:SG	2:B:604:UMP:C6	3.01	0.54
1:A:5:LEU:HD11	1:A:47:THR:HG21	1.90	0.52
1:B:0:FMT:O1	1:B:45:VAL:HG13	2.11	0.51
1:A:7:LEU:O	1:A:11:VAL:HG23	2.14	0.47
1:B:44:LEU:HD21	1:B:52:LEU:HD21	1.97	0.46
1:B:234:ARG:O	1:B:236:PRO:HD3	2.16	0.45
1:B:5:LEU:HD11	1:B:47:THR:HG21	2.00	0.44
1:B:1:MET:HG2	1:B:227:LEU:HD21	2.01	0.42
3:A:732:MTX:H13	3:A:732:MTX:HM1	1.89	0.42
1:B:135:VAL:HA	1:B:138:LEU:HD22	2.02	0.41
1:B:33:GLN:HA	1:B:201:TRP:O	2.21	0.41
1:B:181:TYR:O	1:B:185:VAL:HG23	2.20	0.41
1:A:7:LEU:HD13	1:A:32:HIS:CD2	2.56	0.40
1:B:133:TRP:CZ2	1:B:138:LEU:HD21	2.56	0.40
1:A:33:GLN:HA	1:A:201:TRP:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	263/265 (99%)	256 (97%)	5 (2%)	2 (1%)	19 6
1	В	263/265 (99%)	254 (97%)	8 (3%)	1 (0%)	34 18
All	All	526/530 (99%)	510 (97%)	13 (2%)	3 (1%)	25 11

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	В	94	TYR
1	A	94	TYR
1	A	93	VAL

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric Outliers		Percentiles		
1	A	233/233 (100%)	229 (98%)	4 (2%)	60	46	
1	В	$233/233 \ (100\%)$	224 (96%)	9 (4%)	32	13	
All	All	466/466 (100%)	453 (97%)	13 (3%)	43	25	

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	38	LEU
1	A	116	LEU
1	A	176	PHE
1	A	239	ILE
1	В	17	GLN
1	В	39	GLN
1	В	51	HIS
1	В	96	LYS
1	В	105	ASP
1	В	138	LEU
1	В	176	PHE
1	В	218	LEU
1	В	231	ILE



Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	33	GLN
1	A	118	GLN
1	A	151	GLN
1	A	217	HIS
1	В	17	GLN
1	В	33	GLN
1	В	51	HIS
1	В	108	HIS
1	В	118	GLN
1	В	151	GLN
1	В	162	GLN
1	В	186	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mol	Trme	Chain	hain Res Lin		Bond lengths				Bond angles			
MIOI	Type	Chain	nes	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2		
3	MTX	A	732	-	35,35,35	2.57	5 (14%)	46,49,49	2.34	10 (21%)		
3	MTX	В	733	-	35,35,35	2.45	5 (14%)	46,49,49	2.70	12 (26%)		
2	UMP	В	604	-	21,21,21	1.57	4 (19%)	31,31,31	1.91	9 (29%)		
2	UMP	A	603	-	21,21,21	1.52	4 (19%)	31,31,31	1.93	9 (29%)		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MTX	A	732	-	-	5/25/25/25	0/3/3/3
3	MTX	В	733	-	-	8/25/25/25	0/3/3/3
2	UMP	В	604	-	-	1/10/22/22	0/2/2/2
2	UMP	A	603	-	-	1/10/22/22	0/2/2/2

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(A)	Ideal(Å)
3	A	732	MTX	C7-N8	11.68	1.50	1.31
3	В	733	MTX	C7-N8	10.92	1.49	1.31
3	A	732	MTX	C7-C6	6.02	1.50	1.39
3	В	733	MTX	C7-C6	5.81	1.49	1.39
2	В	604	UMP	O4'-C1'	4.36	1.52	1.42
2	A	603	UMP	O4'-C1'	4.00	1.51	1.42
3	A	732	MTX	CM-N10	2.93	1.51	1.46
2	В	604	UMP	O4'-C4'	2.82	1.51	1.45
2	A	603	UMP	O4'-C4'	2.67	1.51	1.45
3	В	733	MTX	CM-N10	2.63	1.50	1.46
3	В	733	MTX	C6-N5	2.43	1.36	1.32
2	В	604	UMP	P-OP2	-2.26	1.46	1.54
3	A	732	MTX	C15-C14	2.24	1.43	1.39
2	A	603	UMP	P-OP3	-2.19	1.46	1.54
2	В	604	UMP	C6-C5	2.18	1.40	1.35
2	A	603	UMP	C6-C5	2.10	1.39	1.35
3	A	732	MTX	C16-C11	2.03	1.42	1.39
3	В	733	MTX	C9-C6	-2.00	1.48	1.51

All (40) bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$Ideal(^{o})$
3	В	733	MTX	C6-C7-N8	-10.86	112.48	123.13
3	A	732	MTX	C6-C7-N8	-9.71	113.61	123.13
3	В	733	MTX	N1-C2-N3	-6.28	118.84	127.22
2	A	603	UMP	O4'-C1'-N1	5.68	118.01	107.86
2	В	604	UMP	O4'-C1'-N1	5.64	117.93	107.86
3	В	733	MTX	C7-N8-C8A	5.54	122.26	116.69
3	В	733	MTX	CA-N-C	5.53	135.14	121.60
3	A	732	MTX	N1-C2-N3	-5.05	120.48	127.22
3	В	733	MTX	C2-N1-C8A	4.79	120.82	115.36
3	A	732	MTX	C2-N1-C8A	4.33	120.31	115.36
3	В	733	MTX	C4A-C4-N3	-4.31	118.18	121.01
3	A	732	MTX	C7-C6-N5	-4.20	118.10	120.85
3	A	732	MTX	CA-N-C	3.87	131.06	121.60
2	A	603	UMP	C1'-N1-C6	-3.84	113.97	121.55
2	В	604	UMP	C1'-N1-C6	-3.65	114.35	121.55
3	A	732	MTX	N8-C8A-N1	3.56	119.88	115.82
3	В	733	MTX	N8-C8A-N1	3.48	119.79	115.82
3	A	732	MTX	C4A-C4-N3	-3.25	118.88	121.01
2	A	603	UMP	O4'-C1'-C2'	-3.13	100.33	106.25
2	A	603	UMP	C2'-C1'-N1	3.12	120.96	113.77
2	В	604	UMP	P-O5'-C5'	3.03	126.63	118.30
3	A	732	MTX	C9-C6-N5	2.95	121.69	116.96
2	A	603	UMP	O4'-C4'-C3'	-2.87	98.98	105.67
2	В	604	UMP	O4'-C4'-C3'	-2.84	99.04	105.67
2	В	604	UMP	C6-N1-C2	2.74	124.50	120.99
3	A	732	MTX	CM-N10-C14	2.67	124.19	119.57
3	A	732	MTX	C7-N8-C8A	2.67	119.38	116.69
2	В	604	UMP	O4-C4-N3	2.66	123.22	119.31
3	В	733	MTX	CT-CA-N	2.64	116.81	110.55
2	В	604	UMP	O4'-C1'-C2'	-2.50	101.52	106.25
2	A	603	UMP	C1'-N1-C2	2.47	122.51	117.64
3	В	733	MTX	C9-C6-N5	2.45	120.89	116.96
2	В	604	UMP	O4-C4-C5	-2.39	120.95	125.16
2	В	604	UMP	OP3-P-O5'	2.34	112.95	106.73
3	В	733	MTX	C2-N3-C4	2.29	123.27	116.72
2	A	603	UMP	O4-C4-N3	2.11	122.40	119.31
2	A	603	UMP	O5'-P-OP1	2.11	112.38	106.47
3	В	733	MTX	C15-C16-C11	-2.10	118.33	120.78
3	В	733	MTX	CB-CA-N	-2.06	106.72	110.88
2	A	603	UMP	C4'-O4'-C1'	2.03	114.36	109.45

There are no chirality outliers.

All (15) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
3	В	733	MTX	N-CA-CT-O1
3	В	733	MTX	N-CA-CT-O2
3	В	733	MTX	C6-C9-N10-CM
3	В	733	MTX	CT-CA-N-C
3	В	733	MTX	CB-CA-CT-O2
3	В	733	MTX	CB-CA-CT-O1
3	A	732	MTX	CT-CA-N-C
3	A	732	MTX	OE1-CD-CG-CB
3	A	732	MTX	OE2-CD-CG-CB
3	A	732	MTX	N-CA-CT-O2
3	A	732	MTX	CB-CA-N-C
2	A	603	UMP	O4'-C4'-C5'-O5'
2	В	604	UMP	O4'-C4'-C5'-O5'
3	В	733	MTX	OE1-CD-CG-CB
3	В	733	MTX	OE2-CD-CG-CB


There are no ring outliers.

3 monomers are involved in 3 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	732	MTX	1	0
2	В	604	UMP	1	0
2	A	603	UMP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.












Rings

Torsions



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

