

# wwPDB X-ray Structure Validation Summary Report (i)

May 16, 2020 – 10:20 am BST

PDB ID : 4AY4

Title : crystal structure of Bacillus anthracis PurE

Authors: Oliete, R.; Pous, J.; Rodriguez-Puente, S.; Abad-Zapatero, C.; Guasch, A.

Deposited on : 2012-06-18

Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13 EDS : 2.11

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac: 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001)

Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.11

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \text{Whole archive} \\ (\#\text{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar \; resolution} \\ (\#{\rm Entries, \; resolution \; range(\AA)}) \end{array}$
$R_{free}$	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	181	83%	• 14%
1	В	181	87%	•• 12%
1	С	181	84%	• 14%
1	D	181	86%	• 11%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-



ria:

N	/Iol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
	2	ACT	С	1158	-	=	X	=



# 2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 5089 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called N5-CARBOXYAMINOIMIDAZOLE RIBONUCLEOTIDE MUTASE.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	Λ	156	Total	С	N	О	S	0	0	0
1	A	150	1155	729	199	221	6	0	0	U
1	В	159	Total	С	N	О	S	9	0	0
1	Б	109	1178	743	202	227	6	2		
1	С	156	Total	С	N	О	S	4	2	0
1		150	1169	738	202	223	6	4	2	U
1	D	161	Total	С	N	О	S	K	1	0
1	ש	161	1202	757	206	232	7	5		U

There are 80 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	expression tag	UNP Q81ZH8
A	-18	GLY	-	expression tag	UNP Q81ZH8
A	-17	SER	-	expression tag	UNP Q81ZH8
A	-16	SER	-	expression tag	UNP Q81ZH8
A	-15	HIS	_	expression tag	UNP Q81ZH8
A	-14	HIS	-	expression tag	UNP Q81ZH8
A	-13	HIS	-	expression tag	UNP Q81ZH8
A	-12	HIS	-	expression tag	UNP Q81ZH8
A	-11	HIS	-	expression tag	UNP Q81ZH8
A	-10	HIS	-	expression tag	UNP Q81ZH8
A	-9	SER	-	expression tag	UNP Q81ZH8
A	-8	SER	_	expression tag	UNP Q81ZH8
A	-7	GLY	-	expression tag	UNP Q81ZH8
A	-6	LEU	-	expression tag	UNP Q81ZH8
A	-5	VAL	-	expression tag	UNP Q81ZH8
A	-4	PRO	-	expression tag	UNP Q81ZH8
A	-3	ARG		expression tag	UNP Q81ZH8
A	-2	GLY	=	expression tag	UNP Q81ZH8
A	-1	SER	-	expression tag	UNP Q81ZH8
A	0	HIS	_	expression tag	UNP Q81ZH8



Continued from previous page...


Chain	Residue	Modelled  Modelled	Actual	Comment	Reference
В	-19	MET	-	expression tag	UNP Q81ZH8
В	-18	GLY	_	expression tag	UNP Q81ZH8
В	-17	SER	-	expression tag	UNP Q81ZH8
В	-16	SER	-	expression tag	UNP Q81ZH8
В	-15	HIS	-	expression tag	UNP Q81ZH8
В	-14	HIS	_	expression tag	UNP Q81ZH8
В	-13	HIS	_	expression tag	UNP Q81ZH8
В	-12	HIS	-	expression tag	UNP Q81ZH8
В	-11	HIS	-	expression tag	UNP Q81ZH8
В	-10	HIS	-	expression tag	UNP Q81ZH8
В	-9	SER	-	expression tag	UNP Q81ZH8
В	-8	SER	-	expression tag	UNP Q81ZH8
В	-7	GLY	_	expression tag	UNP Q81ZH8
В	-6	LEU	-	expression tag	UNP Q81ZH8
В	-5	VAL	_	expression tag	UNP Q81ZH8
В	-4	PRO	-	expression tag	UNP Q81ZH8
В	-3	ARG	-	expression tag	UNP Q81ZH8
В	-2	GLY	_	expression tag	UNP Q81ZH8
В	-1	SER	_	expression tag	UNP Q81ZH8
В	0	HIS	-	expression tag	UNP Q81ZH8
С	-19	MET	_	expression tag	UNP Q81ZH8
С	-18	GLY	-	expression tag	UNP Q81ZH8
С	-17	SER	-	expression tag	UNP Q81ZH8
С	-16	SER	-	expression tag	UNP Q81ZH8
С	-15	HIS	-	expression tag	UNP Q81ZH8
С	-14	HIS	-	expression tag	UNP Q81ZH8
С	-13	HIS	-	expression tag	UNP Q81ZH8
С	-12	HIS	-	expression tag	UNP Q81ZH8
С	-11	HIS	-	expression tag	UNP Q81ZH8
С	-10	HIS	-	expression tag	UNP Q81ZH8
С	-9	SER	-	expression tag	UNP Q81ZH8
С	-8	SER	-	expression tag	UNP Q81ZH8
С	-7	GLY	-	expression tag	UNP Q81ZH8
С	-6	LEU	-	expression tag	UNP Q81ZH8
С	-5	VAL	-	expression tag	UNP Q81ZH8
С	-4	PRO	-	expression tag	UNP Q81ZH8
С	-3	ARG	-	expression tag	UNP Q81ZH8
С	-2	GLY	-	expression tag	UNP Q81ZH8
С	-1	SER	-	expression tag	UNP Q81ZH8
С	0	HIS	-	expression tag	UNP Q81ZH8
D	-19	MET	-	expression tag	UNP Q81ZH8
D	-18	GLY	-	expression tag	UNP Q81ZH8



 $Continued\ from\ previous\ page...$ 

Chain	Residue	Modelled	Actual	Comment	Reference
D	-17	SER	-	expression tag	UNP Q81ZH8
D	-16	SER	-	expression tag	UNP Q81ZH8
D	-15	HIS	_	expression tag	UNP Q81ZH8
D	-14	HIS	_	expression tag	UNP Q81ZH8
D	-13	HIS	_	expression tag	UNP Q81ZH8
D	-12	HIS	_	expression tag	UNP Q81ZH8
D	-11	HIS	_	expression tag	UNP Q81ZH8
D	-10	HIS	_	expression tag	UNP Q81ZH8
D	-9	SER	_	expression tag	UNP Q81ZH8
D	-8	SER	_	expression tag	UNP Q81ZH8
D	-7	GLY	_	expression tag	UNP Q81ZH8
D	-6	LEU	-	expression tag	UNP Q81ZH8
D	-5	VAL	-	expression tag	UNP Q81ZH8
D	-4	PRO	_	expression tag	UNP Q81ZH8
D	-3	ARG	-	expression tag	UNP Q81ZH8
D	-2	GLY	-	expression tag	UNP Q81ZH8
D	-1	SER	-	expression tag	UNP Q81ZH8
D	0	HIS	_	expression tag	UNP Q81ZH8

 $\bullet$  Molecule 2 is ACETATE ION (three-letter code: ACT) (formula:  $\mathrm{C_2H_3O_2}).$ 



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 4 2 2	0	0
2	В	1	Total C O 4 2 2	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	С	1	Total C O 4 2 2	0	0
2	D	1	Total C O 4 2 2	0	0
2	D	1	Total C O 4 2 2	0	0

#### • Molecule 3 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	108	Total O 108 108	0	0
3	В	82	Total O 82 82	0	0
3	С	91	Total O 91 91	0	0
3	D	84	Total O 84 84	0	0



## 3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: N5-CARBOXYAMINOIMIDAZOLE RIBONUCLEOTIDE MUTASE





# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants	87.00Å 87.00Å 270.00Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.00° 90.00° 120.00°	Depositor
Resolution (Å)	43.93 - 2.00	Depositor
resolution (A)	43.89 - 2.00	EDS
% Data completeness	96.7 (43.93-2.00)	Depositor
(in resolution range)	96.7 (43.89-2.00)	EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	6.26 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
P. P.	0.147 , 0.180	Depositor
$R, R_{free}$	0.161 , $0.192$	DCC
$R_{free}$ test set	2069 reflections $(5.08\%)$	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	22.0	Xtriage
Anisotropy	0.316	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.39, 45.7	EDS
L-test for twinning <sup>2</sup>	$ < L > = 0.48, < L^2> = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o$ , $F_c$ correlation	0.96	EDS
Total number of atoms	5089	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 8.65% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $< L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Clasira	Boı	nd lengths	Bond angles		
MIOI	Mol   Chain		# Z  > 5	RMSZ	# Z >5	
1	A	0.60	1/1173 (0.1%)	0.70	0/1589	
1	В	0.64	$2/1196 \ (0.2\%)$	0.67	0/1620	
1	С	0.63	1/1190 (0.1%)	0.70	0/1611	
1	D	0.59	$1/1220 \ (0.1\%)$	0.71	0/1651	
All	All	0.62	5/4779 (0.1%)	0.69	0/6471	

All (5) bond length outliers are listed below:

Mol	Chain	${f Res}$	Type	${f Atoms}$	$\mathbf{Z}$	${f Observed(\AA)}$	$\operatorname{Ideal}( ext{\AA})$
1	В	101	SER	CB-OG	-5.91	1.34	1.42
1	D	101	SER	CB-OG	-5.65	1.34	1.42
1	A	130	GLN	CD-NE2	-5.64	1.18	1.32
1	С	101	SER	CB-OG	-5.53	1.35	1.42
1	В	123	ASN	CG-ND2	-5.34	1.19	1.32

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

#### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1155	0	1178	3	0
1	В	1178	0	1200	1	0



$\alpha \cdots \tau$	r	•	
Continued	trom	nromanne	naae
$\circ$	110116	picolous	puyc

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	С	1169	0	1196	2	0
1	D	1202	0	1226	2	0
2	A	4	0	3	0	0
2	В	4	0	3	1	0
2	С	4	0	3	2	0
2	D	8	0	6	1	0
3	A	108	0	0	1	0
3	В	82	0	0	1	0
3	С	91	0	0	3	2
3	D	84	0	0	1	2
All	All	5089	0	4815	11	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 11 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$egin{array}{c}  ext{Interatomic} \  ext{distance} \ ( ext{Å}) \end{array}$	$egin{aligned}  ext{Clash} \  ext{overlap } ( ext{Å}) \end{aligned}$
2:C:1158:ACT:H1	3:C:2044:HOH:O	1.95	0.66
1:C:16:GLU:HB2	3:C:2017:HOH:O	2.10	0.52
2:D:1162:ACT:H1	3:D:2029:HOH:O	2.11	0.50
2:C:1158:ACT:H2	3:C:2030:HOH:O	2.11	0.48
1:D:47:PHE:O	1:D:51:GLU:HB2	2.13	0.48

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$egin{aligned} &  ext{Interatomic} \ &  ext{distance} \ &  ext{(Å)} \end{aligned}$	$egin{array}{c} { m Clash} \ { m overlap} \ ({ m \AA}) \end{array}$
3:C:2034:HOH:O	3:D:2034:HOH:O[8_455]	2.07	0.13
3:C:2036:HOH:O	3:D:2022:HOH:O[8_455]	2.08	0.12

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was



analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	$_{ m ntiles}$
1	A	154/181~(85%)	152 (99%)	2 (1%)	0	100	100
1	В	157/181 (87%)	155 (99%)	2 (1%)	0	100	100
1	С	156/181 (86%)	154 (99%)	2 (1%)	0	100	100
1	D	160/181 (88%)	158 (99%)	2 (1%)	0	100	100
All	All	627/724~(87%)	619 (99%)	8 (1%)	0	100	100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	A	121/143~(85%)	121 (100%)	0	100	100
1	В	124/143 (87%)	124 (100%)	0	100	100
1	С	123/143 (86%)	123 (100%)	0	100	100
1	D	127/143 (89%)	127 (100%)	0	100	100
All	All	495/572~(86%)	495 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	89	GLN
1	A	130	GLN
1	В	123	ASN
1	С	136	HIS
1	D	136	HIS



#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

#### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry (i)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trees	Chain	Res	Res Link	Bond lengths			Bond angles		
MIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	ACT	С	1158	-	1,3,3	0.78	0	0,3,3	0.00	-
2	ACT	D	1163	-	1,3,3	1.52	0	0,3,3	0.00	-
2	ACT	D	1162	-	1,3,3	0.96	0	0,3,3	0.00	-
2	ACT	A	1158	-	1,3,3	1.33	0	0,3,3	0.00	-
2	ACT	В	1161	-	1,3,3	0.38	0	0,3,3	0.00	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	С	1158	ACT	2	0



 $Continued\ from\ previous\ page...$ 

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	1162	ACT	1	0
2	В	1161	ACT	1	0

# 5.7 Other polymers (i)

There are no such residues in this entry.

# 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

#### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\#\mathrm{RSRZ}{>}2$	$OWAB(\AA^2)$	Q < 0.9
1	A	156/181~(86%)	-0.39	2 (1%) 77 76	15, 23, 41, 58	0
1	В	159/181 (87%)	-0.38	5 (3%) 49 48	15, 22, 44, 73	1 (0%)
1	С	156/181 (86%)	-0.43	1 (0%) 89 88	14, 21, 38, 58	1 (0%)
1	D	161/181 (88%)	-0.26	5 (3%) 49 48	15, 23, 44, 71	2 (1%)
All	All	632/724 (87%)	-0.37	13 (2%) 63 62	14, 22, 42, 73	4 (0%)

The worst 5 of 13 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	159	GLU	3.7
1	D	1	MET	3.0
1	D	159	GLU	3.0
1	В	135	PHE	2.8
1	В	158	SER	2.7

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

#### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	${f B-factors}({f \AA}^2)$	Q<0.9
2	ACT	С	1158	4/4	0.92	0.14	41,44,47,47	0
2	ACT	A	1158	4/4	0.93	0.18	41,43,43,44	0
2	ACT	D	1162	4/4	0.94	0.13	38,44,44,45	0
2	ACT	D	1163	4/4	0.96	0.10	42,43,45,49	0
2	ACT	В	1161	4/4	0.96	0.13	38,41,42,44	0

# 6.5 Other polymers (i)

There are no such residues in this entry.

