



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 2, 2023 – 09:37 AM EDT

PDB ID : 1B2M
Title : THREE-DIMENSIONAL STRUCTURE OF RIBONULCEASE T1 COM-
PLEXED WITH AN ISOSTERIC PHOSPHONATE ANALOGUE OF GPU:
ALTERNATE SUBSTRATE BINDING MODES AND CATALYSIS.
Authors : Arni, R.K.; Watanabe, L.; Ward, R.J.; Kreitman, R.J.; Kumar, K.; Walz Jr.,
F.G.
Deposited on : 1998-11-27
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.34


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RNA backbone	3102	1079 (2.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	C	2	100%
1	D	2	100%
1	E	2	100%
2	A	104	88% 12%
2	B	104	80% 18% .

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 1774 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called 5'-R(*GP*(U34))-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
1	C	2	Total 40	C 20	N 7	O 12	P 1	0	0	0
1	D	2	Total 40	C 20	N 7	O 12	P 1	0	0	0
1	E	2	Total 40	C 20	N 7	O 12	P 1	0	0	0

- Molecule 2 is a protein called RIBONUCLEASE T1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	A	104	Total 781	C 479	N 127	O 171	S 4	0	0	0
2	B	104	Total 781	C 479	N 127	O 171	S 4	0	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	C	1	Total 1	O 1	0	0
3	D	2	Total 2	O 2	0	0
3	A	48	Total 48	O 48	0	0
3	B	41	Total 41	O 41	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


Note EDS was not executed.

- Molecule 1: 5'-R(*GP*(U34))-3'

Chain C:  100%

G105
U106

- Molecule 1: 5'-R(*GP*(U34))-3'

Chain D:  100%


G105
U106

- Molecule 1: 5'-R(*GP*(U34))-3'

Chain E:  100%


G105
U106

- Molecule 2: RIBONUCLEASE T1

Chain A:  88% 12%

A1 C2 D3 S12 Q20 N43 N44 Y45 P55 I61 Y78 E82 L86 A95 V101 T104

- Molecule 2: RIBONUCLEASE T1

Chain B:  80% 18%

A1 S12 D15 Q20 H27 E28 D29 S35 N36 S37 Y38 Y42 N43 N44 S54 Y57 I61 V78 V79 N84 Q85 L86 A95 N98 V101 T104

4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	69.91Å 90.29Å 33.98Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	8.00 – 2.00	Depositor
% Data completeness (in resolution range)	75.0 (8.00-2.00)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.64	Depositor
Refinement program	X-PLOR 3.1	Depositor
R, R_{free}	0.187 , 0.254	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	1774	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: U34

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	C	1.93	0/22	3.62	3/33 (9.1%)
1	D	2.06	2/22 (9.1%)	3.39	3/33 (9.1%)
1	E	1.92	2/22 (9.1%)	3.50	3/33 (9.1%)
2	A	0.44	0/802	0.67	0/1093
2	B	0.49	0/802	0.69	0/1093
All	All	0.60	4/1670 (0.2%)	0.99	9/2285 (0.4%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	105	G	N9-C8	-5.89	1.33	1.37
1	D	105	G	C8-N7	-5.69	1.27	1.30
1	E	105	G	C8-N7	-5.28	1.27	1.30
1	E	105	G	N9-C8	-5.24	1.34	1.37

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	105	G	N7-C8-N9	13.90	120.05	113.10
1	D	105	G	N7-C8-N9	13.66	119.93	113.10
1	E	105	G	N7-C8-N9	13.64	119.92	113.10
1	C	105	G	C5-N7-C8	-9.45	99.58	104.30
1	C	105	G	C8-N9-C4	-9.11	102.76	106.40
1	E	105	G	C8-N9-C4	-8.84	102.86	106.40
1	D	105	G	C8-N9-C4	-8.70	102.92	106.40
1	E	105	G	C5-N7-C8	-8.12	100.24	104.30
1	D	105	G	C5-N7-C8	-7.09	100.76	104.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	40	0	23	0	0
1	D	40	0	23	0	0
1	E	40	0	23	0	0
2	A	781	0	673	7	0
2	B	781	0	673	10	0
3	A	48	0	0	1	0
3	B	41	0	0	0	0
3	C	1	0	0	0	0
3	D	2	0	0	0	0
All	All	1774	0	1415	17	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (17) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:20:GLN:HE21	2:B:84:ASN:ND2	1.83	0.76
2:B:20:GLN:HE21	2:B:84:ASN:HD22	1.36	0.72
2:B:61:ILE:HB	2:B:78:VAL:HG23	1.82	0.62
2:B:42:TYR:CE2	2:B:44:ASN:HA	2.34	0.61
2:A:95:ALA:HB2	2:A:101:VAL:HG13	1.83	0.60
2:A:3:ASP:HB2	2:A:12:SER:HA	1.87	0.56
2:A:20:GLN:HE22	2:A:86:LEU:HB2	1.73	0.54
2:A:55:PRO:HD2	2:A:82:GLU:HB2	1.89	0.53
2:A:43:ASN:HB2	2:A:45:TYR:CE1	2.44	0.52
2:B:36:ASN:HB2	2:B:38:TYR:CE2	2.44	0.52
2:A:1:ALA:HB1	3:A:178:HOH:O	2.13	0.47
2:B:95:ALA:HB2	2:B:101:VAL:HG13	1.98	0.46
2:B:98:ASN:HD22	2:B:98:ASN:HA	1.57	0.45
2:B:12:SER:O	2:B:15:ASP:HB2	2.18	0.44
2:B:27:HIS:CE1	2:B:57:TYR:CZ	3.05	0.44
2:A:61:ILE:HB	2:A:78:VAL:HG23	2.02	0.41
2:B:57:TYR:O	2:B:79:VAL:HA	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	A	102/104 (98%)	100 (98%)	2 (2%)	0	100	100
2	B	102/104 (98%)	99 (97%)	3 (3%)	0	100	100
All	All	204/208 (98%)	199 (98%)	5 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	A	85/85 (100%)	85 (100%)	0	100	100
2	B	85/85 (100%)	78 (92%)	7 (8%)	11	7
All	All	170/170 (100%)	163 (96%)	7 (4%)	30	28

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	28	GLU
2	B	29	ASP
2	B	35	SER
2	B	54	SER
2	B	84	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	86	LEU
2	B	98	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
2	A	20	GLN
2	A	43	ASN
2	B	27	HIS
2	B	36	ASN
2	B	84	ASN
2	B	98	ASN
2	B	99	ASN

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	C	0/2	-	-
1	D	0/2	-	-
1	E	0/2	-	-
All	All	0/6	-	-

There are no RNA backbone outliers to report.

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

3 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	U34	E	106	1	18,21,23	1.57	1 (5%)	26,30,35	1.32	4 (15%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	U34	C	106	1	18,21,23	1.06	1 (5%)	26,30,35	0.93	2 (7%)
1	U34	D	106	1	18,21,23	1.21	1 (5%)	26,30,35	1.21	2 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	U34	E	106	1	-	0/7/25/30	0/2/2/2
1	U34	C	106	1	-	0/7/25/30	0/2/2/2
1	U34	D	106	1	-	2/7/25/30	0/2/2/2

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	106	U34	C5'-C	-5.88	1.47	1.53
1	C	106	U34	C5'-C	-3.90	1.49	1.53
1	D	106	U34	C5'-C	-3.84	1.49	1.53

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	106	U34	C-C5'-C4'	4.67	120.99	112.47
1	E	106	U34	C5'-C4'-C3'	-2.91	109.79	115.48
1	E	106	U34	C-C5'-C4'	2.61	117.23	112.47
1	C	106	U34	C-C5'-C4'	2.53	117.09	112.47
1	D	106	U34	C4'-O4'-C1'	-2.46	104.05	109.47
1	E	106	U34	C1'-N1-C2	-2.39	113.24	117.57
1	C	106	U34	C5'-C4'-C3'	-2.26	111.07	115.48
1	E	106	U34	C1'-N1-C6	2.20	125.64	120.84

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	D	106	U34	C3'-C4'-C5'-C
1	D	106	U34	O4'-C4'-C5'-C

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.