



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 27, 2023 – 03:13 pm GMT

PDB ID : 8B32
Title : Chalcone synthase from *Hordeum vulgare* complexed with CoA
Authors : Zhang, L.; Groves, M.R.
Deposited on : 2022-09-15
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4298 (1.70-1.70)
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)
RSRZ outliers	127900	4222 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	417	 5% 86% 5% 8%
1	B	417	 8% 87% 5% 8%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 12364 atoms, of which 6008 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Chalcone synthase 2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	382	5908	1854	2976	511	546	21	83	3	0
1	B	382	5895	1851	2968	510	546	20	82	2	0

There are 36 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-17	MET	-	initiating methionine	UNP Q96562
A	-16	GLY	-	expression tag	UNP Q96562
A	-15	SER	-	expression tag	UNP Q96562
A	-14	SER	-	expression tag	UNP Q96562
A	-13	HIS	-	expression tag	UNP Q96562
A	-12	HIS	-	expression tag	UNP Q96562
A	-11	HIS	-	expression tag	UNP Q96562
A	-10	HIS	-	expression tag	UNP Q96562
A	-9	HIS	-	expression tag	UNP Q96562
A	-8	HIS	-	expression tag	UNP Q96562
A	-7	SER	-	expression tag	UNP Q96562
A	-6	GLN	-	expression tag	UNP Q96562
A	-5	ASP	-	expression tag	UNP Q96562
A	-4	PRO	-	expression tag	UNP Q96562
A	-3	ASN	-	expression tag	UNP Q96562
A	-2	SER	-	expression tag	UNP Q96562
A	-1	SER	-	expression tag	UNP Q96562
A	0	SER	-	expression tag	UNP Q96562
B	-17	MET	-	initiating methionine	UNP Q96562
B	-16	GLY	-	expression tag	UNP Q96562
B	-15	SER	-	expression tag	UNP Q96562
B	-14	SER	-	expression tag	UNP Q96562
B	-13	HIS	-	expression tag	UNP Q96562
B	-12	HIS	-	expression tag	UNP Q96562
B	-11	HIS	-	expression tag	UNP Q96562

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-10	HIS	-	expression tag	UNP Q96562
B	-9	HIS	-	expression tag	UNP Q96562
B	-8	HIS	-	expression tag	UNP Q96562
B	-7	SER	-	expression tag	UNP Q96562
B	-6	GLN	-	expression tag	UNP Q96562
B	-5	ASP	-	expression tag	UNP Q96562
B	-4	PRO	-	expression tag	UNP Q96562
B	-3	ASN	-	expression tag	UNP Q96562
B	-2	SER	-	expression tag	UNP Q96562
B	-1	SER	-	expression tag	UNP Q96562
B	0	SER	-	expression tag	UNP Q96562

- Molecule 2 is COENZYME A (three-letter code: COA) (formula: $C_{21}H_{36}N_7O_{16}P_3S$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms							ZeroOcc	AltConf
			Total	C	H	N	O	P	S		
2	A	1	80	21	32	7	16	3	1	3	0
2	B	1	80	21	32	7	16	3	1	3	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	216	Total	O	0	0
			216	216		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	185	Total 185	O 185	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Chalcone synthase 2



- Molecule 1: Chalcone synthase 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	77.81Å 98.96Å 138.11Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.83 – 1.70 45.79 – 1.70	Depositor EDS
% Data completeness (in resolution range)	99.7 (45.83-1.70) 99.7 (45.79-1.70)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.64 (at 1.70Å)	Xtrriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.177 , 0.205 0.186 , 0.209	Depositor DCC
R_{free} test set	5895 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	29.2	Xtrriage
Anisotropy	0.092	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 43.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	12364	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.95% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: COA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.84	1/2988 (0.0%)	0.86	3/4043 (0.1%)
1	B	0.86	1/2983 (0.0%)	0.90	4/4037 (0.1%)
All	All	0.85	2/5971 (0.0%)	0.88	7/8080 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	262	GLU	CD-OE2	8.44	1.34	1.25
1	A	262	GLU	CD-OE1	5.21	1.31	1.25

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	391	VAL	CA-C-O	-7.56	104.23	120.10
1	A	139	MET	CG-SD-CE	-5.84	90.86	100.20
1	B	268	HIS	CB-CA-C	5.33	121.07	110.40
1	A	268	HIS	CB-CA-C	5.29	120.99	110.40
1	A	391	VAL	CA-C-O	-5.26	109.05	120.10
1	B	158	ARG	NE-CZ-NH1	5.24	122.92	120.30
1	B	139	MET	CG-SD-CE	-5.00	92.20	100.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2932	2976	2959	17	0
1	B	2927	2968	2952	14	0
2	A	48	32	32	0	0
2	B	48	32	32	0	0
3	A	216	0	0	1	0
3	B	185	0	0	1	0
All	All	6356	6008	5975	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:10:ARG:HD2	1:B:18:LEU:HD23	1.58	0.83
1:B:14:ARG:HD3	3:B:517:HOH:O	1.84	0.78
1:A:14:ARG:HD3	3:A:519:HOH:O	1.84	0.77
1:B:20:THR:HG21	1:B:237:PRO:HB3	1.74	0.70
1:A:159:LEU:CD1	1:B:175:LEU:HD21	2.29	0.63
1:A:175:LEU:HD21	1:B:159:LEU:CD1	2.35	0.57
1:A:246:GLN:HG3	1:A:382:GLU:HG2	1.90	0.54
1:B:246:GLN:HG3	1:B:382:GLU:HG2	1.92	0.51
1:B:171:THR:OG1	1:B:382:GLU:OE2	2.26	0.50
1:A:159:LEU:HD11	1:B:175:LEU:HD21	1.94	0.50
1:A:171:THR:OG1	1:A:382:GLU:OE2	2.26	0.48
1:A:166:CYS:HB2	1:A:375:PHE:O	2.14	0.48
1:A:326[B]:MET:CE	1:A:330:ARG:CZ	2.92	0.47
1:B:166:CYS:HB2	1:B:375:PHE:O	2.14	0.47
1:A:312:LEU:HB3	1:A:326[B]:MET:CE	2.45	0.46
1:A:159:LEU:HD13	1:B:175:LEU:HD21	1.97	0.45
1:A:267:ILE:HG22	1:A:269:LEU:HG	2.00	0.43
1:A:163:GLN:HE21	1:B:139:MET:CE	2.31	0.43
1:A:175:LEU:HD21	1:B:159:LEU:HD13	1.99	0.42
1:A:326[B]:MET:HB2	1:A:326[B]:MET:HE2	1.71	0.42
1:A:290:GLU:N	1:A:291:PRO:CD	2.83	0.41
1:B:267:ILE:HG22	1:B:269:LEU:HG	2.03	0.41
1:B:290:GLU:N	1:B:291:PRO:CD	2.84	0.41
1:A:312:LEU:HB3	1:A:326[B]:MET:HE1	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	383/417 (92%)	372 (97%)	11 (3%)	0	100	100
1	B	382/417 (92%)	373 (98%)	9 (2%)	0	100	100
All	All	765/834 (92%)	745 (97%)	20 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	313/339 (92%)	312 (100%)	1 (0%)	92	89
1	B	312/339 (92%)	311 (100%)	1 (0%)	92	89
All	All	625/678 (92%)	623 (100%)	2 (0%)	92	89

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	352	ARG
1	B	352	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	163	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	163	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	COA	A	401	-	41,50,50	0.95	1 (2%)	52,75,75	1.27	5 (9%)
2	COA	B	401	-	41,50,50	0.82	1 (2%)	52,75,75	1.02	2 (3%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	COA	A	401	-	-	7/44/64/64	0/3/3/3
2	COA	B	401	-	-	9/44/64/64	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	COA	P3B-O3B	4.70	1.68	1.59
2	B	401	COA	P3B-O3B	3.41	1.65	1.59

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	COA	C7P-C6P-C5P	-5.09	103.88	112.36
2	A	401	COA	OAP-CAP-CBP	-2.99	103.20	110.25
2	B	401	COA	O4B-C1B-C2B	-2.76	102.89	106.93
2	B	401	COA	P2A-O3A-P1A	-2.56	124.04	132.83
2	A	401	COA	P2A-O3A-P1A	-2.38	124.66	132.83
2	A	401	COA	O9A-P3B-O8A	2.33	116.53	107.64
2	A	401	COA	O2A-P1A-O1A	2.15	122.84	112.24

There are no chirality outliers.

All (16) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	401	COA	C3B-O3B-P3B-O9A
2	A	401	COA	CAP-C9P-N8P-C7P
2	B	401	COA	C5B-O5B-P1A-O2A
2	B	401	COA	CCP-O6A-P2A-O5A
2	A	401	COA	O9P-C9P-N8P-C7P
2	A	401	COA	O5P-C5P-N4P-C3P
2	A	401	COA	C6P-C5P-N4P-C3P
2	B	401	COA	C5B-O5B-P1A-O3A
2	B	401	COA	CCP-O6A-P2A-O3A
2	B	401	COA	C5B-O5B-P1A-O1A
2	B	401	COA	CAP-CBP-CCP-O6A
2	B	401	COA	CDP-CBP-CCP-O6A
2	A	401	COA	P1A-O3A-P2A-O5A
2	B	401	COA	CEP-CBP-CCP-O6A
2	A	401	COA	P1A-O3A-P2A-O4A
2	B	401	COA	CCP-O6A-P2A-O4A

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will

also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	382/417 (91%)	0.15	22 (5%) 23 25	26, 31, 49, 81	0
1	B	382/417 (91%)	0.38	34 (8%) 9 11	23, 31, 52, 68	0
All	All	764/834 (91%)	0.27	56 (7%) 15 17	23, 31, 51, 81	0

All (56) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	205	LYS	4.9
1	A	233	LEU	4.5
1	B	51	LEU	4.1
1	B	52	ALA	3.9
1	B	171	THR	3.6
1	B	269	LEU	3.5
1	B	80	ILE	3.5
1	A	11	MET	3.4
1	B	271	LYS	3.3
1	B	208	LEU	3.2
1	B	233	LEU	3.2
1	A	166	CYS	3.0
1	A	132	CYS	3.0
1	A	356	ALA	3.0
1	B	167	PHE	3.0
1	B	232	GLN	2.9
1	B	175	LEU	2.9
1	B	190	VAL	2.8
1	B	206	SER	2.7
1	A	172	VAL	2.7
1	A	131	PHE	2.7
1	A	167	PHE	2.7
1	B	53	ASP	2.7
1	B	89	MET	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	192	CYS	2.6
1	A	165	GLY	2.6
1	B	168	GLY	2.6
1	B	270	LEU	2.6
1	B	204	CYS	2.6
1	B	169	GLY	2.5
1	B	162	TYR	2.5
1	B	130	VAL	2.4
1	B	48	SER	2.4
1	B	77	GLU	2.4
1	A	169	GLY	2.3
1	A	232	GLN	2.3
1	B	131	PHE	2.3
1	B	172	VAL	2.3
1	A	161	MET	2.3
1	A	16	GLU	2.2
1	A	133	THR	2.2
1	A	168	GLY	2.2
1	B	207	HIS	2.2
1	A	234	ASP	2.2
1	A	162	TYR	2.1
1	B	166	CYS	2.1
1	B	12	ALA	2.1
1	A	171	THR	2.1
1	B	192	CYS	2.1
1	B	54	LEU	2.1
1	A	170	ALA	2.1
1	B	132	CYS	2.1
1	A	357	LYS	2.0
1	B	81	LYS	2.0
1	B	161	MET	2.0
1	A	14	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	COA	B	401	48/48	0.77	0.24	46,58,78,86	3
2	COA	A	401	48/48	0.89	0.16	41,49,57,60	3

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.