

# wwPDB X-ray Structure Validation Summary Report (i)

### May 18, 2020 – 04:26 am BST

PDB ID : 2B6E

Title: X-Ray Crystal Structure of Protein HI1161 from Haemophilus influenzae.

Northeast Structural Genomics Consortium Target IR63.

Authors: Kuzin, A.P.; Benach, J.; Chen, Y.; Acton, T.; Xiao, R.; Conover, K.; Ma,

L.-C.; Kellie, R.; Cunningham, K.E.; Montelione, G.; Hunt, J.F.; Tong, L.;

Northeast Structural Genomics Consortium (NESG)

Deposited on : 2005-10-01

Resolution : 1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp

with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul: 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13 EDS : 2.11

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.11

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.


Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \text{Whole archive} \\ (\#\text{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries,\ resolution\ range(\AA)}) \end{array}$
$R_{free}$	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	146	77%	16%	6%
1	В	146	77%	16%	• 5%
1	С	146	79%	16%	5%
1	D	146	7%	21%	)
1	Е	146	79%	14%	• 6%
1	F	146	75%	18%	• 5%





 $Continued\ from\ previous\ page...$ 

Mol	Chain	Length	Quality of chair	n	
1	G	146	13%	26%	• 5%
1	Н	146	13% 77%	16%	6%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	ACY	В	602	-	-	X	-
2	ACY	С	601	-	-	X	-
2	ACY	D	604	-	-	X	-
2	ACY	F	607	-	-	X	-
2	ACY	Н	608	-	-	X	-



# 2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 9023 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Hypothetical UPF0152 protein HI1161.

Mol	Chain	Residues			Atom	.S			ZeroOcc	AltConf	Trace
1	A	137	Total	С	N	О	S	Se	0	0	0
1	A	137	1045	655	187	196	4	3	0	U	0
1	В	139	Total	С	N	О	S	Se	0	0	0
1	Ъ	159	1061	667	189	198	4	3	0	U	U
1	С	139	Total	С	N	О	S	Se	0	0	0
1		109	1061	667	189	198	4	3	0	U	
1	D	146	Total	С	N	О	S	Se	0	0	0
1	D	140	1130	708	208	207	4	3	0		U
1	E	137	Total	С	N	Ο	S	Se	0	0	0
1	L	157	1045	655	187	196	4	3	O	U	
1	F	138	Total	С	N	Ο	S	Se	0	0	0
1	I.	130	1053	661	188	197	4	3	U	U	U
1	G	138	Total	С	Ν	О	S	Se	0	0	0
1	G	100	1053	661	188	197	4	3	0	0	U
1	Н	137	Total	С	N	О	S	Se	0	0	0
1	11	191	1045	655	187	196	4	3			

There are 88 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	MET	MODIFIED RESIDUE	UNP P45083
A	40	MSE	MET	MODIFIED RESIDUE	UNP P45083
A	47	MSE	MET	MODIFIED RESIDUE	UNP P45083
A	139	LEU	_	CLONING ARTIFACT	UNP P45083
A	140	GLU	_	CLONING ARTIFACT	UNP P45083
A	141	HIS	-	EXPRESSION TAG	UNP P45083
A	142	HIS	_	EXPRESSION TAG	UNP P45083
A	143	HIS	_	EXPRESSION TAG	UNP P45083
A	144	HIS	-	EXPRESSION TAG	UNP P45083
A	145	HIS	_	EXPRESSION TAG	UNP P45083
A	146	HIS	_	EXPRESSION TAG	UNP P45083
В	1	MSE	MET	MODIFIED RESIDUE	UNP P45083
В	40	MSE	MET	MODIFIED RESIDUE	UNP P45083



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
В	47	MSE	MET	MODIFIED RESIDUE	UNP P45083
В	139	LEU	_	CLONING ARTIFACT	UNP P45083
В	140	GLU	-	CLONING ARTIFACT	UNP P45083
В	141	HIS	_	EXPRESSION TAG	UNP P45083
В	142	HIS	_	EXPRESSION TAG	UNP P45083
В	143	HIS	_	EXPRESSION TAG	UNP P45083
В	144	HIS	-	EXPRESSION TAG	UNP P45083
В	145	HIS	-	EXPRESSION TAG	UNP P45083
В	146	HIS	-	EXPRESSION TAG	UNP P45083
С	1	MSE	MET	MODIFIED RESIDUE	UNP P45083
С	40	MSE	MET	MODIFIED RESIDUE	UNP P45083
С	47	MSE	MET	MODIFIED RESIDUE	UNP P45083
С	139	LEU	-	CLONING ARTIFACT	UNP P45083
С	140	GLU	-	CLONING ARTIFACT	UNP P45083
С	141	HIS	-	EXPRESSION TAG	UNP P45083
С	142	HIS	-	EXPRESSION TAG	UNP P45083
С	143	HIS	-	EXPRESSION TAG	UNP P45083
С	144	HIS	_	EXPRESSION TAG	UNP P45083
С	145	HIS	-	EXPRESSION TAG	UNP P45083
С	146	HIS	-	EXPRESSION TAG	UNP P45083
D	1	MSE	MET	MODIFIED RESIDUE	UNP P45083
D	40	MSE	MET	MODIFIED RESIDUE	UNP P45083
D	47	MSE	MET	MODIFIED RESIDUE	UNP P45083
D	139	LEU	-	CLONING ARTIFACT	UNP P45083
D	140	GLU	-	CLONING ARTIFACT	UNP P45083
D	141	HIS	-	EXPRESSION TAG	UNP P45083
D	142	HIS	_	EXPRESSION TAG	UNP P45083
D	143	HIS	-	EXPRESSION TAG	UNP P45083
D	144	HIS	-	EXPRESSION TAG	UNP P45083
D	145	HIS	_	EXPRESSION TAG	UNP P45083
D	146	HIS	=	EXPRESSION TAG	UNP P45083
Е	1	MSE	MET	MODIFIED RESIDUE	UNP P45083
Е	40	MSE	MET	MODIFIED RESIDUE	UNP P45083
Е	47	MSE	MET	MODIFIED RESIDUE	UNP P45083
Е	139	LEU		CLONING ARTIFACT	UNP P45083
Е	140	GLU	-	CLONING ARTIFACT	UNP P45083
Е	141	HIS	-	EXPRESSION TAG	UNP P45083
Е	142	HIS	-	EXPRESSION TAG	UNP P45083
Е	143	HIS		EXPRESSION TAG	UNP P45083
Е	144	HIS	-	EXPRESSION TAG	UNP P45083
Е	145	HIS		EXPRESSION TAG	UNP P45083
Е	146	HIS	_	EXPRESSION TAG	UNP P45083




 $Continued\ from\ previous\ page...$ 

Chain	Residue	Modelled	Actual	Comment	Reference
F	1	MSE	MET	MODIFIED RESIDUE	UNP P45083
F	40	MSE	MET	MODIFIED RESIDUE	UNP P45083
F	47	MSE	MET	MODIFIED RESIDUE	UNP P45083
F	139	LEU	_	CLONING ARTIFACT	UNP P45083
F	140	GLU	1	CLONING ARTIFACT	UNP P45083
F	141	HIS	_	EXPRESSION TAG	UNP P45083
F	142	HIS	_	EXPRESSION TAG	UNP P45083
F	143	HIS	-	EXPRESSION TAG	UNP P45083
F	144	HIS	-	EXPRESSION TAG	UNP P45083
F	145	HIS	-	EXPRESSION TAG	UNP P45083
F	146	HIS	-	EXPRESSION TAG	UNP P45083
G	1	MSE	MET	MODIFIED RESIDUE	UNP P45083
G	40	MSE	MET	MODIFIED RESIDUE	UNP P45083
G	47	MSE	MET	MODIFIED RESIDUE	UNP P45083
G	139	LEU	-	CLONING ARTIFACT	UNP P45083
G	140	GLU	-	CLONING ARTIFACT	UNP P45083
G	141	HIS	1	EXPRESSION TAG	UNP P45083
G	142	HIS	-	EXPRESSION TAG	UNP P45083
G	143	HIS	-	EXPRESSION TAG	UNP P45083
G	144	HIS	-	EXPRESSION TAG	UNP P45083
G	145	HIS	-	EXPRESSION TAG	UNP P45083
G	146	HIS	=	EXPRESSION TAG	UNP P45083
Н	1	MSE	MET	MODIFIED RESIDUE	UNP P45083
Н	40	MSE	MET	MODIFIED RESIDUE	UNP P45083
Н	47	MSE	MET	MODIFIED RESIDUE	UNP P45083
Н	139	LEU	-	CLONING ARTIFACT	UNP P45083
Н	140	GLU	=	CLONING ARTIFACT	UNP P45083
Н	141	HIS	-	EXPRESSION TAG	UNP P45083
Н	142	HIS	=	EXPRESSION TAG	UNP P45083
Н	143	HIS	=	EXPRESSION TAG	UNP P45083
Н	144	HIS	-	EXPRESSION TAG	UNP P45083
Н	145	HIS	-	EXPRESSION TAG	UNP P45083
Н	146	HIS	ı	EXPRESSION TAG	UNP P45083

 $\bullet$  Molecule 2 is ACETIC ACID (three-letter code: ACY) (formula:  $\mathrm{C_2H_4O_2}).$ 





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 4 2 2	0	0
2	В	1	Total C O 4 2 2	0	0
2	С	1	Total C O 4 2 2	0	0
2	D	1	Total C O 4 2 2	0	0
2	D	1	Total C O 4 2 2	0	0
2	E	1	Total C O 4 2 2	0	0
2	F	1	Total C O 4 2 2	0	0
2	Н	1	Total C O 4 2 2	0	0

### • Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	86	Total O 86 86	0	0
3	В	79	Total O 79 79	0	0
3	С	68	Total O 68 68	0	0
3	D	96	Total O 96 96	0	0



 $Continued\ from\ previous\ page...$ 


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	Е	66	Total O 66 66	0	0
3	F	55	Total O 55 55	0	0
3	G	22	Total O 22 22	0	0
3	Н	26	Total O 26 26	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Hypothetical UPF0152 protein HI1161









• Molecule 1: Hypothetical UPF0152 protein HI1161



• Molecule 1: Hypothetical UPF0152 protein HI1161



• Molecule 1: Hypothetical UPF0152 protein HI1161





# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	60.65Å 102.88Å 101.36Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.00° 98.36° 90.00°	Depositor
Resolution (Å)	29.97 - 1.90	Depositor
Resolution (A)	29.97 - 1.90	EDS
% Data completeness	87.9 (29.97-1.90)	Depositor
(in resolution range)	98.0 (29.97-1.90)	EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	4.08 (at 1.91Å)	Xtriage
Refinement program	CNS 1.1	Depositor
D D.	0.191 , 0.220	Depositor
$R, R_{free}$	0.196 , $0.225$	DCC
$R_{free}$ test set	9298 reflections $(4.94\%)$	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	18.0	Xtriage
Anisotropy	0.173	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.39, 59.2	EDS
L-test for twinning <sup>2</sup>	$ < L >=0.49, < L^2>=0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	9023	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.04% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $< L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ACY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
MIOI	Chain	RMSZ	# Z >5	RMSZ	# Z  > 5	
1	A	0.31	0/1059	0.67	0/1434	
1	В	0.31	0/1075	0.65	0/1456	
1	С	0.30	0/1075	0.64	0/1456	
1	D	0.30	0/1150	0.63	0/1558	
1	Е	0.30	0/1059	0.61	0/1434	
1	F	0.29	0/1067	0.62	0/1445	
1	G	0.28	0/1067	0.56	0/1445	
1	Н	0.27	0/1059	0.56	0/1434	
All	All	0.30	0/8611	0.62	0/11662	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	Α	1045	0	1066	22	0
1	В	1061	0	1088	19	0
1	С	1061	0	1088	21	0
1	D	1130	0	1136	20	0
1	Ε	1045	0	1066	20	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	F	1053	0	1077	18	0
1	G	1053	0	1077	30	0
1	Н	1045	0	1066	17	0
2	A	4	0	3	1	0
2	В	4	0	3	3	0
2	С	4	0	3	4	0
2	D	8	0	6	5	0
2	Ε	4	0	3	1	0
2	F	4	0	3	2	0
2	Н	4	0	3	2	0
3	A	86	0	0	4	0
3	В	79	0	0	2	0
3	С	68	0	0	2	0
3	D	96	0	0	1	0
3	E	66	0	0	1	0
3	F	55	0	0	0	0
3	G	22	0	0	2	0
3	Н	26	0	0	3	0
All	All	9023	0	8688	158	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 158 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1 Atom-2		$egin{aligned}  ext{Interatomic} \  ext{distance} & ( ext{Å}) \end{aligned}$	$egin{aligned}  ext{Clash} \  ext{overlap } ( ext{Å}) \end{aligned}$
1:B:115:GLN:HE21	2:C:601:ACY:H1	1.18	1.03
2:B:602:ACY:H1	1:C:115:GLN:HE21	1.25	1.00
2:B:602:ACY:H1	1:C:115:GLN:NE2	1.98	0.79
1:B:115:GLN:NE2	2:C:601:ACY:H1	1.98	0.78
1:E:94:ARG:HE	1:E:94:ARG:H	1.30	0.77

There are no symmetry-related clashes.

## 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	$_{ m ntiles}$
1	A	135/146~(92%)	135 (100%)	0	0	100	100
1	В	137/146 (94%)	136 (99%)	1 (1%)	0	100	100
1	С	137/146 (94%)	135 (98%)	2 (2%)	0	100	100
1	D	144/146 (99%)	141 (98%)	3 (2%)	0	100	100
1	E	135/146 (92%)	133 (98%)	2 (2%)	0	100	100
1	F	136/146 (93%)	133 (98%)	3 (2%)	0	100	100
1	G	136/146 (93%)	132 (97%)	3 (2%)	1 (1%)	22	12
1	Н	135/146~(92%)	131 (97%)	4 (3%)	0	100	100
All	All	1095/1168 (94%)	1076 (98%)	18 (2%)	1 (0%)	51	43

#### All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	G	2	LEU

### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	A	117/123~(95%)	115 (98%)	2 (2%)	60	57
1	В	$119/123 \ (97\%)$	115 (97%)	4 (3%)	37	28
1	С	119/123 (97%)	116 (98%)	3 (2%)	47	41
1	D	$126/123 \; (102\%)$	124 (98%)	2 (2%)	62	60
1	E	117/123 (95%)	115 (98%)	2 (2%)	60	57
1	F	118/123~(96%)	115 (98%)	3 (2%)	47	41
1	G	118/123 (96%)	114 (97%)	4 (3%)	37	28
1	Н	117/123 (95%)	116 (99%)	1 (1%)	78	79
All	All	951/984 (97%)	930 (98%)	21 (2%)	52	47



5 of 21 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	83	LEU
1	E	83	LEU
1	G	75	GLU
1	С	94	ARG
1	G	83	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 41 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	11	ASN
1	D	112	GLN
1	Н	11	ASN
1	D	44	HIS
1	D	89	HIS

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry (i)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mol	l Type Chain Res		Res Link		В	Bond lengths			Bond angles		
MIOI	туре	Chain	nes	res   Lilik	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z  > 2	
2	ACY	F	607	_	1,3,3	1.60	0	0,3,3	0.00	-	
2	ACY	Н	608	-	1,3,3	1.36	0	0,3,3	0.00	-	
2	ACY	A	603	_	1,3,3	2.00	0	0,3,3	0.00	-	
2	ACY	D	604	-	1,3,3	1.20	0	0,3,3	0.00	-	
2	ACY	D	606	_	1,3,3	1.47	0	0,3,3	0.00	-	
2	ACY	Е	605	-	1,3,3	1.18	0	0,3,3	0.00	-	
2	ACY	В	602	-	1,3,3	0.86	0	0,3,3	0.00	-	
2	ACY	С	601	-	1,3,3	0.85	0	0,3,3	0.00	-	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

7 monomers are involved in 17 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	F	607	ACY	2	0
2	Н	608	ACY	2	0
2	A	603	ACY	1	0
2	D	604	ACY	5	0
2	Е	605	ACY	1	0
2	В	602	ACY	3	0
2	С	601	ACY	4	0

## 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\#\mathrm{RSRZ}{>}2$	$OWAB(\AA^2)$	Q < 0.9
1	A	134/146 (91%)	0.04	1 (0%) 87 88	8, 15, 40, 67	0
1	В	136/146 (93%)	-0.09	2 (1%) 73 76	8, 14, 36, 53	0
1	С	136/146 (93%)	0.17	5 (3%) 41 44	9, 16, 46, 69	0
1	D	143/146 (97%)	0.21	10 (6%) 16 18	7, 16, 48, 78	0
1	E	134/146 (91%)	0.15	6 (4%) 33 36	12, 21, 48, 61	0
1	F	135/146~(92%)	0.22	8 (5%) 22 25	13, 24, 50, 69	0
1	G	135/146~(92%)	0.71	19 (14%) 2 2	15, 28, 59, 73	0
1	Н	134/146 (91%)	0.82	19 (14%) 2 2	16, 31, 57, 71	0
All	All	$1087/1168 \; (93\%)$	0.28	70 (6%) 19 22	7, 21, 51, 78	0

The worst 5 of 70 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	Н	50	PHE	6.1
1	G	137	ASN	6.1
1	Н	77	GLY	5.2
1	С	139	LEU	4.9
1	G	138	LEU	4.7

## 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates (i)

There are no carbohydrates in this entry.



### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
2	ACY	D	606	4/4	0.69	0.20	26,26,26,28	0
2	ACY	Н	608	4/4	0.72	0.20	28,30,31,32	0
2	ACY	A	603	4/4	0.76	0.35	24,25,27,30	0
2	ACY	Е	605	4/4	0.82	0.12	15,20,22,23	0
2	ACY	D	604	4/4	0.83	0.13	10,14,14,17	0
2	ACY	В	602	4/4	0.88	0.10	12,15,17,19	0
2	ACY	F	607	4/4	0.89	0.11	24,25,27,27	0
2	ACY	С	601	4/4	0.89	0.15	13,16,18,19	0

### 6.5 Other polymers (i)

There are no such residues in this entry.

