Full wwPDB X-ray Structure Validation Report (i)

Jul 13, 2022 – 04:02 PM EDT

PDB ID	:	6B8K
Title	:	Crystal structure of Human galectin-3 CRD in complex with Lactulose
Authors	:	Kishor, C.; Blanchard, H.
Deposited on	:	2017-10-09
Resolution	:	1.28 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.29
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.29

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.28 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1850 (1.30-1.26)
Clashscore	141614	1926 (1.30-1.26)
Ramachandran outliers	138981	1860 (1.30-1.26)
Sidechain outliers	138945	1859 (1.30-1.26)
RSRZ outliers	127900	1807 (1.30-1.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	139	% 80% 18% ••
2	B	2	100%

2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 1415 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Galectin-3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	138	1215	780	221	211	3	0	18	0

- Molecule 2 is an oligosaccharide called beta-D-galactopyranose-(1-4)-beta-D-fructofuranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
			Total	C	O			
2	B	2	23	12	11	0	0	0

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Cl		
3	A	3	3	3	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	174	174	174	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Galectin-3

Chain A:

- Molecule 2: beta-D-galactopyranose-(1-4)-beta-D-fructofuranose

Chain B:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α, β, γ	36.17Å 57.87Å 62.60Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.67 – 1.28 30.67 – 1.28	Depositor EDS
% Data completeness (in resolution range)	99.5 (30.67-1.28) 99.5 (30.67-1.28)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	4.01 (at 1.28Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, R_{free}	0.163 , 0.195 0.168 , 0.196	Depositor DCC
R_{free} test set	1706 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å^2)	11.0	Xtriage
Anisotropy	0.086	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.34 , 41.9	EDS
L-test for twinning[2]	$< \|L\| > = 0.47$, $< L^2 > = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.97	EDS
Total number of atoms	1415	wwPDB-VP
Average B, all atoms (Å^2)	12.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.08% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, FRU, GAL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	1.41	7/1290 (0.5%)	1.43	21/1743 (1.2%)

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	247	TYR	CE1-CZ	-8.91	1.26	1.38
1	A	184	GLU	CD-OE2	-7.25	1.17	1.25
1	A	224	ARG	CZ-NH2	5.86	1.40	1.33
1	A	205	GLU	CD-OE2	5.56	1.31	1.25
1	A	184	GLU	CD-OE1	5.53	1.31	1.25
1	A	244[A]	SER	CA-CB	-5.14	1.45	1.52
1	A	244[B]	SER	CA-CB	-5.14	1.45	1.52

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	129[A]	ARG	NE-CZ-NH2	-9.06	115.77	120.30
1	A	129[B]	ARG	NE-CZ-NH2	-9.06	115.77	120.30
1	A	224	ARG	NE-CZ-NH1	7.97	124.28	120.30
1	A	151	ARG	NE-CZ-NH1	7.95	124.28	120.30
1	A	129[A]	ARG	NE-CZ-NH1	7.92	124.26	120.30
1	A	129[B]	ARG	NE-CZ-NH1	7.92	124.26	120.30
1	A	186	ARG	NE-CZ-NH2	7.91	124.25	120.30
1	A	168[A]	ARG	NE-CZ-NH1	7.74	124.17	120.30
1	A	168[B]	ARG	NE-CZ-NH1	7.74	124.17	120.30
1	A	118	TYR	CB-CG-CD1	-7.30	116.62	121.00
1	A	144	ARG	NE-CZ-NH2	-6.63	116.98	120.30
1	A	184	GLU	OE1-CD-OE2	6.42	131.00	123.30
1	A	247	TYR	CD1-CE1-CZ	6.31	125.48	119.80
1	A	224	ARG	NE-CZ-NH2	-6.15	117.22	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	186	ARG	NE-CZ-NH1	-5.90	117.35	120.30
1	A	169[A]	ARG	NE-CZ-NH2	-5.71	117.45	120.30
1	A	169[B]	ARG	NE-CZ-NH2	-5.71	117.45	120.30
1	A	168[A]	ARG	NE-CZ-NH2	-5.59	117.50	120.30
1	A	168[B]	ARG	NE-CZ-NH2	-5.59	117.50	120.30
1	A	183	ARG	NE-CZ-NH1	5.43	123.01	120.30
1	A	151	ARG	NE-CZ-NH2	-5.12	117.74	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1215	0	1273	21	0
2	B	23	0	20	0	0
3	A	3	0	0	0	0
4	A	174	0	0	8	0
All	All	1415	0	1293	21	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:208[B]:HIS:CE1	1:A:222[B]:ASN:OD1	1.84	1.30
1:A:208[B]:HIS:ND1	1:A:222[B]:ASN:OD1	1.80	1.14
1:A:119[B]:ASN:OD1	4:A:405:HOH:O	1.79	1.00
1:A:113:PRO:HB3	4:A:547:HOH:O	1.76	0.84
1:A:208[B]:HIS:HE1	1:A:222[B]:ASN:OD1	1.65	0.80
1:A:208[A]:HIS:HD2	4:A:401:HOH:O	1.84	0.61
1:A:113:PRO:HD2	4:A:408:HOH:O	2.02	0.59
1:A:113:PRO:N	4:A:408:HOH:O	2.35	0.58
1:A:113:PRO:CD	4:A:408:HOH:O	2.51	0.58

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:205:GLU:OE1	1:A:208[A]:HIS:HE1	1.89	0.56
1:A:134[B]:ILE:HD12	1:A:236:ILE:CD1	2.38	0.54
1:A:176[A]:LYS:HE2	1:A:179:ASN:HA	1.90	0.53
1:A:208[B]:HIS:CE1	1:A:220[B]:GLN:NE2	2.79	0.51
1:A:205:GLU:OE1	1:A:208[A]:HIS:CE1	2.65	0.49
1:A:158:HIS:O	1:A:173:CYS:HA	2.15	0.46
1:A:196[B]:LYS:NZ	4:A:404:HOH:O	2.49	0.46
1:A:134[B]:ILE:HD12	1:A:236:ILE:HD11	1.98	0.45
1:A:134[B]:ILE:HD12	1:A:236:ILE:HD13	1.99	0.45
1:A:208[B]:HIS:CE1	1:A:220[B]:GLN:HE22	2.35	0.45
1:A:150[B]:GLN:OE1	1:A:233[B]:LYS:HG3	2.19	0.43
1:A:208[B]:HIS:CD2	4:A:403:HOH:O	2.74	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	154/139 (111%)	150 (97%)	4 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	142/124 (114%)	141 (99%)	1 (1%)	84 60

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	113	PRO

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	201	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	FRU	B	1	2	11,12,12	3.56	7 (63%)	10,18,18	4.57	7 (70%)
2	GAL	B	2	2	11,11,12	2.34	6 (54%)	15,15,17	1.99	4 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FRU	B	1	2	-	2/5/24/24	0/1/1/1
2	GAL	B	2	2	-	0/2/19/22	0/1/1/1

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1	FRU	O5-C2	7.22	1.54	1.43
2	B	1	FRU	C6-C5	-5.46	1.33	1.51
2	B	1	FRU	C4-C5	4.57	1.64	1.53
2	B	2	GAL	C1-C2	-3.81	1.43	1.52
2	B	1	FRU	O5-C5	3.70	1.51	1.43
2	B	1	FRU	C1-C2	-3.45	1.46	1.52
2	B	2	GAL	O2-C2	3.14	1.50	1.43
2	B	2	GAL	C2-C3	3.14	1.57	1.52
2	B	2	GAL	O5-C1	2.72	1.48	1.43
2	B	2	GAL	O5-C5	2.67	1.48	1.43
2	B	1	FRU	O4-C4	2.43	1.48	1.43
2	B	2	GAL	C4-C5	-2.14	1.48	1.53
2	B	1	FRU	C4-C3	-2.04	1.44	1.52

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	B	1	FRU	O5-C5-C6	9.91	136.44	108.85
2	B	1	FRU	O6-C6-C5	-6.56	88.79	111.29
2	B	1	FRU	C6-C5-C4	-5.54	101.73	115.09
2	B	1	FRU	C5-C4-C3	-4.57	87.24	101.91
2	B	2	GAL	O5-C5-C6	-4.25	100.55	107.20
2	B	2	GAL	C1-O5-C5	-3.60	107.31	112.19
2	B	2	GAL	C1-C2-C3	3.33	113.76	109.67
2	B	2	GAL	O2-C2-C3	-2.70	104.74	110.14
2	B	1	FRU	O5-C5-C4	2.15	110.84	105.49
2	B	1	FRU	O1-C1-C2	-2.11	107.39	111.86
2	B	1	FRU	O4-C4-C5	2.06	117.00	111.05

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	1	FRU	O5-C5-C6-O6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	B	1	FRU	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.

Oligosaccharide Chain B



5.6 Ligand geometry (i)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	A	138/139 (99%)	-0.08	1 (0%) 87 84	6, 10, 17, 30	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	113	PRO	5.3

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	FRU	B	1	12/12	0.88	0.11	15,22,28,28	0
2	GAL	B	2	11/12	0.95	0.06	10,12,15,19	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain B:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)



6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
3	CL	A	302	1/1	0.91	0.14	37,37,37,37	0
3	CL	A	303	1/1	0.98	0.05	21,21,21,21	0
3	CL	A	301	1/1	0.99	0.06	10,10,10,10	0

6.5 Other polymers (i)

There are no such residues in this entry.