



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 26, 2023 – 05:57 AM EDT

PDB ID : 6B9J  
Title : Structure of vaccinia virus D8 protein bound to human Fab vv138  
Authors : Zajonc, D.M.  
Deposited on : 2017-10-10  
Resolution : 2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 2.35.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1957 (2.90-2.90)
Clashscore	141614	2172 (2.90-2.90)
Ramachandran outliers	138981	2115 (2.90-2.90)
Sidechain outliers	138945	2117 (2.90-2.90)
RSRZ outliers	127900	1906 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	X	241	 93% • 5%
1	Y	241	 92% • 5%
2	A	217	 94% • •
2	H	217	 94% • •
3	B	215	 90% 8% ••

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	L	215	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	NA	X	302	-	-	-	X
5	NA	Y	302	-	-	-	X

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 9990 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called IMV membrane protein.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
1	X	228	1847	1196	300	351	0	1	0
1	Y	228	1858	1203	302	353	0	1	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
X	236	HIS	-	expression tag	UNP Q1M1K6
X	237	HIS	-	expression tag	UNP Q1M1K6
X	238	HIS	-	expression tag	UNP Q1M1K6
X	239	HIS	-	expression tag	UNP Q1M1K6
X	240	HIS	-	expression tag	UNP Q1M1K6
X	241	HIS	-	expression tag	UNP Q1M1K6
Y	236	HIS	-	expression tag	UNP Q1M1K6
Y	237	HIS	-	expression tag	UNP Q1M1K6
Y	238	HIS	-	expression tag	UNP Q1M1K6
Y	239	HIS	-	expression tag	UNP Q1M1K6
Y	240	HIS	-	expression tag	UNP Q1M1K6
Y	241	HIS	-	expression tag	UNP Q1M1K6


- Molecule 2 is a protein called Fab vv138 Heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	H	212	1584	1005	254	318	7	0	0	0
2	A	212	1556	984	247	318	7	0	0	0

- Molecule 3 is a protein called Fab vv138 Light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	L	209	Total	C	N	O	S	0	0	0
			1577	985	265	323	4			
3	B	213	Total	C	N	O	S	0	0	0
			1501	931	256	310	4			

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	X	1	Total	C	O	0	0
			6	3	3		
4	Y	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	X	3	Total	Na	0	0
			3	3		
5	H	2	Total	Na	0	0
			2	2		
5	Y	2	Total	Na	0	0
			2	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	X	14	Total O 14 14	0	0
6	H	9	Total O 9 9	0	0
6	L	5	Total O 5 5	0	0
6	Y	9	Total O 9 9	0	0
6	A	7	Total O 7 7	0	0
6	B	4	Total O 4 4	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: IMV membrane protein




- Molecule 1: IMV membrane protein





- Molecule 2: Fab vv138 Heavy chain




- Molecule 2: Fab vv138 Heavy chain






- Molecule 3: Fab vv138 Light chain





- Molecule 3: Fab vv138 Light chain





## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	234.28Å 253.84Å 73.76Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	117.14 – 2.90 117.14 – 2.90	Depositor EDS
% Data completeness (in resolution range)	99.2 (117.14-2.90) 99.2 (117.14-2.90)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.16	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.72 (at 2.91Å)	Xtrriage
Refinement program	REFMAC 5.8.0135	Depositor
R, $R_{free}$	0.241 , 0.269 0.242 , 0.265	Depositor DCC
$R_{free}$ test set	2466 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	74.1	Xtrriage
Anisotropy	0.067	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 73.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	9990	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	90.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.83% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	X	0.39	0/1900	0.60	0/2586
1	Y	0.39	0/1912	0.60	0/2600
2	A	0.38	0/1595	0.59	0/2185
2	H	0.38	0/1623	0.59	0/2216
3	B	0.42	0/1531	0.62	0/2095
3	L	0.41	0/1611	0.62	0/2192
All	All	0.39	0/10172	0.60	0/13874

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	X	1847	0	1780	2	0
1	Y	1858	0	1803	2	0
2	A	1556	0	1454	5	0
2	H	1584	0	1527	4	0
3	B	1501	0	1346	9	0
3	L	1577	0	1482	11	0
4	X	6	0	8	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	Y	6	0	8	0	0
5	H	2	0	0	0	0
5	X	3	0	0	0	0
5	Y	2	0	0	0	0
6	A	7	0	0	0	0
6	B	4	0	0	0	0
6	H	9	0	0	0	0
6	L	5	0	0	0	0
6	X	14	0	0	0	0
6	Y	9	0	0	0	0
All	All	9990	0	9408	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:L:136:CYS:HG	3:L:196:CYS:HG	0.94	0.87
3:B:136:CYS:HG	3:B:196:CYS:CB	1.95	0.78
3:L:136:CYS:HG	3:L:196:CYS:CB	1.97	0.77
3:B:126:GLN:O	3:B:129:SER:OG	2.05	0.75
3:L:126:GLN:O	3:L:129:SER:OG	2.03	0.74
2:H:200:CYS:SG	2:H:213:LYS:HB3	2.31	0.71
2:A:200:CYS:SG	2:A:213:LYS:HB3	2.32	0.69
2:H:144:CYS:HG	2:H:200:CYS:CB	2.06	0.65
2:H:144:CYS:SG	2:H:200:CYS:CB	2.85	0.65
2:A:144:CYS:SG	2:A:200:CYS:CB	2.86	0.64
3:L:149:GLN:HG2	3:L:156:LEU:HD11	1.86	0.58
3:L:149:GLN:CG	3:L:156:LEU:HD11	2.34	0.56
3:B:150:TRP:CB	3:B:181:LEU:HD11	2.35	0.56
3:L:139:ASN:HD22	3:L:140:ASN:N	2.05	0.55
2:H:144:CYS:SG	2:H:200:CYS:HB3	2.48	0.54
3:B:139:ASN:HD22	3:B:140:ASN:N	2.06	0.54
2:A:144:CYS:SG	2:A:200:CYS:HB3	2.48	0.54
3:L:55:ARG:NH1	3:L:63:PHE:O	2.41	0.53
3:B:55:ARG:NH1	3:B:63:PHE:O	2.42	0.53
1:Y:78:SER:O	1:Y:86:LYS:HE2	2.14	0.48
1:X:78:SER:O	1:X:86:LYS:HE2	2.14	0.47
3:L:136:CYS:HB2	3:L:150:TRP:CZ2	2.50	0.46
2:A:103:ILE:HD11	3:B:50:TYR:HB3	1.98	0.45

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:X:22:LYS:HB3	1:X:23:PRO:CD	2.47	0.45
1:Y:22:LYS:HB3	1:Y:23:PRO:CD	2.48	0.43
3:B:137:LEU:HD12	3:B:138:LEU:N	2.34	0.43
3:L:25:ALA:HB3	3:L:70:THR:HG22	2.01	0.42
3:L:6:GLN:HG2	3:L:89:CYS:SG	2.60	0.41
3:L:137:LEU:HD12	3:L:138:LEU:N	2.35	0.41
3:B:6:GLN:HG2	3:B:89:CYS:SG	2.60	0.41
3:B:25:ALA:HB3	3:B:70:THR:HG22	2.02	0.41
2:A:144:CYS:HG	2:A:200:CYS:HG	0.40	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	X	225/241 (93%)	211 (94%)	13 (6%)	1 (0%)	34	66
1	Y	225/241 (93%)	211 (94%)	13 (6%)	1 (0%)	34	66
2	A	208/217 (96%)	192 (92%)	16 (8%)	0	100	100
2	H	208/217 (96%)	194 (93%)	14 (7%)	0	100	100
3	B	209/215 (97%)	194 (93%)	15 (7%)	0	100	100
3	L	203/215 (94%)	194 (96%)	9 (4%)	0	100	100
All	All	1278/1346 (95%)	1196 (94%)	80 (6%)	2 (0%)	47	78

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	X	84	VAL
1	Y	84	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	X	204/222 (92%)	204 (100%)	0	100	100
1	Y	208/222 (94%)	206 (99%)	2 (1%)	76	92
2	A	170/185 (92%)	165 (97%)	5 (3%)	42	76
2	H	177/185 (96%)	172 (97%)	5 (3%)	43	76
3	B	153/187 (82%)	149 (97%)	4 (3%)	46	77
3	L	173/187 (92%)	169 (98%)	4 (2%)	50	80
All	All	1085/1188 (91%)	1065 (98%)	20 (2%)	59	85

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	H	63	LYS
2	H	116	SER
2	H	139	THR
2	H	148	ASP
2	H	174	LEU
3	L	70	THR
3	L	139	ASN
3	L	187	ASP
3	L	201	GLN
1	Y	13	LYS
1	Y	213	HIS
2	A	85	SER
2	A	116	SER
2	A	139	THR
2	A	148	ASP
2	A	174	LEU
3	B	24	ARG
3	B	70	THR
3	B	139	ASN
3	B	187	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such

sidechains are listed below:

Mol	Chain	Res	Type
1	X	9	ASN
1	X	59	ASN
2	H	168	HIS
2	H	175	GLN
3	L	139	ASN
1	Y	9	ASN
1	Y	59	ASN
2	A	168	HIS
3	B	139	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 7 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	GOL	X	301	-	5,5,5	0.63	0	5,5,5	0.56	0
4	GOL	Y	301	-	5,5,5	0.72	0	5,5,5	0.93	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	X	301	-	-	2/4/4/4	-
4	GOL	Y	301	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	X	301	GOL	O1-C1-C2-C3
4	X	301	GOL	O1-C1-C2-O2
4	Y	301	GOL	O1-C1-C2-C3
4	Y	301	GOL	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	X	228/241 (94%)	0.45	3 (1%) 77 77	39, 60, 96, 152	0
1	Y	228/241 (94%)	0.45	2 (0%) 84 84	43, 60, 96, 150	0
2	A	212/217 (97%)	0.41	9 (4%) 36 32	47, 101, 138, 158	0
2	H	212/217 (97%)	0.55	11 (5%) 27 23	47, 94, 129, 144	0
3	B	213/215 (99%)	0.83	21 (9%) 7 5	74, 125, 159, 180	0
3	L	209/215 (97%)	0.93	25 (11%) 4 3	73, 108, 137, 154	0
All	All	1302/1346 (96%)	0.60	71 (5%) 25 21	39, 93, 141, 180	0

All (71) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	B	138	LEU	4.2
2	H	100	SER	3.9
3	B	161	SER	3.9
2	H	188	VAL	3.7
2	H	138	GLY	3.6
3	B	196	CYS	3.4
3	B	135	VAL	3.3
3	B	150	TRP	3.3
3	L	176	SER	3.3
3	L	175	TYR	3.1
2	H	164	THR	3.0
3	L	204	SER	3.0
3	L	137	LEU	2.9
3	L	170	SER	2.9
2	H	185	VAL	2.8
3	L	199	THR	2.8
3	L	115	PRO	2.8
3	B	120	PHE	2.8
2	H	215	VAL	2.7

*Continued on next page...*



*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
3	B	159	GLY	2.7
3	B	115	PRO	2.7
3	B	95	SER	2.7
3	L	203	LEU	2.7
3	L	147	LYS	2.6
3	B	48	LEU	2.5
3	L	150	TRP	2.5
3	L	120	PHE	2.5
2	H	142	LEU	2.5
3	L	206	PRO	2.5
1	Y	213	HIS	2.5
2	A	188	VAL	2.5
2	H	170	PHE	2.4
3	L	25	ALA	2.4
3	L	138	LEU	2.4
3	B	104	THR	2.4
2	A	169	THR	2.4
3	B	146	ALA	2.3
3	L	134	VAL	2.3
3	L	93	GLY	2.3
1	X	20	ARG	2.3
3	B	203	LEU	2.3
3	B	2	ILE	2.3
3	L	129	SER	2.3
3	L	198	VAL	2.3
3	B	30	THR	2.3
3	B	207	VAL	2.3
3	B	177	LEU	2.3
3	L	135	VAL	2.2
3	L	164	SER	2.2
3	L	78	ARG	2.2
3	L	185	LYS	2.2
2	H	45	LEU	2.2
2	A	186	VAL	2.2
2	A	18	VAL	2.2
2	H	186	VAL	2.1
2	A	163	LEU	2.1
2	A	158	TRP	2.1
1	X	79	ASN	2.1
2	A	215	VAL	2.1
1	X	202	LEU	2.1
2	H	193	LEU	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	A	128	LEU	2.1
3	B	137	LEU	2.1
3	B	85	ALA	2.1
2	A	100	SER	2.1
3	L	117	VAL	2.1
3	B	94	SER	2.1
1	Y	20	ARG	2.1
3	B	148	VAL	2.0
3	L	179	SER	2.0
3	L	201	GLN	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	GOL	Y	301	6/6	0.54	0.32	60,69,71,71	0
5	NA	X	303	1/1	0.66	0.28	78,78,78,78	0
5	NA	X	302	1/1	0.70	0.47	73,73,73,73	0
5	NA	Y	302	1/1	0.72	0.44	70,70,70,70	0
5	NA	X	304	1/1	0.77	0.21	79,79,79,79	0
4	GOL	X	301	6/6	0.81	0.24	55,64,65,68	0
5	NA	H	302	1/1	0.85	0.17	76,76,76,76	0
5	NA	Y	303	1/1	0.86	0.33	70,70,70,70	0
5	NA	H	301	1/1	0.94	0.20	72,72,72,72	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.