



Full wwPDB EM Validation Report ⓘ

Dec 25, 2022 – 06:37 am GMT

PDB ID : 8BBE
EMDB ID : EMD-15954
Title : Structure of the IFT-A complex; IFT-A2 module
Authors : Hesketh, S.J.; Mukhopadhyay, A.G.; Nakamura, D.; Toropova, K.; Roberts, A.J.
Deposited on : 2022-10-12
Resolution : 3.50 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev43
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.9
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.31.3

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	C	1241	
2	D	1500	
3	E	1184	
4	F	209	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 20181 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Intraflagellar transport protein 122 homolog.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	C	702	5595	3569	951	1040	35	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	208	UNK	ASN	conflict	UNP Q9HBG6
C	209	UNK	ARG	conflict	UNP Q9HBG6
C	210	UNK	TYR	conflict	UNP Q9HBG6
C	211	UNK	ILE	conflict	UNP Q9HBG6
C	212	UNK	GLN	conflict	UNP Q9HBG6
C	213	UNK	GLU	conflict	UNP Q9HBG6

- Molecule 2 is a protein called SNAP-tag,Tetratricopeptide repeat protein 21B.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	D	648	5253	3318	916	978	41	0	0

- Molecule 3 is a protein called WD repeat-containing protein 35.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
3	E	1112	8863	5657	1497	1650	59	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	-2	PHE	-	expression tag	UNP Q9P2L0
E	-1	GLN	-	expression tag	UNP Q9P2L0
E	0	GLY	-	expression tag	UNP Q9P2L0

- Molecule 4 is a protein called Intraflagellar transport protein 43 homolog.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
4	F	57	470	296	74	98	2	0	0


There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
F	0	GLY	-	expression tag	UNP Q96FT9

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Intraflagellar transport protein 122 homolog





• Molecule 3: WD repeat-containing protein 35






• Molecule 4: Intraflagellar transport protein 43 homolog



GLY MET
GLU
GLY TRP ASP
LEU LEU LEU ASP
SER LEU LEU ASP
VAL ARG VAL GLU
LYS GLU ALA
SER LEU LEU ARG
LYS ARG TYR
SER ARG SER
LEU LEU ALA
LYS THR
SER ARG
GLU ARG
ALA
GLN
GLU
SER
LEU
ALA
GLN
SER
ASP
HIS
LEU
ASN
GLY
ILE
SER
LEU
THR

GLN
GLY
TRP
ALA
GLY
ASP
SER
VAL
LYS
ALA
SER
LYS
PHE
ARG
SER
ARG
LYS
ALA
SER
GLU
GLU
LEU
ILE
GLU
MET
ASP
GLY
PHE
ARG
ARG
LEU
ARG
PRO
GLN
GLU
SER
LEU
SER
GLY
TYR
HIS
ASN
GLY
ASN
ILE
ILE
PRO
PRO
ASP
LEU
GLU
GLU
VAL
GLN
SER
GLU
ASP
PHE
VAL
LEU
GLN
VAL
ALA



ALA
PRO
SER
ILE
ILE
LYS
ARG
VAL
M130
T131
Y132
R133
D134
L135
D136
N137
D138
L139
M140
K141
Y142
S143
A144
I145
Q146
T147
L148
D149
G150
E151
I152
D153
L154
K155
L156
L157
T158
K159
V160
L161
A162
P163
E164
H165
E166
V167
R168
E169
D170
D171
V172
G173
W174
D175
W176
L179
F180

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	242645	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	48	Depositor
Minimum defocus (nm)	1500	Depositor
Maximum defocus (nm)	3500	Depositor
Magnification	Not provided	
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor
Maximum map value	0.701	Depositor
Minimum map value	-0.291	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.009	Depositor
Recommended contour level	0.16	Depositor
Map size (Å)	546.304, 546.304, 546.304	wwPDB
Map dimensions	512, 512, 512	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.067, 1.067, 1.067	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	C	0.65	0/5679	1.04	19/7662 (0.2%)
2	D	0.67	0/5347	0.98	21/7203 (0.3%)
3	E	0.65	0/9049	1.00	31/12229 (0.3%)
4	F	0.67	0/479	1.04	2/650 (0.3%)
All	All	0.66	0/20554	1.01	73/27744 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	6
2	D	0	1
3	E	0	3
All	All	0	10

There are no bond length outliers.

All (73) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	61	TYR	CB-CG-CD2	-10.31	114.81	121.00
3	E	669	ARG	NE-CZ-NH2	9.34	124.97	120.30
3	E	938	ARG	NE-CZ-NH2	8.64	124.62	120.30
2	D	997	ARG	NE-CZ-NH2	8.57	124.58	120.30
3	E	1122	ARG	NE-CZ-NH2	8.37	124.48	120.30
1	C	189	ARG	NE-CZ-NH2	8.26	124.43	120.30
2	D	1044	ARG	NE-CZ-NH2	7.98	124.29	120.30
2	D	672	ARG	NE-CZ-NH2	7.79	124.19	120.30
2	D	1007	ARG	NE-CZ-NH2	7.70	124.15	120.30
3	E	521	ARG	NE-CZ-NH2	7.63	124.11	120.30
1	C	250	ARG	NE-CZ-NH2	7.55	124.08	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	F	168	ARG	NE-CZ-NH2	7.48	124.04	120.30
1	C	684	ARG	NE-CZ-NH2	7.27	123.94	120.30
2	D	981	ARG	NE-CZ-NH2	7.22	123.91	120.30
3	E	593	ARG	NE-CZ-NH2	7.17	123.89	120.30
1	C	159	ARG	NE-CZ-NH2	7.13	123.86	120.30
3	E	1038	ARG	NE-CZ-NH2	7.03	123.82	120.30
2	D	1228	ARG	NE-CZ-NH2	6.97	123.78	120.30
1	C	647	ARG	NE-CZ-NH2	6.96	123.78	120.30
1	C	453	ARG	NE-CZ-NH2	6.91	123.76	120.30
2	D	1175	ARG	NE-CZ-NH2	6.83	123.72	120.30
3	E	1079	ARG	NE-CZ-NH2	6.79	123.70	120.30
3	E	669	ARG	NE-CZ-NH1	-6.78	116.91	120.30
3	E	353	ARG	NE-CZ-NH2	6.60	123.60	120.30
2	D	1050	ARG	NE-CZ-NH2	6.49	123.55	120.30
1	C	587	ARG	NE-CZ-NH2	6.46	123.53	120.30
2	D	832	ARG	NE-CZ-NH2	6.45	123.53	120.30
1	C	171	ARG	NE-CZ-NH2	6.41	123.51	120.30
3	E	721	ARG	NE-CZ-NH2	6.34	123.47	120.30
1	C	384	ARG	NE-CZ-NH2	6.33	123.47	120.30
3	E	700	ARG	NE-CZ-NH2	6.13	123.36	120.30
3	E	350	ARG	NE-CZ-NH2	6.12	123.36	120.30
3	E	868	ARG	NE-CZ-NH2	6.11	123.35	120.30
3	E	240	ARG	NE-CZ-NH2	6.07	123.34	120.30
3	E	672	ARG	NE-CZ-NH2	6.05	123.33	120.30
2	D	1015	ARG	NE-CZ-NH2	5.84	123.22	120.30
3	E	119	ARG	NE-CZ-NH2	5.84	123.22	120.30
2	D	691	ARG	NE-CZ-NH2	5.83	123.22	120.30
2	D	723	ARG	NE-CZ-NH2	5.75	123.17	120.30
3	E	1025	ARG	NE-CZ-NH2	5.74	123.17	120.30
3	E	594	ARG	NE-CZ-NH2	5.71	123.16	120.30
2	D	898	ARG	NE-CZ-NH2	5.70	123.15	120.30
1	C	474	ARG	NE-CZ-NH2	5.69	123.14	120.30
3	E	966	ARG	NE-CZ-NH2	5.65	123.12	120.30
2	D	1281	ARG	NE-CZ-NH2	5.60	123.10	120.30
3	E	774	ARG	NE-CZ-NH2	5.60	123.10	120.30
1	C	518	ARG	NE-CZ-NH2	5.59	123.10	120.30
3	E	528	ARG	NE-CZ-NH2	5.58	123.09	120.30
3	E	766	ARG	NE-CZ-NH2	5.49	123.04	120.30
1	C	197	ARG	NE-CZ-NH2	5.48	123.04	120.30
3	E	813	ARG	NE-CZ-NH2	5.47	123.03	120.30
1	C	465	ARG	NE-CZ-NH2	5.43	123.02	120.30
4	F	142	TYR	CB-CG-CD2	-5.42	117.75	121.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	425	ARG	NE-CZ-NH2	5.37	122.98	120.30
2	D	858	ARG	NE-CZ-NH2	5.30	122.95	120.30
2	D	1053	ARG	NE-CZ-NH2	5.28	122.94	120.30
1	C	673	ARG	NE-CZ-NH2	5.27	122.93	120.30
3	E	148	ARG	NE-CZ-NH2	5.26	122.93	120.30
3	E	620	PHE	CB-CG-CD2	-5.23	117.14	120.80
2	D	718	ARG	NE-CZ-NH2	5.19	122.90	120.30
3	E	1018	ARG	NE-CZ-NH2	5.17	122.89	120.30
1	C	526	ARG	NE-CZ-NH2	5.15	122.88	120.30
1	C	61	TYR	CA-CB-CG	5.15	123.19	113.40
2	D	1311	ARG	NE-CZ-NH2	5.14	122.87	120.30
3	E	767	ARG	NE-CZ-NH2	5.10	122.85	120.30
1	C	362	ARG	NE-CZ-NH2	5.09	122.85	120.30
3	E	547	TYR	CB-CG-CD2	-5.07	117.96	121.00
3	E	1038	ARG	NE-CZ-NH1	-5.06	117.77	120.30
2	D	1177	ARG	NE-CZ-NH2	5.06	122.83	120.30
3	E	733	ARG	NE-CZ-NH2	5.06	122.83	120.30
2	D	926	ARG	NE-CZ-NH2	5.04	122.82	120.30
3	E	616	ARG	NE-CZ-NH2	5.02	122.81	120.30
2	D	998	ARG	NE-CZ-NH2	5.01	122.81	120.30

There are no chirality outliers.

All (10) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	288	TYR	Sidechain
1	C	453	ARG	Sidechain
1	C	465	ARG	Sidechain
1	C	61	TYR	Sidechain
1	C	726	TYR	Sidechain
1	C	90	TYR	Sidechain
2	D	732	TYR	Sidechain
3	E	1122	ARG	Sidechain
3	E	339	TYR	Sidechain
3	E	733	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	5595	0	5567	2	0
2	D	5253	0	5275	1	0
3	E	8863	0	8810	4	0
4	F	470	0	441	0	0
All	All	20181	0	20093	7	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All (7) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:E:567:LEU:HB2	3:E:596:VAL:HG21	1.93	0.49
3:E:309:LEU:HD11	3:E:318:ILE:HD11	1.97	0.46
3:E:1113:PHE:CG	3:E:1118:SER:HB3	2.53	0.44
1:C:169:ILE:HD13	1:C:266:TYR:CD1	2.53	0.42
2:D:937:CYS:SG	2:D:961:LEU:HD11	2.59	0.42
1:C:576:LEU:C	1:C:577:ASN:HD22	2.23	0.41
3:E:368:LYS:HZ1	3:E:656:GLU:CD	2.23	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	692/1241 (56%)	659 (95%)	33 (5%)	0	100	100
2	D	646/1500 (43%)	623 (96%)	23 (4%)	0	100	100
3	E	1102/1184 (93%)	1053 (96%)	49 (4%)	0	100	100
4	F	55/209 (26%)	55 (100%)	0	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	2495/4134 (60%)	2390 (96%)	105 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	610/1085 (56%)	609 (100%)	1 (0%)	93	98
2	D	556/1279 (44%)	556 (100%)	0	100	100
3	E	961/1020 (94%)	961 (100%)	0	100	100
4	F	53/181 (29%)	53 (100%)	0	100	100
All	All	2180/3565 (61%)	2179 (100%)	1 (0%)	100	100

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	597	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
3	E	1116	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-15954. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)

6.1.1 Primary map



6.1.2 Raw map




The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 256



Y Index: 256




Z Index: 256


6.2.2 Raw map



X Index: 256



Y Index: 256



Z Index: 256

The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)

6.3.1 Primary map



X Index: 254




Y Index: 235




Z Index: 262


6.3.2 Raw map



X Index: 254



Y Index: 301



Z Index: 221


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal surface views [i](#)


6.4.1 Primary map



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.16. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.4.2 Raw map



X



Y



Z

These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.5 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.5.1 emd_15954_msk_1.map [i](#)



X



Y




Z

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 140 nm³; this corresponds to an approximate mass of 126 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum i




*Reported resolution corresponds to spatial frequency of 0.286 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.286 Å⁻¹

8.2 Resolution estimates


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.50	-	-
Author-provided FSC curve	3.70	4.18	3.74
Unmasked-calculated*	4.34	8.80	4.43

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 4.34 differs from the reported value 3.5 by more than 10 %

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-15954 and PDB model 8BBE. Per-residue inclusion information can be found in section 3 on page 5.

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.16 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.16).











9.4 Atom inclusion [i](#)



At the recommended contour level, 72% of all backbone atoms, 56% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary [i](#)

The table lists the average atom inclusion at the recommended contour level (0.16) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.5591	 0.3060
C	 0.7012	 0.4050
D	 0.3672	 0.1580
E	 0.6000	 0.3390
F	 0.2338	 0.1710

