



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 22, 2020 – 08:32 AM BST

PDB ID : 4BER  
Title : Crystal structure of the Legionella pneumophila FIC domain-containing effector AnkX protein in complex with cytidine monophosphate  
Authors : Campanacci, V.; Mukherjee, S.; Roy, C.R.; Cherfils, J.  
Deposited on : 2013-03-12  
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	484	<p>4% 77% 18% ..</p>
1	B	484	<p>% 81% 15% ..</p>

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 7755 atoms, of which 36 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called PHOSPHOCHOLINE TRANSFERASE ANKX.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	475	Total 3789	C 2429	N 633	O 710	S 17	0	0	0
1	B	475	Total 3789	C 2428	N 633	O 711	S 17	0	0	0

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	expression tag	UNP Q5ZXN6
A	247	PRO	LEU	engineered mutation	UNP Q5ZXN6
B	1	GLY	-	expression tag	UNP Q5ZXN6
B	247	PRO	LEU	engineered mutation	UNP Q5ZXN6

- Molecule 2 is CYTIDINE-5'-MONOPHOSPHATE (three-letter code: C5P) (formula:  $C_9H_{14}N_3O_8P$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
2	A	1	Total	C	H	N	O	P	0	0
			35	9	14	3	8	1		
2	B	1	Total	C	H	N	O	P	0	0
			35	9	14	3	8	1		

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	O P	0	0
			5	4 1		
3	A	1	Total	O P	0	0
			5	4 1		
3	B	1	Total	O P	0	0
			5	4 1		
3	B	1	Total	O P	0	0
			5	4 1		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	H	O	0	0
			14	3	8	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	36	Total	O	0	0
			36	36		
5	B	37	Total	O	0	0
			37	37		


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: PHOSPHOCHOLINE TRANSFERASE ANKX



- Molecule 1: PHOSPHOCHOLINE TRANSFERASE ANKX



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	58.33Å 103.56Å 224.10Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.28 – 2.60 46.28 – 2.60	Depositor EDS
% Data completeness (in resolution range)	99.8 (46.28-2.60) 99.7 (46.28-2.60)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.36 (at 2.61Å)	Xtrriage
Refinement program	BUSTER 2.10.0	Depositor
R, $R_{free}$	0.206 , 0.239 0.212 , 0.245	Depositor DCC
$R_{free}$ test set	2136 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	51.3	Xtrriage
Anisotropy	0.579	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 47.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	7755	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	66.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.60% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, PO4, C5P

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.49	0/3874	0.70	0/5232
1	B	0.49	0/3875	0.71	2/5234 (0.0%)
All	All	0.49	0/7749	0.71	2/10466 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	429	LEU	CA-CB-CG	5.97	129.04	115.30
1	B	429	LEU	N-CA-CB	5.63	121.66	110.40

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3789	0	3773	42	0
1	B	3789	0	3772	43	0
2	A	21	14	12	1	0
2	B	21	14	12	0	0
3	A	10	0	0	0	0
3	B	10	0	0	0	0
4	A	6	8	8	1	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	36	0	0	0	0
5	B	37	0	0	0	0
All	All	7719	36	7577	84	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (84) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:103:GLU:N	1:A:103:GLU:OE1	1.88	1.05
1:A:260:GLU:OE1	1:A:262:ASN:HB2	1.85	0.75
1:B:437:ASN:HB3	1:B:440:ILE:HG12	1.75	0.68
1:B:59:ILE:HG12	1:B:76:LEU:HD11	1.77	0.67
1:A:374:ASN:H	1:A:377:GLN:HE21	1.42	0.67
1:B:228:LEU:O	1:B:230:PRO:HD3	1.95	0.66
1:B:374:ASN:H	1:B:377:GLN:HE21	1.43	0.66
1:A:33:SER:HB3	1:A:34:LYS:HE2	1.78	0.66
1:B:245:ASN:HD21	1:B:256:ALA:H	1.44	0.66
1:A:228:LEU:O	1:A:230:PRO:HD3	1.95	0.66
1:A:245:ASN:HD21	1:A:256:ALA:H	1.44	0.66
1:A:336:LEU:O	1:A:340:THR:HG23	1.96	0.66
1:B:427:THR:OG1	1:B:429:LEU:HD12	1.96	0.64
1:A:217:ILE:O	1:A:221:HIS:HD2	1.80	0.64
1:B:217:ILE:O	1:B:221:HIS:HD2	1.79	0.64
1:A:59:ILE:HD11	1:A:246:ILE:HD13	1.83	0.60
1:B:405:MET:HE3	1:B:408:LEU:HD23	1.84	0.60
1:B:14:LEU:HB2	1:B:249:MET:CE	2.33	0.58
1:A:427:THR:H	1:A:430:HIS:CD2	2.21	0.57
1:B:336:LEU:O	1:B:340:THR:HG23	2.06	0.56
1:B:14:LEU:HB2	1:B:249:MET:HE3	1.88	0.56
1:B:427:THR:H	1:B:430:HIS:CD2	2.24	0.56
1:A:102:ASP:N	1:A:103:GLU:OE1	2.41	0.54
1:A:405:MET:CE	1:A:443:LYS:HD2	2.40	0.52
1:A:2:VAL:HG22	1:B:4:ILE:HG12	1.91	0.51
1:A:263:VAL:HG11	1:A:388:PRO:HG3	1.93	0.51
1:A:188:LYS:HD3	1:A:189:ASN:H	1.75	0.51
1:B:467:THR:HG22	1:B:470:HIS:H	1.75	0.51
1:A:108:GLY:HA3	1:A:179:THR:HG21	1.93	0.51
1:B:108:GLY:HA3	1:B:179:THR:HG21	1.93	0.50
1:A:3:LYS:HE3	1:A:4:ILE:H	1.77	0.50

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:263:VAL:O	1:A:267:TYR:HB2	2.11	0.50
1:B:237:ARG:HA	1:B:241:ASN:HD22	1.77	0.49
1:B:263:VAL:O	1:B:267:TYR:HB2	2.12	0.49
1:B:282:ASN:O	1:B:286:ILE:HG12	2.14	0.48
1:B:405:MET:CE	1:B:443:LYS:HD2	2.43	0.48
1:A:405:MET:HE3	1:A:408:LEU:HD23	1.94	0.48
1:A:245:ASN:ND2	1:A:256:ALA:H	2.11	0.48
1:A:188:LYS:HD3	1:A:189:ASN:N	2.28	0.48
1:A:174:TYR:CG	1:A:180:ASN:HB2	2.49	0.47
1:A:237:ARG:HA	1:A:241:ASN:HD22	1.79	0.47
1:A:29:GLY:HA2	1:A:32:TRP:CE2	2.49	0.47
1:A:134:GLU:HB2	1:A:184:LEU:HB3	1.96	0.47
1:A:282:ASN:O	1:A:286:ILE:HG12	2.15	0.47
1:A:44:ARG:NH1	2:A:1000:C5P:O2	2.47	0.47
1:A:467:THR:HG22	1:A:470:HIS:H	1.79	0.47
1:B:134:GLU:HB2	1:B:184:LEU:HB3	1.96	0.47
1:A:427:THR:H	1:A:430:HIS:HD2	1.60	0.46
1:B:174:TYR:CG	1:B:180:ASN:HB2	2.49	0.46
1:B:29:GLY:HA2	1:B:32:TRP:CE2	2.50	0.46
1:B:245:ASN:O	1:B:249:MET:HG3	2.15	0.46
1:B:42:GLU:HG2	1:B:49:LEU:HD12	1.98	0.46
1:A:259:TYR:H	4:A:3000:GOL:H12	1.81	0.46
1:A:211:LEU:HG	1:A:339:LYS:HE2	1.98	0.46
1:A:42:GLU:HG2	1:A:49:LEU:HD12	1.97	0.45
1:B:245:ASN:ND2	1:B:256:ALA:H	2.11	0.45
1:B:405:MET:HE1	1:B:443:LYS:HD2	1.98	0.44
1:A:405:MET:HE1	1:A:443:LYS:HD2	1.99	0.44
1:A:101:THR:OG1	1:A:102:ASP:N	2.50	0.44
1:B:14:LEU:HB2	1:B:249:MET:HE1	2.00	0.44
1:B:405:MET:CE	1:B:408:LEU:HD23	2.48	0.43
1:A:59:ILE:HD11	1:A:246:ILE:CD1	2.47	0.43
1:B:309:THR:HG23	1:B:312:ARG:HH21	1.82	0.43
1:B:400:VAL:HG13	1:B:440:ILE:HD12	2.01	0.42
1:B:157:PRO:HA	1:B:160:ILE:HG23	2.02	0.41
1:B:220:LYS:HG2	1:B:224:MET:HE2	2.02	0.41
1:A:344:LEU:O	1:A:347:LEU:HB2	2.20	0.41
1:B:254:PRO:HD3	1:B:340:THR:HG21	2.03	0.41
1:A:57:CYS:O	1:A:61:LEU:HD22	2.20	0.41
1:B:24:ARG:NH1	1:B:296:ILE:HD11	2.36	0.41
1:B:215:LEU:HA	1:B:215:LEU:HD12	1.94	0.41
1:B:11:LEU:HA	1:B:249:MET:HE1	2.03	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:427:THR:H	1:B:430:HIS:HD2	1.65	0.41
1:B:427:THR:HG1	1:B:429:LEU:HD12	1.83	0.41
1:A:332:ASN:HD22	1:A:333:PRO:HD2	1.86	0.41
1:A:157:PRO:HA	1:A:160:ILE:HG23	2.03	0.41
1:A:405:MET:HE2	1:A:443:LYS:HD2	2.02	0.41
1:B:437:ASN:HB3	1:B:440:ILE:CG1	2.47	0.41
1:B:237:ARG:HA	1:B:241:ASN:HB2	2.03	0.41
1:B:332:ASN:HD22	1:B:333:PRO:HD2	1.85	0.40
1:A:254:PRO:HD2	1:A:280:ILE:HG23	2.02	0.40
1:A:59:ILE:O	1:A:63:VAL:HG13	2.21	0.40
1:B:254:PRO:HD2	1:B:280:ILE:HG23	2.02	0.40
1:B:127:PHE:CE1	1:B:228:LEU:HB2	2.57	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	471/484 (97%)	455 (97%)	16 (3%)	0	100	100
1	B	471/484 (97%)	455 (97%)	15 (3%)	1 (0%)	47	71
All	All	942/968 (97%)	910 (97%)	31 (3%)	1 (0%)	51	75

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	327	ASP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	412/422 (98%)	360 (87%)	52 (13%)	4 8
1	B	413/422 (98%)	375 (91%)	38 (9%)	9 17
All	All	825/844 (98%)	735 (89%)	90 (11%)	6 11

All (90) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	LYS
1	A	25	LEU
1	A	34	LYS
1	A	53	LEU
1	A	59	ILE
1	A	61	LEU
1	A	63	VAL
1	A	69	GLU
1	A	82	LYS
1	A	86	LYS
1	A	89	GLU
1	A	143	PHE
1	A	146	ARG
1	A	149	LYS
1	A	160	ILE
1	A	163	LEU
1	A	166	GLN
1	A	186	VAL
1	A	188	LYS
1	A	203	LYS
1	A	211	LEU
1	A	215	LEU
1	A	223	ARG
1	A	229	HIS
1	A	240	VAL
1	A	288	GLU
1	A	296	ILE

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>
1	A	312	ARG
1	A	315	LEU
1	A	331	LEU
1	A	332	ASN
1	A	334	GLU
1	A	336	LEU
1	A	338	LEU
1	A	340	THR
1	A	341	GLN
1	A	347	LEU
1	A	349	GLU
1	A	360	LEU
1	A	368	LEU
1	A	371	SER
1	A	372	ASN
1	A	374	ASN
1	A	383	GLU
1	A	384	GLN
1	A	415	LYS
1	A	437	ASN
1	A	449	LEU
1	A	451	GLN
1	A	452	GLU
1	A	455	ILE
1	A	467	THR
1	B	53	LEU
1	B	59	ILE
1	B	69	GLU
1	B	82	LYS
1	B	90	GLU
1	B	160	ILE
1	B	163	LEU
1	B	166	GLN
1	B	186	VAL
1	B	211	LEU
1	B	215	LEU
1	B	223	ARG
1	B	229	HIS
1	B	288	GLU
1	B	296	ILE
1	B	297	THR
1	B	327	ASP

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>
1	B	331	LEU
1	B	332	ASN
1	B	336	LEU
1	B	338	LEU
1	B	340	THR
1	B	347	LEU
1	B	351	TYR
1	B	360	LEU
1	B	368	LEU
1	B	371	SER
1	B	374	ASN
1	B	384	GLN
1	B	401	ILE
1	B	415	LYS
1	B	429	LEU
1	B	435	CYS
1	B	437	ASN
1	B	441	MET
1	B	449	LEU
1	B	455	ILE
1	B	467	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (29) such sidechains are listed below:

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>
1	A	92	GLN
1	A	178	ASN
1	A	199	GLN
1	A	221	HIS
1	A	241	ASN
1	A	245	ASN
1	A	251	GLN
1	A	332	ASN
1	A	374	ASN
1	A	377	GLN
1	A	380	GLN
1	A	430	HIS
1	A	437	ASN
1	A	439	GLN
1	A	463	ASN
1	A	464	HIS
1	B	178	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	199	GLN
1	B	221	HIS
1	B	241	ASN
1	B	245	ASN
1	B	251	GLN
1	B	332	ASN
1	B	374	ASN
1	B	377	GLN
1	B	380	GLN
1	B	430	HIS
1	B	437	ASN
1	B	439	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	PO4	B	2000	-	4,4,4	1.24	1 (25%)	6,6,6	0.47	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	PO4	A	2001	-	4,4,4	1.36	1 (25%)	6,6,6	0.42	0
4	GOL	A	3000	-	5,5,5	0.15	0	5,5,5	0.45	0
2	C5P	B	1000	-	19,22,22	0.79	1 (5%)	24,33,33	1.33	2 (8%)
2	C5P	A	1000	-	19,22,22	0.76	1 (5%)	24,33,33	1.23	2 (8%)
3	PO4	A	2000	-	4,4,4	1.29	1 (25%)	6,6,6	0.41	0
3	PO4	B	2001	-	4,4,4	1.19	1 (25%)	6,6,6	0.38	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	3000	-	-	2/4/4/4	-
2	C5P	B	1000	-	-	3/8/26/26	0/2/2/2
2	C5P	A	1000	-	-	3/8/26/26	0/2/2/2

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	2001	PO4	P-O1	2.58	1.56	1.50
2	B	1000	C5P	C6-N1	2.46	1.38	1.35
2	A	1000	C5P	C6-N1	2.43	1.38	1.35
3	A	2000	PO4	P-O1	2.41	1.56	1.50
3	B	2000	PO4	P-O1	2.31	1.56	1.50
3	B	2001	PO4	P-O1	2.18	1.56	1.50

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1000	C5P	C2-N3-C4	4.34	120.74	116.34
2	A	1000	C5P	C2-N3-C4	4.23	120.62	116.34
2	B	1000	C5P	P-O5'-C5'	3.98	129.26	118.30
2	A	1000	C5P	P-O5'-C5'	3.20	127.10	118.30

There are no chirality outliers.

All (8) torsion outliers are listed below:




Mol	Chain	Res	Type	Atoms
4	A	3000	GOL	O1-C1-C2-O2
4	A	3000	GOL	O1-C1-C2-C3
2	B	1000	C5P	C5'-O5'-P-O3P
2	A	1000	C5P	C5'-O5'-P-O3P
2	A	1000	C5P	C5'-O5'-P-O2P
2	B	1000	C5P	C5'-O5'-P-O1P
2	A	1000	C5P	C5'-O5'-P-O1P
2	B	1000	C5P	C5'-O5'-P-O2P

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	3000	GOL	1	0
2	A	1000	C5P	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	475/484 (98%)	-0.04	17 (3%) 42 35	37, 62, 101, 123	1 (0%)
1	B	475/484 (98%)	-0.06	6 (1%) 77 73	39, 62, 103, 136	1 (0%)
All	All	950/968 (98%)	-0.05	23 (2%) 59 53	37, 62, 102, 136	2 (0%)

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	87	LYS	4.3
1	A	88	VAL	3.8
1	A	143	PHE	3.5
1	A	331	LEU	3.4
1	A	89	GLU	3.3
1	A	86	LYS	3.3
1	A	305	LEU	2.9
1	A	378	ILE	2.8
1	A	306	GLU	2.8
1	B	330	ASP	2.8
1	A	1	GLY	2.7
1	B	94	LYS	2.7
1	B	328	PHE	2.6
1	A	4	ILE	2.5
1	A	85	GLY	2.5
1	A	146	ARG	2.5
1	B	292	ARG	2.5
1	B	149	LYS	2.5
1	B	93	GLU	2.3
1	A	449	LEU	2.3
1	A	147	PHE	2.2
1	A	372	ASN	2.1
1	A	158	GLU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	GOL	A	3000	6/6	0.76	0.31	36,48,55,55	0
3	PO4	A	2000	5/5	0.79	0.34	120,120,121,122	0
3	PO4	A	2001	5/5	0.89	0.23	113,113,115,115	0
3	PO4	B	2000	5/5	0.90	0.24	125,125,125,125	0
2	C5P	A	1000	21/21	0.94	0.13	57,59,64,66	0
2	C5P	B	1000	21/21	0.96	0.12	57,62,65,65	0
3	PO4	B	2001	5/5	0.97	0.22	81,81,82,83	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around C5P A 1000:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.