Full wwPDB X-ray Structure Validation Report (i)

Jan 6, 2022 – 03:16 pm GMT

PDB ID	:	7BE8
Title	:	Escherichia coli YtfE (Mn)
Authors	:	Silva, L.S.O.; Matias, P.M.; Romao, C.V.; Saraiva, L.M.
Deposited on	:	2020-12-22
Resolution	:	2.02 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.24
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0267
CCP4	:	7.1.010 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.24

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.02 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	10434 (2.04-2.00)
Clashscore	141614	11643 (2.04-2.00)
Ramachandran outliers	138981	11493 (2.04-2.00)
Sidechain outliers	138945	11492 (2.04-2.00)
RSRZ outliers	127900	10220 (2.04-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	220	6% 97% •
1	B	220	35% 90% 8% •

2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 7001 atoms, of which 3415 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Iron-sulfur cluster repair protein YtfE.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	219	3436	1093	1699	300	332	12	0	2	0
1	B	216	3401	1075	1692	294	328	12	0	2	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	30	ALA	CYS	engineered mutation	UNP P69506
A	31	ALA	CYS	engineered mutation	UNP P69506
B	30	ALA	CYS	engineered mutation	UNP P69506
B	31	ALA	CYS	engineered mutation	UNP P69506

- Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mn		
2	A	2	2	2	0	0
2	B	2	2	2	0	0

- Molecule 3 is OXYGEN ATOM (three-letter code: O) (formula: O).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	1	1	1	0	0
3	B	1	1	1	0	0

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
4	A	1	5	4	1	0	0
			Total	O	S		
4	A	1	5	4	1	0	0

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
5	A	1	12	3	6	3	0	0
			Total	C	H	O		
5	A	1	11	3	5	3	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
5	B	1	14	3	8	3	0	0
			Total	C	H	O		
5	B	1	11	3	5	3	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
6	A	77	77	77	0	0
			Total	O		
6	B	23	23	23	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Iron-sulfur cluster repair protein YtfE

Chain A:





• Molecule 1: Iron-sulfur cluster repair protein YtfE

Chain B:







4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α, β, γ	60.15Å 50.12Å 88.24Å 90.00° 100.30° 90.00°	Depositor
Resolution (Å)	59.18 – 2.02 59.18 – 2.02	Depositor EDS
% Data completeness (in resolution range)	77.9 (59.18-2.02) 78.0 (59.18-2.02)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.55 (at 2.02Å)	Xtriage
Refinement program	REFMAC 1.14_3260, PHENIX 1.14_3260	Depositor
R, R_{free}	0.239 , 0.287 0.252 , 0.288	Depositor DCC
R_{free} test set	1268 reflections (4.74%)	wwPDB-VP
Wilson B-factor (Å^2)	36.8	Xtriage
Anisotropy	0.156	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	(Not available) , (Not available)	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.92	EDS
Total number of atoms	7001	wwPDB-VP
Average B, all atoms (Å^2)	61.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 16.16% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: MN, O, GOL, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.24	0/1778	0.38	0/2405
1	B	0.24	0/1749	0.38	0/2366
All	All	0.24	0/3527	0.38	0/4771

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1737	1699	1728	5	0
1	B	1709	1692	1701	11	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	10	0	0	0	0
5	A	12	11	16	1	0
5	B	12	13	16	0	0
6	A	77	0	0	2	0
6	B	23	0	0	1	0
All	All	3586	3415	3461	16	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (16) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:143:GLN:NE2	6:B:401:HOH:O	2.21	0.72
1:B:41:ALA:O	1:B:45:GLU:N	2.22	0.71
1:A:162:GLU:OE1	6:A:402:HOH:O	2.15	0.65
1:A:90:GLN:OE1	6:A:401:HOH:O	2.13	0.65
1:B:47:ASP:OD1	1:B:49:GLU:N	2.35	0.59
1:B:63:ILE:HD12	1:B:63:ILE:H	1.70	0.56
1:B:44:LYS:HB3	1:B:46:LEU:HD12	1.88	0.54
1:B:14:SER:O	1:B:63:ILE:HD11	2.08	0.54
1:A:91:LEU:HD13	1:A:123:HIS:HA	1.90	0.53
1:B:54:GLU:O	1:B:58:LEU:HD13	2.12	0.50
1:B:77:ASP:O	1:B:81:VAL:HG22	2.13	0.48
1:A:155:VAL:HG22	5:A:307:GOL:H12	1.98	0.45
1:B:68:ARG:O	1:B:215:ARG:NH1	2.51	0.43
1:B:177:VAL:O	1:B:177:VAL:HG12	2.19	0.43
1:A:177:VAL:O	1:A:177:VAL:HG12	2.19	0.42

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	219/220 (100%)	211 (96%)	8 (4%)	0	100	100
1	B	216/220 (98%)	207 (96%)	9 (4%)	0	100	100
All	All	435/440 (99%)	418 (96%)	17 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	188/187 (100%)	188 (100%)	0	100	100
1	B	186/187 (100%)	185 (100%)	1 (0%)	88	91
All	All	374/374 (100%)	373 (100%)	1 (0%)	92	94

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	140	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 12 ligands modelled in this entry, 6 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
4	SO4	A	305	-	4,4,4	0.14	0	6,6,6	0.08	0
5	GOL	A	307	-	5,5,5	0.90	0	5,5,5	0.94	0
5	GOL	B	305	-	5,5,5	0.90	0	5,5,5	1.01	0
5	GOL	A	306	-	5,5,5	0.87	0	5,5,5	0.99	0
4	SO4	A	304	-	4,4,4	0.14	0	6,6,6	0.06	0
5	GOL	B	304	-	5,5,5	0.86	0	5,5,5	1.00	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	B	304	-	-	2/4/4/4	-
5	GOL	A	306	-	-	2/4/4/4	-
5	GOL	B	305	-	-	0/4/4/4	-
5	GOL	A	307	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	306	GOL	O1-C1-C2-C3
5	B	304	GOL	O1-C1-C2-C3
5	B	304	GOL	O1-C1-C2-O2
5	A	306	GOL	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	307	GOL	1	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	219/220 (99%)	1.00	14 (6%)	19	18	25, 38, 59, 68	0
1	B	216/220 (98%)	2.34	77 (35%)	0	0	36, 55, 152, 164	0
All	All	435/440 (98%)	1.66	91 (20%)	1	0	25, 46, 143, 164	0

All (91) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	43	ARG	20.8
1	B	46	LEU	15.2
1	B	50	VAL	14.5
1	B	62	PRO	12.8
1	B	61	GLN	12.4
1	B	55	LEU	11.7
1	B	40	ALA	11.0
1	B	36	THR	9.6
1	B	45	GLU	9.5
1	B	41	ALA	9.5
1	B	51	ILE	8.7
1	B	44	LYS	8.6
1	B	47	ASP	8.6
1	B	38	ALA	7.5
1	B	59	ALA	7.5
1	B	7	PRO	7.5
1	B	42	ALA	7.4
1	B	35	GLN	6.7
1	B	56	ALA	6.6
1	B	11	LEU	6.4
1	B	58	LEU	6.3
1	B	12	ALA	6.2
1	B	60	GLU	5.7
1	B	25	TYR	5.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	17	ARG	5.3
1	B	54	GLU	5.3
1	B	66	ASP	5.2
1	B	63	ILE	5.2
1	B	189	ALA	5.2
1	B	49	GLU	5.2
1	B	24	LYS	4.9
1	B	26	ASP	4.9
1	B	109	PRO	4.8
1	A	26	ASP	4.8
1	A	63	ILE	4.4
1	B	21	LEU	4.4
1	B	57	LYS	4.3
1	B	39	ARG	4.2
1	B	8	LEU	4.0
1	B	31	ALA	4.0
1	B	34	LYS	3.8
1	B	14	SER	3.6
1	B	33	GLY	3.6
1	B	64	GLU	3.5
1	B	48	VAL	3.5
1	B	52	GLU	3.5
1	A	43	ARG	3.4
1	B	148	GLN	3.4
1	B	70	ALA	3.3
1	B	187	TRP	3.3
1	B	192	ASN	3.2
1	B	181	PRO	3.2
1	B	9	GLY	3.1
1	B	5	ASP	3.1
1	A	64	GLU	3.1
1	B	28	ASP	3.1
1	A	65	LYS	3.1
1	B	27	MET	3.1
1	B	110	SER	3.0
1	B	65	LYS	3.0
1	B	20	ALA	3.0
1	A	181	PRO	2.9
1	B	23	ARG	2.9
1	B	175	ASN	2.9
1	B	107	ASP	2.8
1	B	37	LEU	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	177	VAL	2.8
1	B	13	LEU	2.8
1	A	66	ASP	2.7
1	B	184[A]	CYS	2.7
1	B	53	ALA	2.7
1	A	46	LEU	2.6
1	A	57	LYS	2.6
1	B	183	ALA	2.6
1	B	22	PHE	2.6
1	B	101	VAL	2.5
1	B	182	GLU	2.5
1	A	183	ALA	2.4
1	A	36	THR	2.3
1	B	193	GLY	2.3
1	A	131	MET	2.3
1	B	30	ALA	2.3
1	A	51	ILE	2.3
1	B	219	GLY	2.3
1	A	138	PHE	2.3
1	B	15	ILE	2.2
1	B	138	PHE	2.2
1	B	68	ARG	2.2
1	B	136	ILE	2.1
1	B	165	GLU	2.1
1	B	29	TYR	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
5	GOL	A	306	6/6	0.62	0.17	63,75,87,87	0
5	GOL	B	305	6/6	0.62	0.29	131,139,168,168	0
4	SO4	A	304	5/5	0.68	0.29	58,64,79,90	0
5	GOL	B	304	6/6	0.71	0.19	61,75,91,93	0
5	GOL	A	307	6/6	0.78	0.34	27,49,60,60	0
4	SO4	A	305	5/5	0.83	0.21	68,70,80,87	0
2	MN	B	302	1/1	0.92	0.07	51,51,51,51	0
3	O	B	303	1/1	0.95	0.34	52,52,52,52	0
2	MN	A	302	1/1	0.96	0.10	38,38,38,38	0
3	O	A	303	1/1	0.98	0.26	35,35,35,35	0
2	MN	B	301	1/1	0.99	0.11	38,38,38,38	0
2	MN	A	301	1/1	0.99	0.17	26,26,26,26	0

6.5 Other polymers (i)

There are no such residues in this entry.