



wwPDB EM Validation Summary Report ⓘ

Nov 20, 2022 – 05:48 pm GMT

PDB ID : 4BIJ
EMDB ID : EMD-2355
Title : Threading model of T7 large terminase
Authors : Dauden, M.I.; Martin-Benito, J.; Sanchez-Ferrero, J.C.; Pulido-Cid, M.; Valpuesta, J.M.; Carrascosa, J.L.
Deposited on : 2013-04-10
Resolution : 16.00 Å(reported)
Based on initial model : 3CPE

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev43
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.9
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.31.2


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 16.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	476	22% (Poor fit) 35% (0 outliers) 46% (1 outlier) 14% (2 outliers) 5% (3+ outliers)
1	B	476	22% (Poor fit) 33% (0 outliers) 45% (1 outlier) 16% (2 outliers) 6% (3+ outliers)
1	C	476	22% (Poor fit) 36% (0 outliers) 45% (1 outlier) 14% (2 outliers) 5% (3+ outliers)
1	D	476	22% (Poor fit) 35% (0 outliers) 45% (1 outlier) 14% (2 outliers) 6% (3+ outliers)
1	E	476	22% (Poor fit) 35% (0 outliers) 43% (1 outlier) 16% (2 outliers) 5% (3+ outliers)

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 18855 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called DNA MATURASE B.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	474	Total 3771	2402	648	703	18	0	0
1	B	474	Total 3771	2402	648	703	18	0	0
1	C	474	Total 3771	2402	648	703	18	0	0
1	D	474	Total 3771	2402	648	703	18	0	0
1	E	474	Total 3771	2402	648	703	18	0	0

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: DNA MATURASE B





• Molecule 1: DNA MATURASE B






• Molecule 1: DNA MATURASE B







• Molecule 1: DNA MATURASE B



• Molecule 1: DNA MATURASE B





4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C5	Depositor
Number of particles used	3650	Depositor
Resolution determination method	Not provided	
CTF correction method	EACH PLATE	Depositor
Microscope	FEI TECNAI F20	Depositor
Voltage (kV)	100	Depositor
Electron dose ($e^-/\text{\AA}^2$)	Not provided	
Minimum defocus (nm)	1500	Depositor
Maximum defocus (nm)	3200	Depositor
Magnification	67000	Depositor
Image detector	FEI EAGLE (4k x 4k)	Depositor
Maximum map value	0.094	Depositor
Minimum map value	-0.051	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.009	Depositor
Recommended contour level	0.0383	Depositor
Map size (\AA)	352.0, 352.0, 352.0	wwPDB
Map dimensions	80, 80, 80	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (\AA)	4.4, 4.4, 4.4	Depositor

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.07	22/3850 (0.6%)	1.57	92/5212 (1.8%)
1	B	1.08	22/3850 (0.6%)	1.57	96/5212 (1.8%)
1	C	1.06	23/3850 (0.6%)	1.56	92/5212 (1.8%)
1	D	1.03	22/3850 (0.6%)	1.56	96/5212 (1.8%)
1	E	1.11	22/3850 (0.6%)	1.57	95/5212 (1.8%)
All	All	1.07	111/19250 (0.6%)	1.57	471/26060 (1.8%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	13
1	B	0	14
1	C	0	14
1	D	0	14
1	E	0	13
All	All	0	68

The worst 5 of 111 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	291	PRO	N-CD	-23.77	1.14	1.47
1	B	134	PRO	N-CD	20.15	1.76	1.47
1	E	300	PRO	N-CD	19.45	1.75	1.47
1	C	291	PRO	N-CD	-19.44	1.20	1.47
1	B	291	PRO	N-CD	-19.19	1.21	1.47

The worst 5 of 471 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	291	PRO	N-CD-CG	15.74	126.81	103.20

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	291	PRO	N-CD-CG	15.70	126.75	103.20
1	E	77	TRP	CH2-CZ2-CE2	14.73	132.13	117.40
1	A	77	TRP	CH2-CZ2-CE2	14.71	132.12	117.40
1	B	77	TRP	CH2-CZ2-CE2	14.71	132.10	117.40

There are no chirality outliers.

5 of 68 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	144	GLY	Peptide
1	A	194	SER	Mainchain
1	A	21	ALA	Mainchain
1	A	82	LEU	Peptide
1	A	91	LYS	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3771	0	3778	650	0
1	B	3771	0	3776	649	0
1	C	3771	0	3778	670	0
1	D	3771	0	3778	685	0
1	E	3771	0	3778	661	0
All	All	18855	0	18888	2904	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 77.

The worst 5 of 2904 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:346:TYR:N	1:E:173:ALA:CB	1.69	1.56
1:D:173:ALA:CB	1:E:346:TYR:N	1.69	1.54
1:D:346:TYR:CE1	1:D:349:CYS:HB3	1.41	1.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:173:ALA:CB	1:C:346:TYR:N	1.69	1.52
1:C:173:ALA:CB	1:D:346:TYR:N	1.69	1.51

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	468/476 (98%)	394 (84%)	46 (10%)	28 (6%)	1	17
1	B	468/476 (98%)	394 (84%)	46 (10%)	28 (6%)	1	17
1	C	468/476 (98%)	394 (84%)	46 (10%)	28 (6%)	1	17
1	D	468/476 (98%)	394 (84%)	46 (10%)	28 (6%)	1	17
1	E	468/476 (98%)	394 (84%)	46 (10%)	28 (6%)	1	17
All	All	2340/2380 (98%)	1970 (84%)	230 (10%)	140 (6%)	3	17

5 of 140 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	83	LYS
1	A	92	GLU
1	A	125	ILE
1	A	132	ALA
1	A	182	GLN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	409/411 (100%)	372 (91%)	37 (9%)	9	30
1	B	409/411 (100%)	347 (85%)	62 (15%)	3	14
1	C	409/411 (100%)	370 (90%)	39 (10%)	8	27
1	D	409/411 (100%)	363 (89%)	46 (11%)	6	22
1	E	409/411 (100%)	365 (89%)	44 (11%)	6	23
All	All	2045/2055 (100%)	1817 (89%)	228 (11%)	9	22

5 of 228 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	204	THR
1	E	395	ASP
1	D	42	MET
1	E	348	ASP
1	E	162	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 33 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	182	GLN
1	E	325	ASN
1	E	358	GLN
1	C	335	ASN
1	C	325	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-2355. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map




The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 45



Y Index: 37



Z Index: 37

The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal surface views [i](#)

6.4.1 Primary map



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.0383. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.5 Mask visualisation

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [\(i\)](#)



The volume at the recommended contour level is 435 nm³; this corresponds to an approximate mass of 393 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [i](#)



*Reported resolution corresponds to spatial frequency of 0.062\AA^{-1}


8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-2355 and PDB model 4BIJ. Per-residue inclusion information can be found in section 3 on page 4.


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.0383 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [\(i\)](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [\(i\)](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.0383).













9.4 Atom inclusion [i](#)



At the recommended contour level, 76% of all backbone atoms, 74% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary [i](#)

The table lists the average atom inclusion at the recommended contour level (0.0383) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.7426	 0.0450
A	 0.7436	 0.0410
B	 0.7417	 0.0440
C	 0.7420	 0.0480
D	 0.7425	 0.0460
E	 0.7431	 0.0440

