



Full wwPDB X-ray Structure Validation Report ⓘ

May 14, 2020 – 01:46 pm BST

PDB ID : 2BJN
Title : X-ray Structure of human TPC6
Authors : Kummel, D.; Mueller, J.J.; Roske, Y.; Misselwitz, R.; Bussow, K.; Heinemann, U.
Deposited on : 2005-02-04
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4298 (1.70-1.70)
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	160	 78% 11% 11%
1	B	160	 79% 11% 7%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SO4	A	1161	-	-	X	-
2	SO4	B	1160	-	-	X	-

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 2504 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TRAFFICKING PROTEIN PARTICLE COMPLEX SUB-UNIT 6B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	143	1142	727	199	205	11	0	3	1
1	B	149	1187	757	204	216	10	0	5	1

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	A	1	5	4	1	0	0
2	A	1	5	4	1	0	0
2	B	1	5	4	1	0	0
2	B	1	5	4	1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total O S 5 4 1	0	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0


- Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	58	Total O 58 58	0	0
4	B	68	Total O 68 68	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: TRAFFICKING PROTEIN PARTICLE COMPLEX SUBUNIT 6B

Chain A:  78% 11% • 11%



- Molecule 1: TRAFFICKING PROTEIN PARTICLE COMPLEX SUBUNIT 6B

Chain B:  79% 11% • 7%



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	99.42Å 57.01Å 60.41Å 90.00° 117.16° 90.00°	Depositor
Resolution (Å)	22.98 – 1.70 22.98 – 1.70	Depositor EDS
% Data completeness (in resolution range)	98.6 (22.98-1.70) 98.6 (22.98-1.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.42 (at 1.70Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.175 , 0.209 0.188 , 0.223	Depositor DCC
R_{free} test set	1697 reflections (5.19%)	wwPDB-VP
Wilson B-factor (Å ²)	26.5	Xtrriage
Anisotropy	0.042	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 57.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2504	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.76% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.76	1/1177 (0.1%)	0.82	1/1572 (0.1%)
1	B	0.85	0/1227	0.92	5/1642 (0.3%)
All	All	0.81	1/2404 (0.0%)	0.87	6/3214 (0.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	64	LEU	C-O	5.91	1.34	1.23

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	41	MET	CG-SD-CE	-10.74	83.02	100.20
1	A	41	MET	CG-SD-CE	-10.24	83.81	100.20
1	B	16	MET	CG-SD-CE	-8.93	85.92	100.20
1	B	10[A]	LEU	CA-CB-CG	-5.35	103.00	115.30
1	B	10[B]	LEU	CA-CB-CG	-5.35	103.00	115.30
1	B	156	MET	CG-SD-CE	5.10	108.36	100.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1142	0	1163	23	0
1	B	1187	0	1217	23	1
2	A	10	0	0	12	0
2	B	15	0	0	4	1
3	A	12	0	16	1	0
3	B	12	0	16	0	0
4	A	58	0	0	0	0
4	B	68	0	0	5	0
All	All	2504	0	2412	45	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:32:GLY:HA2	2:A:1161:SO4:O2	1.67	0.93
1:B:156:MET:HB3	2:B:1160:SO4:O3	1.71	0.89
1:B:156:MET:SD	4:B:2061:HOH:O	2.32	0.87
1:B:156:MET:HG3	4:B:2061:HOH:O	1.79	0.83
1:A:35:ILE:HG12	2:A:1161:SO4:O1	1.85	0.76
1:A:34:CYS:HB2	2:A:1161:SO4:O4	1.89	0.73
1:A:32:GLY:HA2	2:A:1161:SO4:S	2.31	0.70
1:B:154:GLN:NE2	1:B:156:MET:SD	2.65	0.69
1:B:156:MET:HB3	2:B:1160:SO4:S	2.34	0.67
1:B:31:ASN:C	1:B:31:ASN:HD22	2.01	0.64
1:B:61:LYS:HG2	1:B:62:ASP:N	2.13	0.63
1:A:29:VAL:O	2:A:1161:SO4:O2	2.18	0.62
1:A:35:ILE:CG1	2:A:1161:SO4:O1	2.48	0.62
1:B:85:ASN:OD1	1:B:87:ARG:HG3	2.01	0.60
1:A:92:GLY:HA2	1:A:157:ILE:HG23	1.86	0.58
1:A:34:CYS:CB	2:A:1161:SO4:O4	2.51	0.58
1:A:34:CYS:CA	2:A:1161:SO4:O4	2.53	0.57
1:A:29:VAL:O	2:A:1161:SO4:S	2.63	0.57
1:A:32:GLY:CA	2:A:1161:SO4:S	2.89	0.57
1:A:35:ILE:HD11	3:A:1163:GOL:H2	1.87	0.57
1:A:29:VAL:O	2:A:1161:SO4:O3	2.22	0.57
1:A:52:ARG:HG3	1:B:3:MET:HG2	1.88	0.55
1:B:31:ASN:ND2	1:B:33:ARG:H	2.05	0.54
1:A:32:GLY:HA2	2:A:1161:SO4:O1	2.08	0.54
1:B:31:ASN:HD22	1:B:32:GLY:N	2.06	0.53
1:B:31:ASN:ND2	1:B:31:ASN:C	2.63	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:156:MET:CB	2:B:1160:SO4:O3	2.53	0.51
1:A:92:GLY:CA	1:A:157:ILE:HG23	2.40	0.51
1:B:156:MET:CE	4:B:2061:HOH:O	2.58	0.49
1:B:116:HIS:C	1:B:116:HIS:CD2	2.87	0.47
1:A:92:GLY:HA2	1:A:157:ILE:CG2	2.46	0.46
1:A:87:ARG:NE	1:A:89:ASN:HD21	2.13	0.45
1:A:87:ARG:NE	1:A:89:ASN:ND2	2.65	0.45
1:B:156:MET:CG	4:B:2061:HOH:O	2.38	0.45
1:B:13:HIS:CE1	1:B:103:LEU:HD13	2.52	0.45
1:A:50:ILE:HD13	1:A:50:ILE:HA	1.82	0.44
1:B:140:ILE:HD13	1:B:140:ILE:N	2.34	0.43
1:A:87:ARG:HE	1:A:89:ASN:HD21	1.66	0.42
1:B:54:THR:N	2:B:1161:SO4:O4	2.36	0.41
1:B:55:LYS:HD3	1:B:60:PHE:CE2	2.55	0.41
1:B:10[A]:LEU:HD22	1:B:103:LEU:HD23	2.02	0.41
1:A:81:LYS:NZ	1:A:98:ASP:OD1	2.48	0.41
1:A:50:ILE:CG1	1:A:131:GLY:HA2	2.51	0.40
1:B:156:MET:HE2	4:B:2061:HOH:O	2.20	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:89:ASN:ND2	2:B:1160:SO4:O4[2_657]	2.16	0.04

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	141/160 (88%)	135 (96%)	6 (4%)	0	100 100
1	B	150/160 (94%)	144 (96%)	5 (3%)	1 (1%)	22 8

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	291/320 (91%)	279 (96%)	11 (4%)	1 (0%)	34 24

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	61	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	127/138 (92%)	123 (97%)	4 (3%)	40 21
1	B	133/138 (96%)	129 (97%)	4 (3%)	41 22
All	All	260/276 (94%)	252 (97%)	8 (3%)	39 21

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	85	ASN
1	A	106	GLN
1	A	146	SER
1	A	157	ILE
1	B	31	ASN
1	B	81	LYS
1	B	104	LEU
1	B	140	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	85	ASN
1	A	89	ASN
1	B	31	ASN
1	B	40	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	116	HIS
1	B	158	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GOL	A	1163	2	5,5,5	0.48	0	5,5,5	0.39	0
2	SO4	B	1160	-	4,4,4	0.19	0	6,6,6	0.62	0
2	SO4	A	1160	-	4,4,4	0.55	0	6,6,6	0.36	0
3	GOL	A	1162	-	5,5,5	0.28	0	5,5,5	0.83	0
3	GOL	B	1163	-	5,5,5	0.49	0	5,5,5	0.22	0
2	SO4	B	1162	-	4,4,4	0.44	0	6,6,6	0.59	0
3	GOL	B	1164	-	5,5,5	0.46	0	5,5,5	0.75	0
2	SO4	B	1161	-	4,4,4	0.24	0	6,6,6	0.36	0
2	SO4	A	1161	1,3	4,4,4	0.26	0	6,6,6	0.27	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	B	1164	-	-	2/4/4/4	-
3	GOL	A	1163	2	-	2/4/4/4	-
3	GOL	B	1163	-	-	0/4/4/4	-
3	GOL	A	1162	-	-	4/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1163	GOL	O1-C1-C2-C3
3	A	1162	GOL	C1-C2-C3-O3
3	B	1164	GOL	O1-C1-C2-C3
3	A	1163	GOL	O1-C1-C2-O2
3	A	1162	GOL	O2-C2-C3-O3
3	A	1162	GOL	O1-C1-C2-O2
3	B	1164	GOL	O1-C1-C2-O2
3	A	1162	GOL	O1-C1-C2-C3

There are no ring outliers.

4 monomers are involved in 18 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1163	GOL	1	0
2	B	1160	SO4	3	1
2	B	1161	SO4	1	0
2	A	1161	SO4	12	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers

Unable to reproduce the depositors R factor - this section is therefore empty.