



# Full wwPDB X-ray Structure Validation Report ⓘ

Dec 3, 2023 – 10:40 am GMT

PDB ID : 2BMZ  
Title : Banana Lectin bound to Xyl-b1,3 Man-a-O-Methyl (XM)  
Authors : Meagher, J.L.; Winter, H.C.; Ezell, P.; Goldstein, I.J.; Stuckey, J.A.  
Deposited on : 2005-03-17  
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtrriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36



## 2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 2373 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RIPENING-ASSOCIATED PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	141	1031	664	171	193	3	1	0	0
1	B	138	1011	653	167	189	2	1	0	0

- Molecule 2 is an oligosaccharide called beta-D-xylopyranose-(1-3)-methyl alpha-D-mannopyranoside.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
			Total	C	O			
2	C	2	22	12	10	0	0	0
2	D	2	22	12	10	0	0	0
2	E	2	22	12	10	0	0	0
2	F	2	22	12	10	0	0	0

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0

- Molecule 4 is CADMIUM ION (three-letter code: CD) (formula: Cd).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	2	Total Cd 2 2	0	0

- Molecule 5 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	100	Total O 100 100	0	0
5	B	121	Total O 121 121	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


Note EDS was not executed.

- Molecule 1: RIPENING-ASSOCIATED PROTEIN

Chain A:  84% 16%



- Molecule 1: RIPENING-ASSOCIATED PROTEIN

Chain B:  84% 13%



- Molecule 2: beta-D-xylopyranose-(1-3)-methyl alpha-D-mannopyranoside

Chain C:  100%



- Molecule 2: beta-D-xylopyranose-(1-3)-methyl alpha-D-mannopyranoside

Chain D:  50% 50%



- Molecule 2: beta-D-xylopyranose-(1-3)-methyl alpha-D-mannopyranoside

Chain E:  50% 50%



- Molecule 2: beta-D-xylopyranose-(1-3)-methyl alpha-D-mannopyranoside

Chain F:  50% 50%



## 4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	81.62Å 81.62Å 146.81Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	9.99 – 2.40	Depositor
% Data completeness (in resolution range)	97.3 (9.99-2.40)	Depositor
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.228 , 0.256	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	2373	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	36.0	wwPDB-VP

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: CD, MMA, SO4, XYP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.40	0/1058	0.67	1/1427 (0.1%)
1	B	0.39	0/1038	0.68	1/1401 (0.1%)
All	All	0.40	0/2096	0.68	2/2828 (0.1%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	9	ALA	N-CA-C	-7.67	90.28	111.00
1	B	9	ALA	N-CA-C	-6.30	93.98	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1031	0	1001	14	0
1	B	1011	0	980	12	0
2	C	22	0	13	0	0
2	D	22	0	13	0	0
2	E	22	0	13	0	0
2	F	22	0	13	0	0
3	A	10	0	0	0	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	10	0	0	0	0
4	A	2	0	0	0	0
5	A	100	0	0	0	0
5	B	121	0	0	0	0
All	All	2373	0	2033	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:89:LEU:HD11	1:B:135:ILE:HG22	1.78	0.64
1:B:6:LYS:HD3	1:B:138:TYR:CZ	2.34	0.63
1:A:32:PHE:HA	1:A:61:THR:O	2.03	0.57
1:B:20:MET:HB2	1:B:55:TYR:OH	2.06	0.55
1:A:54:HIS:HB3	1:B:84:HIS:CD2	2.44	0.53
1:B:8:GLY:O	1:B:9:ALA:CB	2.58	0.52
1:A:8:GLY:O	1:A:9:ALA:CB	2.57	0.51
1:A:72:TYR:CD2	1:A:120:LYS:HE2	2.45	0.51
1:B:96:THR:C	1:B:98:LYS:H	2.14	0.51
1:A:57:GLY:HA3	1:A:130:LYS:O	2.13	0.48
1:A:42:VAL:O	1:A:52:THR:HA	2.14	0.48
1:A:96:THR:C	1:A:98:LYS:H	2.15	0.48
1:B:79:GLU:HG2	1:B:111:PRO:HA	1.96	0.48
1:B:6:LYS:HD3	1:B:138:TYR:CE2	2.49	0.47
1:A:52:THR:O	1:A:53:ARG:HD2	2.16	0.45
1:A:54:HIS:HB3	1:B:84:HIS:NE2	2.31	0.45
1:B:30:LYS:HG2	1:B:64:GLU:HB2	1.99	0.45
1:A:31:ILE:HD12	1:A:31:ILE:N	2.33	0.44
1:A:89:LEU:HD21	1:A:134:ALA:HA	2.00	0.44
1:B:57:GLY:HA3	1:B:130:LYS:O	2.17	0.43
1:A:20:MET:CE	1:A:23:ALA:HB2	2.50	0.42
1:A:53:ARG:HB3	1:A:55:TYR:CZ	2.56	0.41
1:A:32:PHE:CZ	1:A:62:PRO:HB3	2.56	0.40
1:B:121:ILE:HG23	1:B:137:VAL:HB	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

There are no protein backbone outliers to report in this entry.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	100/100 (100%)	98 (98%)	2 (2%)	55	74
1	B	98/100 (98%)	97 (99%)	1 (1%)	76	88
All	All	198/200 (99%)	195 (98%)	3 (2%)	65	80

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	80	VAL
1	A	133	ASP
1	B	110	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	2	ASN
1	A	68	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates i

8 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	MMA	C	1	2	13,13,13	1.15	1 (7%)	18,18,18	0.65	0
2	XYP	C	2	2	9,9,10	1.50	2 (22%)	10,12,14	0.63	0
2	MMA	D	1	2	13,13,13	1.05	0	18,18,18	0.71	0
2	XYP	D	2	2	9,9,10	1.35	1 (11%)	10,12,14	0.66	0
2	MMA	E	1	2	13,13,13	1.05	0	18,18,18	0.72	0
2	XYP	E	2	2	9,9,10	1.46	2 (22%)	10,12,14	0.65	0
2	MMA	F	1	2	13,13,13	1.08	0	18,18,18	0.72	0
2	XYP	F	2	2	9,9,10	1.50	1 (11%)	10,12,14	0.65	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MMA	C	1	2	-	0/4/24/24	0/1/1/1
2	XYP	C	2	2	-	-	0/1/1/1
2	MMA	D	1	2	-	0/4/24/24	0/1/1/1
2	XYP	D	2	2	-	-	0/1/1/1
2	MMA	E	1	2	-	0/4/24/24	0/1/1/1
2	XYP	E	2	2	-	-	0/1/1/1
2	MMA	F	1	2	-	0/4/24/24	0/1/1/1
2	XYP	F	2	2	-	-	0/1/1/1

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	2	XYP	O5-C5	2.99	1.48	1.42
2	F	2	XYP	O5-C5	2.98	1.48	1.42
2	D	2	XYP	O5-C5	2.78	1.48	1.42

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	2	XYP	O5-C5	2.76	1.48	1.42
2	E	2	XYP	O5-C1	2.20	1.47	1.42
2	C	2	XYP	O5-C1	2.09	1.46	1.42
2	C	1	MMA	O5-C1	2.01	1.47	1.41

There are no bond angle outliers.


There are no chirality outliers.



There are no torsion outliers.


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.







## 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	SO4	B	1145	-	4,4,4	0.31	0	6,6,6	0.07	0
3	SO4	B	1143	-	4,4,4	0.29	0	6,6,6	0.08	0
3	SO4	A	1144	-	4,4,4	0.27	0	6,6,6	0.07	0
3	SO4	A	1141	-	4,4,4	0.29	0	6,6,6	0.05	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

### 6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.