



Full wwPDB X-ray Structure Validation Report i

May 28, 2020 – 09:29 pm BST

PDB ID : 2BNL
Title : The structure of the N-terminal domain of RsbR
Authors : Murray, J.W.; Delumeau, O.; Lewis, R.J.
Deposited on : 2005-03-28
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 7278 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MODULATOR PROTEIN RSBR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	129	Total	C 1075	N 689	O 168	Se 216	2	0	0
1	B	128	Total	C 1066	N 683	O 166	Se 215	2	0	0
1	C	134	Total	C 1118	N 715	O 177	Se 224	2	0	0
1	D	129	Total	C 1075	N 689	O 168	Se 216	2	0	0
1	E	129	Total	C 1075	N 689	O 168	Se 216	2	0	0
1	F	129	Total	C 1075	N 689	O 168	Se 216	2	0	0

- Molecule 2 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Na 1	0	0
2	C	1	Total	Na 1	0	0
2	F	1	Total	Na 1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	127	Total	O 127	0	0
3	B	133	Total	O 133	0	0
3	C	134	Total	O 134	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	D	145	Total O 145 145	0	0
3	E	113	Total O 113 113	0	0
3	F	139	Total O 139 139	0	0


3 Residue-property plots [\(i\)](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: MODULATOR PROTEIN RSBR




- Molecule 1: MODULATOR PROTEIN RSBR



- Molecule 1: MODULATOR PROTEIN RSBR




- Molecule 1: MODULATOR PROTEIN RSBR



- Molecule 1: MODULATOR PROTEIN RSBR





4 Data and refinement statistics i

Property	Value	Source
Space group	P 32 1 2	Depositor
Cell constants a, b, c, α , β , γ	136.06 Å 136.06 Å 113.30 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	22.24 – 2.00 22.24 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.9 (22.24-2.00) 99.9 (22.24-2.00)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^{\text{1}}$	3.79 (at 2.01 Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R , R_{free}	0.154 , 0.198 0.168 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	30.6	Xtriage
Anisotropy	0.455	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 53.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.029 for -h,-k,l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7278	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 30.10 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.3838e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section:
NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.57	16/1092 (1.5%)	1.07	5/1475 (0.3%)
1	B	1.55	15/1083 (1.4%)	1.02	1/1464 (0.1%)
1	C	1.54	10/1137 (0.9%)	1.08	3/1538 (0.2%)
1	D	1.67	18/1092 (1.6%)	1.21	8/1475 (0.5%)
1	E	1.45	13/1092 (1.2%)	1.11	8/1475 (0.5%)
1	F	1.59	13/1092 (1.2%)	1.11	3/1475 (0.2%)
All	All	1.56	85/6588 (1.3%)	1.10	28/8902 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	1

All (85) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	81	MSE	SE-CE	-11.69	1.26	1.95
1	F	81	MSE	SE-CE	-11.51	1.27	1.95
1	C	48	GLU	CD-OE1	11.33	1.38	1.25
1	A	81	MSE	SE-CE	-10.66	1.32	1.95
1	B	81	MSE	SE-CE	-9.85	1.37	1.95
1	D	93	LYS	CE-NZ	9.66	1.73	1.49
1	C	48	GLU	CG-CD	9.26	1.65	1.51
1	C	49	TYR	CE2-CZ	8.69	1.49	1.38
1	D	93	LYS	CD-CE	8.66	1.73	1.51
1	A	17	GLU	CG-CD	8.59	1.64	1.51
1	D	17	GLU	CG-CD	8.53	1.64	1.51
1	B	48	GLU	CD-OE1	8.48	1.34	1.25

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	F	134	SER	CB-OG	8.22	1.52	1.42
1	D	17	GLU	CD-OE1	8.09	1.34	1.25
1	E	73	ARG	CZ-NH2	8.03	1.43	1.33
1	F	93	LYS	CE-NZ	7.82	1.68	1.49
1	E	93	LYS	CE-NZ	7.79	1.68	1.49
1	D	108	GLU	CG-CD	7.43	1.63	1.51
1	F	48	GLU	CD-OE1	7.38	1.33	1.25
1	E	90	GLU	CB-CG	-7.26	1.38	1.52
1	B	48	GLU	CG-CD	7.14	1.62	1.51
1	A	111	GLU	CG-CD	7.02	1.62	1.51
1	C	90	GLU	CB-CG	-7.02	1.38	1.52
1	A	111	GLU	CB-CG	6.90	1.65	1.52
1	B	111	GLU	CD-OE2	6.89	1.33	1.25
1	C	64	GLU	CG-CD	6.80	1.62	1.51
1	A	136	GLU	CD-OE2	6.73	1.33	1.25
1	A	90	GLU	CB-CG	-6.72	1.39	1.52
1	B	8	TYR	CZ-OH	6.60	1.49	1.37
1	E	48	GLU	CG-CD	6.59	1.61	1.51
1	D	28	GLU	CG-CD	6.55	1.61	1.51
1	D	69	GLU	CG-CD	6.54	1.61	1.51
1	F	69	GLU	CG-CD	6.54	1.61	1.51
1	D	93	LYS	CG-CD	6.50	1.74	1.52
1	B	108	GLU	CG-CD	6.38	1.61	1.51
1	B	64	GLU	CD-OE2	-6.34	1.18	1.25
1	C	60	GLU	CG-CD	6.33	1.61	1.51
1	C	17	GLU	CG-CD	6.30	1.61	1.51
1	B	93	LYS	CE-NZ	6.22	1.64	1.49
1	D	20	GLN	CB-CG	6.12	1.69	1.52
1	D	135	TRP	CE3-CZ3	6.09	1.48	1.38
1	E	9	GLN	CB-CG	6.07	1.69	1.52
1	D	101	ASP	CB-CG	6.02	1.64	1.51
1	D	9	GLN	CB-CG	5.99	1.68	1.52
1	B	111	GLU	CG-CD	5.96	1.60	1.51
1	E	90	GLU	CG-CD	5.89	1.60	1.51
1	A	136	GLU	CG-CD	5.88	1.60	1.51
1	B	90	GLU	CB-CG	-5.81	1.41	1.52
1	A	69	GLU	CG-CD	5.77	1.60	1.51
1	E	57	VAL	CB-CG2	5.76	1.65	1.52
1	D	17	GLU	CD-OE2	5.76	1.31	1.25
1	E	81	MSE	SE-CE	-5.73	1.61	1.95
1	E	135	TRP	CB-CG	-5.71	1.40	1.50
1	E	111	GLU	CB-CG	5.65	1.62	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	64	GLU	CD-OE1	-5.62	1.19	1.25
1	D	31	GLU	CG-CD	5.62	1.60	1.51
1	A	73	ARG	CZ-NH2	5.61	1.40	1.33
1	F	69	GLU	CD-OE2	5.56	1.31	1.25
1	B	90	GLU	CG-CD	5.55	1.60	1.51
1	C	111	GLU	CG-CD	5.54	1.60	1.51
1	F	73	ARG	CZ-NH1	5.51	1.40	1.33
1	A	47	LYS	CD-CE	5.46	1.65	1.51
1	F	111	GLU	CG-CD	5.44	1.60	1.51
1	F	136	GLU	CD-OE2	5.40	1.31	1.25
1	F	82	LYS	CD-CE	5.39	1.64	1.51
1	A	73	ARG	CG-CD	5.38	1.65	1.51
1	F	57	VAL	CB-CG2	5.35	1.64	1.52
1	D	48	GLU	CG-CD	5.35	1.59	1.51
1	B	73	ARG	CB-CG	5.33	1.67	1.52
1	A	48	GLU	CD-OE2	5.30	1.31	1.25
1	D	5	GLN	CG-CD	5.30	1.63	1.51
1	A	31	GLU	CG-CD	5.30	1.59	1.51
1	D	28	GLU	CB-CG	-5.29	1.42	1.52
1	A	17	GLU	CD-OE2	5.28	1.31	1.25
1	F	48	GLU	CB-CG	-5.26	1.42	1.52
1	B	3	SER	N-CA	5.21	1.56	1.46
1	F	87	ALA	CA-CB	5.18	1.63	1.52
1	B	48	GLU	CB-CG	-5.16	1.42	1.52
1	D	47	LYS	CD-CE	5.16	1.64	1.51
1	A	41	VAL	CB-CG1	-5.15	1.42	1.52
1	E	17	GLU	CG-CD	5.15	1.59	1.51
1	A	17	GLU	CD-OE1	5.13	1.31	1.25
1	E	132	SER	CB-OG	5.10	1.48	1.42
1	C	120	PHE	CE2-CZ	5.07	1.47	1.37
1	E	20	GLN	CG-CD	5.02	1.62	1.51

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	93	LYS	CD-CE-NZ	12.62	140.72	111.70
1	F	118	ARG	NE-CZ-NH1	10.60	125.60	120.30
1	E	59	ASP	CB-CG-OD1	10.27	127.54	118.30
1	F	81	MSE	CG-SE-CE	-9.13	78.82	98.90
1	E	73	ARG	NE-CZ-NH1	-8.09	116.26	120.30
1	C	51	ASP	CB-CG-OD2	-7.89	111.20	118.30
1	A	73	ARG	NE-CZ-NH2	-7.71	116.45	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	50	ILE	CG1-CB-CG2	-7.56	94.78	111.40
1	F	118	ARG	NE-CZ-NH2	-7.44	116.58	120.30
1	D	59	ASP	CB-CG-OD1	7.15	124.74	118.30
1	D	117	ASP	CB-CG-OD2	-6.50	112.45	118.30
1	E	39	ASP	CB-CG-OD1	6.33	124.00	118.30
1	A	50	ILE	CG1-CB-CG2	-6.10	97.97	111.40
1	A	19	LEU	CB-CG-CD1	6.07	121.31	111.00
1	B	118	ARG	NE-CZ-NH2	-6.04	117.28	120.30
1	E	39	ASP	CB-CG-OD2	-5.94	112.95	118.30
1	D	39	ASP	CB-CG-OD1	5.88	123.60	118.30
1	E	59	ASP	CB-CG-OD2	-5.82	113.06	118.30
1	E	24	ASP	CB-CG-OD2	-5.78	113.10	118.30
1	D	19	LEU	CB-CG-CD1	5.72	120.72	111.00
1	C	103	ARG	CG-CD-NE	-5.52	100.20	111.80
1	A	81	MSE	CG-SE-CE	-5.45	86.91	98.90
1	E	24	ASP	CB-CG-OD1	5.20	122.98	118.30
1	C	39	ASP	CB-CG-OD2	-5.16	113.66	118.30
1	A	79	LEU	CB-CG-CD1	-5.09	102.35	111.00
1	E	73	ARG	NH1-CZ-NH2	5.07	124.98	119.40
1	D	101	ASP	CB-CG-OD1	5.06	122.85	118.30
1	D	73	ARG	NE-CZ-NH1	5.04	122.82	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	101	ASP	Peptide

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1075	0	1043	25	0
1	B	1066	0	1029	17	0
1	C	1118	0	1086	15	0
1	D	1075	0	1043	23	0
1	E	1075	0	1043	11	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	F	1075	0	1042	30	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	F	1	0	0	0	0
3	A	127	0	0	5	0
3	B	133	0	0	1	0
3	C	134	0	0	5	0
3	D	145	0	0	6	1
3	E	113	0	0	3	0
3	F	139	0	0	6	1
All	All	7278	0	6286	109	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (109) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:93:LYS:NZ	1:E:93:LYS:CE	1.68	1.55
1:D:93:LYS:CD	1:D:93:LYS:CG	1.74	1.54
1:F:93:LYS:NZ	1:F:93:LYS:CE	1.68	1.54
1:A:81:MSE:CE	1:A:81:MSE:SE	1.32	1.51
1:D:93:LYS:NZ	1:D:93:LYS:CE	1.73	1.51
1:F:81:MSE:CE	1:F:81:MSE:SE	1.27	1.46
1:C:81:MSE:SE	1:C:81:MSE:CE	1.26	1.45
1:F:81:MSE:CE	1:F:81:MSE:CG	2.19	1.19
1:F:81:MSE:HE3	1:F:81:MSE:SE	1.84	1.14
1:A:81:MSE:SE	1:A:81:MSE:HE1	1.88	1.14
1:C:81:MSE:SE	1:C:81:MSE:HE3	1.83	1.13
1:A:81:MSE:CE	1:A:81:MSE:CG	2.29	1.10
1:A:81:MSE:HE3	1:A:81:MSE:SE	1.88	1.10
1:E:35:TYR:HB2	3:E:2036:HOH:O	1.53	1.09
1:F:81:MSE:HE2	1:F:81:MSE:SE	1.84	1.09
1:C:81:MSE:SE	1:C:81:MSE:HE1	1.83	1.09
1:F:81:MSE:SE	1:F:81:MSE:HE1	1.84	1.07
1:A:81:MSE:HE2	1:A:81:MSE:SE	1.88	1.06
1:C:81:MSE:SE	1:C:81:MSE:HE2	1.83	1.05
1:C:73:ARG:HD2	3:C:2082:HOH:O	1.68	0.93
1:C:81:MSE:CE	1:C:81:MSE:CG	2.49	0.90
1:A:136:GLU:HG2	1:B:135:TRP:CE2	2.09	0.88
1:A:17:GLU:HG3	3:A:2019:HOH:O	1.73	0.87

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:21:LEU:HD22	1:F:98:LYS:HZ2	1.36	0.87
1:E:135:TRP:CE2	1:F:136:GLU:HG2	2.11	0.84
1:B:48:GLU:OE1	1:B:73:ARG:NH1	2.16	0.77
1:F:81:MSE:CE	1:F:81:MSE:HG3	2.16	0.75
1:F:9:GLN:OE1	3:F:2015:HOH:O	2.05	0.74
1:D:93:LYS:HD2	3:D:2106:HOH:O	1.86	0.74
1:F:21:LEU:HD22	1:F:98:LYS:NZ	2.03	0.72
1:A:136:GLU:HG2	1:B:135:TRP:CD2	2.24	0.72
1:F:81:MSE:HE3	1:F:81:MSE:CG	2.11	0.70
1:F:21:LEU:CD2	1:F:98:LYS:HZ2	2.06	0.68
1:F:81:MSE:HE2	1:F:81:MSE:CG	2.11	0.67
1:E:73:ARG:HD2	3:E:2023:HOH:O	1.94	0.67
1:E:35:TYR:C	1:E:35:TYR:CD1	2.68	0.66
1:A:130:GLN:OE1	3:A:2121:HOH:O	2.14	0.66
1:A:81:MSE:HG3	1:A:81:MSE:CE	2.23	0.66
1:A:43:GLU:OE1	3:A:2047:HOH:O	2.12	0.66
1:D:81:MSE:CE	1:D:84:LEU:HD23	2.26	0.65
1:A:69:GLU:HG3	3:A:2080:HOH:O	1.97	0.65
1:C:76:GLN:OE1	3:C:2086:HOH:O	2.14	0.65
1:D:81:MSE:HE3	1:D:131:TYR:CG	2.34	0.62
1:C:81:MSE:HE1	1:C:128:PHE:CD1	2.37	0.60
1:F:81:MSE:HE2	1:F:81:MSE:HG3	1.84	0.58
1:B:21:LEU:HD21	1:B:94:ARG:HB3	1.84	0.58
1:F:47:LYS:C	1:F:47:LYS:HD3	2.24	0.58
1:D:81:MSE:HE1	1:D:84:LEU:HD23	1.85	0.58
1:F:57:VAL:HG12	3:F:2077:HOH:O	2.02	0.58
1:E:5:GLN:HG3	1:E:9:GLN:OE1	2.04	0.57
1:A:136:GLU:CG	1:B:135:TRP:CE2	2.87	0.57
3:E:2096:HOH:O	1:F:93:LYS:HE2	2.05	0.57
1:E:135:TRP:CD2	1:F:136:GLU:HG2	2.40	0.56
1:D:118:ARG:HD2	3:D:2004:HOH:O	2.06	0.55
1:F:5:GLN:NE2	3:F:2009:HOH:O	2.33	0.55
1:D:21:LEU:HD21	1:D:94:ARG:HB3	1.89	0.55
1:F:102:LYS:C	3:F:2116:HOH:O	2.45	0.55
1:F:17:GLU:HG2	1:F:98:LYS:HZ3	1.71	0.55
1:C:48:GLU:HG3	3:C:2055:HOH:O	2.08	0.54
1:A:136:GLU:HG2	1:B:135:TRP:NE1	2.21	0.54
1:B:81:MSE:HE1	1:B:128:PHE:CD1	2.43	0.54
1:D:81:MSE:HA	1:D:81:MSE:HE2	1.89	0.54
1:A:81:MSE:HE1	1:A:128:PHE:CD1	2.44	0.53
1:C:103:ARG:HG3	1:C:107:GLN:HB2	1.90	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:81:MSE:CG	1:A:81:MSE:HE2	2.28	0.51
1:A:136:GLU:HG2	1:B:135:TRP:CG	2.46	0.51
1:F:17:GLU:HG2	1:F:98:LYS:NZ	2.25	0.51
1:F:108:GLU:N	3:F:2119:HOH:O	2.43	0.51
1:A:81:MSE:HE3	1:A:81:MSE:CG	2.28	0.51
1:A:136:GLU:HG2	1:B:135:TRP:CD1	2.46	0.50
1:F:136:GLU:HG3	3:F:2135:HOH:O	2.11	0.49
1:D:81:MSE:CE	1:D:81:MSE:HA	2.44	0.48
1:F:17:GLU:OE2	1:F:98:LYS:HD3	2.14	0.47
1:D:133:ILE:O	1:D:134:SER:C	2.53	0.47
1:D:19:LEU:HB2	3:D:2019:HOH:O	2.14	0.46
1:A:44:ASN:HB3	1:A:73:ARG:HH12	1.80	0.46
1:D:101:ASP:HB2	3:D:2116:HOH:O	2.15	0.46
1:D:22:TRP:HZ3	1:D:50:ILE:HG12	1.81	0.45
1:C:117:ASP:OD2	3:C:2119:HOH:O	2.21	0.45
1:A:19:LEU:HD11	1:A:43:GLU:HG3	1.98	0.45
1:F:47:LYS:HD3	1:F:47:LYS:O	2.17	0.45
1:A:136:GLU:CG	1:B:135:TRP:NE1	2.80	0.44
1:F:21:LEU:HD22	1:F:98:LYS:CE	2.47	0.44
1:A:81:MSE:HB3	1:B:129:ASN:HD22	1.83	0.44
1:D:5:GLN:HG2	1:D:9:GLN:NE2	2.33	0.44
1:E:135:TRP:NE1	1:F:136:GLU:HG2	2.31	0.44
1:F:21:LEU:HB2	1:F:98:LYS:HZ1	1.82	0.44
1:E:108:GLU:HB2	1:E:111:GLU:HG3	2.00	0.44
1:C:103:ARG:HD2	1:C:107:GLN:O	2.18	0.44
1:D:93:LYS:CG	1:D:93:LYS:CE	2.87	0.43
1:B:73:ARG:NH1	1:B:73:ARG:HB2	2.33	0.43
1:C:44:ASN:HB3	1:C:73:ARG:HH12	1.84	0.43
1:E:37:LEU:HD13	1:E:41:VAL:HG11	2.01	0.43
1:B:3:SER:N	3:B:2003:HOH:O	2.53	0.42
1:D:81:MSE:HE2	1:D:84:LEU:HD23	1.98	0.42
1:D:98:LYS:HD3	1:D:98:LYS:HA	1.83	0.42
1:D:37:LEU:HD13	1:D:41:VAL:HG11	2.02	0.42
1:C:98:LYS:HA	1:C:98:LYS:HD3	1.93	0.42
1:E:19:LEU:HD11	1:E:43:GLU:HG3	2.02	0.42
1:D:92:TRP:CZ2	1:D:93:LYS:HE2	2.55	0.41
1:D:19:LEU:HD23	3:D:2023:HOH:O	2.19	0.41
1:D:81:MSE:HE3	1:D:131:TYR:CD2	2.55	0.41
1:B:73:ARG:HH11	1:B:73:ARG:CB	2.34	0.41
1:A:136:GLU:HG3	3:A:2123:HOH:O	2.20	0.41
1:C:17:GLU:HB2	3:C:2023:HOH:O	2.20	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:101:ASP:OD1	3:D:2115:HOH:O	2.22	0.41
1:B:121:SER:N	1:B:122:PRO:HD2	2.36	0.41
1:B:81:MSE:CE	1:B:85:ALA:HB2	2.50	0.41
1:A:136:GLU:OE2	1:B:135:TRP:NE1	2.54	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:D:2086:HOH:O	3:F:2063:HOH:O[2_664]	2.07	0.13

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	125/136 (92%)	123 (98%)	2 (2%)	0	100 100
1	B	124/136 (91%)	120 (97%)	4 (3%)	0	100 100
1	C	132/136 (97%)	130 (98%)	2 (2%)	0	100 100
1	D	125/136 (92%)	124 (99%)	1 (1%)	0	100 100
1	E	125/136 (92%)	122 (98%)	3 (2%)	0	100 100
1	F	125/136 (92%)	124 (99%)	1 (1%)	0	100 100
All	All	756/816 (93%)	743 (98%)	13 (2%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	120/123 (98%)	120 (100%)	0	100 100
1	B	119/123 (97%)	117 (98%)	2 (2%)	60 65
1	C	125/123 (102%)	124 (99%)	1 (1%)	81 86
1	D	120/123 (98%)	118 (98%)	2 (2%)	60 65
1	E	120/123 (98%)	119 (99%)	1 (1%)	81 86
1	F	120/123 (98%)	119 (99%)	1 (1%)	81 86
All	All	724/738 (98%)	717 (99%)	7 (1%)	76 81

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	101	ASP
1	B	111	GLU
1	C	103	ARG
1	D	34	SER
1	D	81	MSE
1	E	102	LYS
1	F	47	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	76	GLN
1	B	129	ASN
1	D	9	GLN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [\(i\)](#)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data i

6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	127/136 (93%)	0.25	7 (5%) 25 24	27, 34, 47, 65	0
1	B	126/136 (92%)	0.29	8 (6%) 20 19	29, 35, 47, 60	0
1	C	132/136 (97%)	0.19	8 (6%) 21 20	27, 34, 47, 53	0
1	D	127/136 (93%)	0.19	5 (3%) 39 38	27, 32, 46, 63	0
1	E	127/136 (93%)	0.35	11 (8%) 10 9	28, 35, 48, 64	0
1	F	127/136 (93%)	0.24	9 (7%) 16 15	28, 33, 45, 62	0
All	All	766/816 (93%)	0.25	48 (6%) 20 19	27, 34, 47, 65	0

All (48) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	35	TYR	4.8
1	C	34	SER	4.4
1	E	34	SER	4.4
1	B	101	ASP	4.3
1	F	101	ASP	4.1
1	B	3	SER	3.9
1	F	135	TRP	3.8
1	D	135	TRP	3.5
1	C	106	ASP	3.5
1	A	135	TRP	3.4
1	F	108	GLU	3.4
1	D	102	LYS	3.3
1	A	101	ASP	3.3
1	E	135	TRP	3.2
1	A	102	LYS	3.1
1	A	11	ILE	3.0
1	E	102	LYS	3.0
1	F	102	LYS	2.9
1	A	34	SER	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	53	LEU	2.8
1	E	88	LEU	2.8
1	E	101	ASP	2.7
1	C	108	GLU	2.7
1	F	11	ILE	2.7
1	C	53	LEU	2.6
1	C	107	GLN	2.6
1	E	53	LEU	2.6
1	B	135	TRP	2.6
1	E	84	LEU	2.5
1	C	88	LEU	2.5
1	D	53	LEU	2.4
1	B	53	LEU	2.4
1	E	70	LEU	2.4
1	B	34	SER	2.4
1	D	50	ILE	2.3
1	B	77	ILE	2.2
1	D	127	ILE	2.2
1	F	49	TYR	2.2
1	B	11	ILE	2.2
1	F	95	LEU	2.2
1	A	49	TYR	2.2
1	B	54	LEU	2.1
1	E	127	ILE	2.1
1	C	104	LEU	2.1
1	C	95	LEU	2.1
1	F	84	LEU	2.0
1	F	74	ALA	2.0
1	E	49	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NA	C	1137	1/1	0.96	0.29	33,33,33,33	0
2	NA	F	1137	1/1	0.97	0.32	39,39,39,39	0
2	NA	B	1137	1/1	0.98	0.24	36,36,36,36	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.