

# Full wwPDB X-ray Structure Validation Report (i)

#### May 14, 2020 - 09:53 pm BST

PDB ID	:	1BRA
$\operatorname{Title}$	:	RELOCATING A NEGATIVE CHARGE IN THE BINDING POCKET OF
		TRYPSIN
Authors	:	Perona, J.J.; Mcgrath, M.E.; Fletterick, R.J.
Deposited on	:	1992-12-17
$\operatorname{Resolution}$	:	2.20  Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution		
	$(\# { m Entries})$	$(\# { m Entries}, { m resolution} \ { m range}({ m \AA}))$		
Clashscore	141614	$5594\ (2.20-2.20)$		
Ramachandran outliers	138981	5503(2.20-2.20)		
Sidechain outliers	138945	5504 (2.20-2.20)		

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain					
1	А	223	62%	32%	5% •			



## 2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 1813 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called TRYPSIN.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
1	А	223	Total 1666	C 1041	N 285	O 326	S 14	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	189	GLY	ASP	CONFLICT	UNP P00763
А	226	ASP	GLY	CONFLICT	UNP P00763

• Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	1	Total Ca 1 1	0	0

• Molecule 3 is BENZAMIDINE (three-letter code: BEN) (formula:  $C_7H_8N_2$ ).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	А	1	Total 9	$\begin{array}{c} \mathrm{C} \\ 7 \end{array}$	N 2	0	0

• Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	137	Total O 137 137	0	0



## 3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are colorcoded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: TRYPSIN





## 4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source	
Space group	I 2 3	Depositor	
Cell constants	124.40Å 124.40Å 124.40Å	Depositor	
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $90.00^{\circ}$	Depositor	
Resolution (Å)	6.00 - 2.20	Depositor	
% Data completeness	(Not available) $(6.00-2.20)$	Depositor	
(in resolution range)	(100 available) (0.00 2.20)		
$R_{merge}$	(Not available)	Depositor	
R <sub>sym</sub>	(Not available)	Depositor	
Refinement program	PROLSQ	Depositor	
$R, R_{free}$	0.156 , (Not available)	Depositor	
Estimated twinning fraction	No twinning to report.	Xtriage	
Total number of atoms	1813	wwPDB-VP	
Average B, all atoms $(Å^2)$	13.0	wwPDB-VP	



# 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, BEN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
		RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	А	0.96	0/1701	2.07	57/2318~(2.5%)	

There are no bond length outliers.

All (57) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	А	62	ARG	NE-CZ-NH2	-17.67	111.47	120.30
1	А	62	ARG	NE-CZ-NH1	16.39	128.49	120.30
1	А	178	ASP	CB-CG-OD1	-10.87	108.52	118.30
1	А	223	ASP	CB-CG-OD2	10.85	128.07	118.30
1	А	190	SER	CB-CA-C	10.37	129.80	110.10
1	А	77	GLU	CG-CD-OE2	-9.86	98.57	118.30
1	А	77	GLU	OE1-CD-OE2	9.46	134.66	123.30
1	А	82	PHE	CB-CG-CD2	-9.20	114.36	120.80
1	А	102	ASP	CB-CG-OD1	9.08	126.47	118.30
1	А	178	ASP	CA-CB-CG	-7.91	96.00	113.40
1	А	240	ASP	CB-CG-OD1	7.65	125.19	118.30
1	А	199	VAL	CG1-CB-CG2	7.28	122.55	110.90
1	А	146	SER	CA-CB-OG	-7.27	91.58	111.20
1	А	48	ASN	N-CA-CB	7.20	123.55	110.60
1	А	95	ASP	CB-CG-OD1	7.00	124.60	118.30
1	А	151	GLU	OE1-CD-OE2	6.87	131.54	123.30
1	А	96	ARG	NE-CZ-NH1	6.58	123.59	120.30
1	А	199	VAL	N-CA-CB	6.58	125.97	111.50
1	А	240	ASP	CB-CG-OD2	-6.55	112.41	118.30
1	А	194	ASP	CB-CG-OD1	-6.47	112.48	118.30
1	A	67	ARG	NE-CZ-NH1	6.38	123.49	120.30
1	А	73	ILE	CA-CB-CG2	6.38	123.65	110.90
1	А	229	THR	CA-CB-OG1	-6.38	95.61	109.00
1	A	95	ASP	CB-CG-OD2	-6.34	112.60	118.30

Continued on next page...



Mol	Chain	$\mathbf{Res}$	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	А	102	ASP	CB-CG-OD2	-6.22	112.70	118.30
1	А	154	LEU	O-C-N	6.11	132.48	122.70
1	А	82	PHE	CB-CG-CD1	6.10	125.07	120.80
1	А	151	GLU	CA-CB-CG	-6.06	100.08	113.40
1	А	227	VAL	CB-CA-C	6.04	122.88	111.40
1	А	60	LYS	N-CA-CB	6.03	121.46	110.60
1	А	42	CYS	CA-C-O	-6.02	107.47	120.10
1	А	199	VAL	CB-CA-C	6.02	122.83	111.40
1	А	120	THR	N-CA-CB	-5.97	98.96	110.30
1	А	244	ALA	CB-CA-C	-5.92	101.22	110.10
1	А	175	LYS	CA-CB-CG	-5.90	100.42	113.40
1	А	71	HIS	N-CA-CB	5.78	121.01	110.60
1	А	229	THR	O-C-N	5.77	131.93	122.70
1	А	193	GLY	CA-C-O	-5.71	110.32	120.60
1	А	37	SER	N-CA-CB	-5.69	101.96	110.50
1	А	239	GLN	CA-CB-CG	-5.58	101.13	113.40
1	А	162	LEU	CB-CA-C	5.51	120.67	110.20
1	А	67	ARG	NE-CZ-NH2	-5.49	117.56	120.30
1	А	203	GLY	CA-C-O	-5.49	110.73	120.60
1	А	78	GLY	CA-C-O	-5.48	110.73	120.60
1	А	67	ARG	CG-CD-NE	-5.44	100.38	111.80
1	А	42	CYS	O-C-N	5.40	132.38	123.20
1	А	245	ASN	CA-C-O	-5.39	108.78	120.10
1	А	120	THR	OG1-CB-CG2	-5.35	97.69	110.00
1	А	49	ASP	CB-CG-OD2	-5.32	113.51	118.30
1	А	81	GLN	N-CA-CB	5.32	120.18	110.60
1	А	178	ASP	CA-C-O	-5.30	108.97	120.10
1	A	155	LEU	CB-CG-CD2	-5.30	102.00	111.00
1	A	62	ARG	CD-NE-CZ	-5.29	116.20	123.60
1	A	223	ASP	CB-CG-OD1	-5.22	113.60	118.30
1	A	166	ALA	CB-CA-C	5.13	117.79	110.10
1	A	162	LEU	N-CA-CB	-5.11	100.18	110.40
1	A	42	CYS	C-N-CA	-5.10	111.60	122.30

Continued from previous page...

There are no chirality outliers.

There are no planarity outliers.

#### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	1666	0	1603	55	0
2	А	1	0	0	0	0
3	А	9	0	7	2	0
4	А	137	0	0	5	0
All	All	1813	0	1610	55	0

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:64:GLN:HE22	1:A:67:ARG:HH21	1.02	0.94
1:A:72:ASN:HD22	1:A:74:ASN:H	1.12	0.90
1:A:64:GLN:NE2	1:A:67:ARG:HH21	1.70	0.89
1:A:72:ASN:ND2	1:A:74:ASN:H	1.72	0.86
1:A:32:SER:OG	1:A:40:HIS:HD2	1.69	0.76
1:A:172:TYR:HB3	1:A:175:LYS:NZ	2.01	0.74
1:A:115:ASN:HD22	1:A:117:ARG:H	1.38	0.72
1:A:79:ASN:HB3	1:A:117:ARG:HG3	1.76	0.68
1:A:72:ASN:HD22	1:A:74:ASN:N	1.91	0.66
1:A:213:VAL:HA	1:A:228:TYR:CD1	2.30	0.66
1:A:172:TYR:HB3	1:A:175:LYS:HZ3	1.62	0.62
1:A:213:VAL:HG22	1:A:228:TYR:HE1	1.63	0.62
1:A:73:ILE:HD13	1:A:141:TRP:CE2	2.39	0.57
1:A:64:GLN:HE21	1:A:67:ARG:HE	1.53	0.57
1:A:64:GLN:NE2	1:A:67:ARG:NH2	2.48	0.56
1:A:34:ASN:ND2	1:A:38:GLY:H	2.02	0.56
1:A:183:VAL:HB	1:A:228:TYR:CE2	2.40	0.56
1:A:74:ASN:ND2	1:A:153:ASP:OD1	2.35	0.53
1:A:117:ARG:HD2	1:A:117:ARG:N	2.23	0.53
1:A:25:ASN:HB3	1:A:117:ARG:HB3	1.91	0.53
1:A:91:HIS:HB2	1:A:103:ILE:HG23	1.89	0.52
1:A:115:ASN:ND2	1:A:117:ARG:H	2.03	0.52
1:A:64:GLN:NE2	1:A:67:ARG:HE	2.07	0.52
1:A:72:ASN:ND2	1:A:74:ASN:N	2.51	0.51
1:A:24:GLU:HG2	1:A:25:ASN:ND2	2.27	0.50
1:A:120:THR:HG22	4:A:272:HOH:O	2.12	0.49
1:A:213:VAL:CG1	3:A:246:BEN:H5	2.42	0.49

Continued on next page...



A 4 1	A.4 a.m. D	Interatomic	Clash
Atom-1	Atom-2	distance $(\text{\AA})$	overlap (Å)
1:A:64:GLN:HE22	1:A:67:ARG:NH2	1.87	0.49
1:A:115:ASN:HD22	1:A:117:ARG:N	2.09	0.47
1:A:117:ARG:HD3	4:A:381:HOH:O	2.15	0.47
1:A:72:ASN:C	1:A:72:ASN:HD22	2.18	0.47
1:A:213:VAL:HA	1:A:228:TYR:HD1	1.78	0.47
1:A:236:ASP:HB2	4:A:384:HOH:O	2.14	0.47
1:A:132:ALA:HA	1:A:162:LEU:HD22	1.96	0.47
1:A:115:ASN:HD21	1:A:117:ARG:HG2	1.81	0.46
1:A:213:VAL:HG13	3:A:246:BEN:H5	1.97	0.46
1:A:135:GLN:NE2	4:A:315:HOH:O	2.50	0.45
1:A:160:ALA:HA	1:A:161:PRO:HD3	1.83	0.44
1:A:110:SER:HB2	1:A:111:PRO:HD2	1.99	0.44
1:A:80:GLU:HB3	1:A:82:PHE:CE1	2.52	0.44
1:A:97:LYS:HE2	1:A:97:LYS:O	2.18	0.44
1:A:37:SER:HB2	1:A:41:PHE:CE1	2.53	0.44
1:A:171:SER:C	1:A:173:PRO:HD3	2.39	0.43
1:A:117:ARG:H	1:A:117:ARG:HD2	1.84	0.42
1:A:56:ALA:HA	1:A:104:MET:HB2	2.01	0.42
1:A:86:ALA:HB2	1:A:109:SER:HA	2.00	0.42
1:A:123:LEU:HB3	1:A:124:PRO:HD2	2.01	0.42
1:A:73:ILE:HD13	1:A:141:TRP:CZ2	2.55	0.41
1:A:199:VAL:HG11	1:A:228:TYR:CD1	2.56	0.41
1:A:37:SER:HB2	1:A:41:PHE:CD1	2.55	0.41
1:A:192:GLN:HB3	4:A:383:HOH:O	2.21	0.41
1:A:221(A):LEU:HA	1:A:222:PRO:HD3	1.88	0.41
1:A:32:SER:OG	1:A:40:HIS:CD2	2.60	0.41
1:A:182:CYS:HA	1:A:226:ASP:O	2.21	0.40
1:A:91:HIS:CG	1:A:92:PRO:HD2	2.57	0.40

Continued from previous page...

There are no symmetry-related clashes.

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	А	221/223 (99%)	217 (98%)	4 (2%)	0	100 100	

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	А	185/185~(100%)	173~(94%)	12~(6%)	17 19	

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	$\mathbf{Res}$	Type
1	А	34	ASN
1	А	37	SER
1	А	72	ASN
1	А	87	LYS
1	А	97	LYS
1	А	115	ASN
1	А	117	ARG
1	А	165	GLN
1	А	175	LYS
1	А	185	LEU
1	А	217	TYR
1	A	227	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	23	GLN
1	А	30	GLN
1	А	34	ASN
1	А	40	HIS
1	А	50	GLN
1	А	64	GLN

Continued on next page...



Continued from previous page...

Mol	Chain	$\mathbf{Res}$	Type
1	А	72	ASN
1	А	84	ASN
1	А	101	ASN
1	А	115	ASN
1	А	135	GLN
1	А	165	GLN
1	А	210	GLN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

#### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

#### 5.6 Ligand geometry (i)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Tune	Chain	Dog	Link	Bond lengths		Bond angles			
WIOI	туре	Unain	nes		Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	BEN	А	246	-	9,9,9	1.72	1 (11%)	7,11,11	1.21	1 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	BEN	А	246	-	-	2/4/4/4	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	$\operatorname{Ideal}(\operatorname{\AA})$
3	А	246	BEN	С1-С	-4.26	1.39	1.47

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
3	А	246	BEN	C1-C-N2	2.44	121.72	118.05

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	А	246	BEN	N2-C-C1-C2
3	А	246	BEN	N1-C-C1-C2

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	А	246	BEN	2	0

#### 5.7 Other polymers (i)

There are no such residues in this entry.

#### 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

## 6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

## 6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

## 6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

