



Full wwPDB X-ray Structure Validation Report ⓘ

May 13, 2020 – 07:12 am BST

PDB ID : 4BRE
Title : Legionella pneumophila NTPDase1 crystal form II (closed) in complex with transition state mimic adenosine 5'phosphovanadate
Authors : Zebisch, M.; Schaefer, P.; Lauble, P.; Straeter, N.
Deposited on : 2013-06-04
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3398 (1.60-1.60)
Clashscore	141614	3665 (1.60-1.60)
Ramachandran outliers	138981	3564 (1.60-1.60)
Sidechain outliers	138945	3563 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	368	 86% 10% ••
1	B	368	 90% 6% •

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 6544 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ECTONUCLEOSIDE TRIPHOSPHATE DIPHOSPHOHYDROLASE I.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	358	2984	1908	486	574	16	0	23	0
1	B	358	2984	1904	489	575	16	0	23	0


There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	34	MET	-	expression tag	UNP Q5ZUA2
A	137	ASP	GLU	conflict	UNP Q5ZUA2
A	149	VAL	ALA	conflict	UNP Q5ZUA2
A	394	LEU	-	expression tag	UNP Q5ZUA2
A	395	GLU	-	expression tag	UNP Q5ZUA2
A	396	HIS	-	expression tag	UNP Q5ZUA2
A	397	HIS	-	expression tag	UNP Q5ZUA2
A	398	HIS	-	expression tag	UNP Q5ZUA2
A	399	HIS	-	expression tag	UNP Q5ZUA2
A	400	HIS	-	expression tag	UNP Q5ZUA2
A	401	HIS	-	expression tag	UNP Q5ZUA2
B	34	MET	-	expression tag	UNP Q5ZUA2
B	137	ASP	GLU	conflict	UNP Q5ZUA2
B	149	VAL	ALA	conflict	UNP Q5ZUA2
B	394	LEU	-	expression tag	UNP Q5ZUA2
B	395	GLU	-	expression tag	UNP Q5ZUA2
B	396	HIS	-	expression tag	UNP Q5ZUA2
B	397	HIS	-	expression tag	UNP Q5ZUA2
B	398	HIS	-	expression tag	UNP Q5ZUA2
B	399	HIS	-	expression tag	UNP Q5ZUA2
B	400	HIS	-	expression tag	UNP Q5ZUA2
B	401	HIS	-	expression tag	UNP Q5ZUA2

- Molecule 2 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Na 1 1	0	0

- Molecule 3 is ADENOSINE-5'-PHOSPHOVANADATE (three-letter code: 50T) (formula: $C_{10}H_{17}N_5O_{11}PV$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	N	O	P			V
3	A	1	28	10	5	11	1	1	0	0
3	B	1	47	20	10	15	1	1	0	1

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total Mg 1 1	0	0
4	A	1	Total Mg 1 1	0	0

- Molecule 5 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: $C_6H_{13}NO_4S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
5	A	1	12	6	1	4	1	0	0
5	B	1	12	6	1	4	1	0	0


- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
6	A	237	240	240	0	3
6	B	232	234	234	0	2

3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: ECTONUCLEOSIDE TRIPHOSPHATE DIPHOSPHOHYDROLASE I

Chain A:  86% 10% ..



- Molecule 1: ECTONUCLEOSIDE TRIPHOSPHATE DIPHOSPHOHYDROLASE I

Chain B:  90% 6% .



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	62.68Å 86.20Å 70.88Å 90.00° 105.94° 90.00°	Depositor
Resolution (Å)	29.33 – 1.60 29.33 – 1.60	Depositor EDS
% Data completeness (in resolution range)	99.9 (29.33-1.60) 99.9 (29.33-1.60)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.62 (at 1.60Å)	Xtrriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.172 , 0.208 0.226 , 0.259	Depositor DCC
R_{free} test set	1404 reflections (1.47%)	wwPDB-VP
Wilson B-factor (Å ²)	18.4	Xtrriage
Anisotropy	0.623	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.44 , 51.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6544	wwPDB-VP
Average B, all atoms (Å ²)	11.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.67% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, 50T, MG, MES

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.90	0/3128	0.93	4/4262 (0.1%)
1	B	0.91	1/3125 (0.0%)	0.95	5/4256 (0.1%)
All	All	0.90	1/6253 (0.0%)	0.94	9/8518 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	189	GLY	N-CA	5.41	1.54	1.46

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	253	ASP	CB-CG-OD2	7.26	124.83	118.30
1	A	356	TYR	CB-CG-CD2	-6.48	117.11	121.00
1	A	277	LYS	CD-CE-NZ	-6.12	97.62	111.70
1	B	56	ARG	NE-CZ-NH1	-6.11	117.24	120.30
1	A	56	ARG	NE-CZ-NH2	-5.92	117.34	120.30
1	B	253	ASP	CB-CG-OD2	5.77	123.49	118.30
1	B	137	ASP	CB-CG-OD1	5.56	123.30	118.30
1	B	265	LYS	CD-CE-NZ	5.42	124.17	111.70
1	B	253	ASP	CB-CA-C	5.32	121.04	110.40

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	391[B]	HIS	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2984	0	2893	26	0
1	B	2984	0	2889	19	0
2	A	1	0	0	0	0
3	A	28	0	12	1	0
3	B	47	0	24	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	12	0	13	1	0
5	B	12	0	13	0	0
6	A	240	0	0	9	0
6	B	234	0	0	3	0
All	All	6544	0	5844	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:333[B]:GLN:OE1	1:A:335[B]:ASP:OD1	1.55	1.24
1:B:323[B]:GLN:OE1	1:B:327:ASN:ND2	1.73	1.19
1:A:392[B]:ARG:NH2	6:A:2238:HOH:O	1.90	1.03
1:A:283[B]:GLN:OE1	6:A:2168:HOH:O	2.02	0.76
1:B:323[B]:GLN:CD	1:B:327:ASN:HD22	1.92	0.73
1:A:293[A]:GLU:OE1	6:A:2181:HOH:O	2.05	0.73
1:A:137:ASP:HB3	1:A:141:ARG:HH12	1.57	0.68
1:A:231[B]:GLN:HG3	6:A:2091:HOH:O	1.94	0.67
1:B:231[B]:GLN:HG3	6:B:2103:HOH:O	1.95	0.65

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:231[B]:GLN:CG	6:B:2103:HOH:O	2.45	0.64
1:A:109:HIS:HB3	1:A:145:GLN:HB3	1.82	0.60
1:A:231[B]:GLN:CG	6:A:2091:HOH:O	2.48	0.60
1:A:200:LYS:HG3	6:A:2096:HOH:O	2.01	0.59
1:B:231[A]:GLN:HG3	1:B:353:LEU:HD21	1.85	0.58
1:A:91:ASN:O	1:A:95:ILE:HG13	2.04	0.58
1:B:231[B]:GLN:OE1	1:B:357[B]:TYR:OH	2.23	0.56
1:B:231[B]:GLN:HG2	1:B:356:TYR:CE2	2.41	0.56
1:B:303:LEU:HD13	1:B:357[A]:TYR:HE2	1.71	0.56
3:A:1396:50T:N1	6:A:2187:HOH:O	2.34	0.54
1:A:137:ASP:HB3	1:A:141:ARG:NH1	2.23	0.54
1:B:161:LEU:HD22	1:B:210[A]:VAL:HG11	1.91	0.53
1:A:323[B]:GLN:OE1	1:A:327:ASN:ND2	2.42	0.52
1:A:77:TRP:CD2	1:A:107:PRO:HD3	2.45	0.51
5:A:1398:MES:H32	5:A:1398:MES:H81	1.52	0.48
1:A:328:GLN:HG3	6:A:2194:HOH:O	2.14	0.48
1:A:136:LEU:C	1:A:136:LEU:HD23	2.33	0.48
1:A:115:PHE:O	1:A:151:ALA:HA	2.15	0.45
1:A:231[B]:GLN:OE1	1:A:357[B]:TYR:OH	2.34	0.45
1:B:76:ILE:HD12	1:B:108:ILE:HD13	1.98	0.45
1:A:231[B]:GLN:HG2	1:A:356:TYR:CE2	2.53	0.44
1:A:111:ILE:HB	1:A:112:PRO:CD	2.47	0.44
1:B:231[B]:GLN:CD	1:B:357[B]:TYR:OH	2.56	0.44
1:B:231[B]:GLN:HG2	1:B:356:TYR:HE2	1.81	0.43
1:A:392[B]:ARG:CZ	6:A:2238:HOH:O	2.50	0.43
1:B:393:ALA:HB2	6:B:2232:HOH:O	2.19	0.43
1:A:111:ILE:HB	1:A:112:PRO:HD2	2.00	0.42
1:A:179:ASN:O	1:A:179:ASN:OD1	2.37	0.42
1:A:57:LEU:HD22	1:A:103:LEU:HD23	2.02	0.41
1:B:231[A]:GLN:HE22	1:B:346:TYR:HE2	1.68	0.41
1:B:234[B]:MET:HG2	1:B:356:TYR:CE1	2.55	0.41
1:B:210[B]:VAL:HG23	1:B:221:LEU:HB2	2.03	0.41
1:B:265:LYS:HE3	1:B:265:LYS:HB3	1.88	0.41
1:A:394:LEU:HA	1:A:394:LEU:HD23	1.81	0.41
1:A:85:PHE:O	1:A:88:ILE:HG12	2.21	0.41
1:A:103:LEU:HD12	1:A:139:TRP:CH2	2.56	0.41
1:B:231[A]:GLN:HG3	1:B:353:LEU:CD2	2.51	0.40
1:B:58:HIS:NE2	1:B:383:ASP:OD2	2.54	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	379/368 (103%)	365 (96%)	14 (4%)	0	100	100
1	B	379/368 (103%)	365 (96%)	14 (4%)	0	100	100
All	All	758/736 (103%)	730 (96%)	28 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	341/328 (104%)	330 (97%)	11 (3%)	39	15
1	B	341/328 (104%)	336 (98%)	5 (2%)	65	44
All	All	682/656 (104%)	666 (98%)	16 (2%)	52	25

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	40	GLU
1	A	56	ARG
1	A	95	ILE
1	A	109	HIS
1	A	180	LYS
1	A	200	LYS
1	A	212	LEU
1	A	234[A]	MET

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	234[B]	MET
1	A	265	LYS
1	A	386	ILE
1	B	56	ARG
1	B	95	ILE
1	B	147[A]	GLN
1	B	147[B]	GLN
1	B	265	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	109	HIS
1	A	179	ASN
1	A	327	ASN
1	A	328	GLN
1	B	319	GLN
1	B	327	ASN
1	B	370	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 3 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	MES	A	1398	-	12,12,12	2.04	2 (16%)	14,16,16	3.25	7 (50%)
3	50T	A	1396	4	21,30,30	1.42	3 (14%)	22,49,49	1.51	2 (9%)
5	MES	B	1397	-	12,12,12	2.01	1 (8%)	14,16,16	3.37	5 (35%)
3	50T	B	1395[B]	-	21,30,30	1.31	2 (9%)	22,49,49	2.59	5 (22%)
3	50T	B	1395[A]	-	21,30,30	1.39	2 (9%)	22,49,49	1.71	5 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	MES	A	1398	-	-	5/6/14/14	0/1/1/1
3	50T	A	1396	4	-	1/6/33/33	0/3/3/3
5	MES	B	1397	-	-	3/6/14/14	0/1/1/1
3	50T	B	1395[B]	-	-	1/6/33/33	0/3/3/3
3	50T	B	1395[A]	-	-	1/6/33/33	0/3/3/3

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	1397	MES	C8-S	-6.46	1.68	1.77
5	A	1398	MES	C8-S	-6.13	1.68	1.77
3	B	1395[B]	50T	C6-N6	4.55	1.50	1.34
3	A	1396	50T	C6-N6	3.76	1.47	1.34
3	B	1395[A]	50T	C6-N6	3.68	1.47	1.34
3	A	1396	50T	C2-N3	3.32	1.37	1.32
3	B	1395[A]	50T	C2-N3	2.89	1.36	1.32
3	A	1396	50T	O3'-C3'	2.13	1.48	1.43
3	B	1395[B]	50T	C6-C5	2.10	1.51	1.43
5	A	1398	MES	O1S-S	2.00	1.50	1.45

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	1397	MES	O1S-S-C8	-8.39	96.81	106.92

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1395[B]	50T	O4'-C1'-C2'	-7.98	95.27	106.93
5	A	1398	MES	O2S-S-C8	7.12	115.49	106.92
5	A	1398	MES	O3S-S-C8	-6.78	94.81	105.77
5	B	1397	MES	O3S-S-C8	6.42	116.15	105.77
3	B	1395[B]	50T	N3-C2-N1	-5.47	120.13	128.68
3	B	1395[A]	50T	N3-C2-N1	-4.59	121.51	128.68
3	A	1396	50T	N3-C2-N1	-4.51	121.62	128.68
5	B	1397	MES	O2S-S-C8	4.40	112.21	106.92
5	A	1398	MES	C5-N4-C3	4.20	118.29	108.83
3	B	1395[B]	50T	C5-C6-N6	4.11	126.60	120.35
3	B	1395[B]	50T	C3'-C2'-C1'	-3.53	95.67	100.98
3	A	1396	50T	O4'-C1'-C2'	-3.51	101.80	106.93
3	B	1395[A]	50T	N6-C6-N1	3.51	125.85	118.57
5	B	1397	MES	O3S-S-O2S	-3.06	103.80	111.27
3	B	1395[A]	50T	C5-C6-N6	-3.03	115.74	120.35
5	A	1398	MES	O1S-S-C8	3.00	110.53	106.92
3	B	1395[B]	50T	C2-N1-C6	2.99	123.87	118.75
3	B	1395[A]	50T	O4'-C1'-C2'	-2.87	102.73	106.93
5	A	1398	MES	O3S-S-O1S	2.66	117.78	111.27
5	A	1398	MES	C2-C3-N4	2.65	114.12	110.10
5	B	1397	MES	C6-O1-C2	2.45	118.08	109.89
5	A	1398	MES	O2S-S-O1S	-2.09	106.70	113.95
3	B	1395[A]	50T	C1'-N9-C4	-2.00	123.12	126.64

There are no chirality outliers.

All (11) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	B	1397	MES	C8-C7-N4-C5
5	A	1398	MES	N4-C7-C8-S
5	A	1398	MES	C7-C8-S-O2S
5	A	1398	MES	C7-C8-S-O3S
5	B	1397	MES	N4-C7-C8-S
5	B	1397	MES	C8-C7-N4-C3
5	A	1398	MES	C7-C8-S-O1S
3	A	1396	50T	O4'-C4'-C5'-O5'
3	B	1395[B]	50T	O4'-C4'-C5'-O5'
3	B	1395[A]	50T	O4'-C4'-C5'-O5'
5	A	1398	MES	C8-C7-N4-C3


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	1398	MES	1	0
3	A	1396	50T	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.


6.3 Carbohydrates [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands [i](#)




Unable to reproduce the depositors R factor - this section is therefore empty.

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around 50T B 1395 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around 50T B 1395 (A):**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers

Unable to reproduce the depositor's R factor - this section is therefore empty.