



Full wwPDB X-ray Structure Validation Report i

Jan 30, 2023 – 12:34 pm GMT

PDB ID : 8BSG
Title : COMPLEX OF LEPORINE SERUM ALBUMIN WITH DICLOFENAC
Authors : Bujacz, A.; Talaj, J.; Zielinski, K.
Deposited on : 2022-11-25
Resolution : 1.89 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The types of validation reports are described at
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.31.3
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.31.3


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.89 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



2 Entry composition [\(i\)](#)


There are 7 unique types of molecules in this entry. The entry contains 5163 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Albumin.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	584	4669	2958	782	892	37	0	7	0

- Molecule 2 is 2-[2,6-DICHLOROPHENYL]AMINO]BENZENEACETIC ACID (three-letter code: DIF) (formula: C₁₄H₁₁Cl₂NO₂) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	Cl	N	O		
2	A	1	19	14	2	1	2	0	0
2	A	1	19	14	2	1	2	0	0

- Molecule 3 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0

- Molecule 4 is (20S)-2,5,8,11,14,17-HEXAMETHYL-3,6,9,12,15,18-HEXAOXAHENICOSAN E-1,20-DIOL (three-letter code: POG) (formula: C₂₁H₄₄O₈).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			22	16	6		
4	A	1	Total	C	O	0	0
			14	10	4		

- Molecule 5 is S-1,2-PROPANEDIOL (three-letter code: PGO) (formula: C₃H₈O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			5	3	2		
5	A	1	Total	C	O	0	0
			5	3	2		

- Molecule 6 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total Na 1 1	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	385	Total O 385 385	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Albumin



4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	74.78Å 80.13Å 104.10Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.65 – 1.89 43.65 – 1.89	Depositor EDS
% Data completeness (in resolution range)	99.7 (43.65-1.89) 99.7 (43.65-1.89)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ ¹	1.78 (at 1.89Å)	Xtriage
Refinement program	REFMAC 5.8.0103	Depositor
R , R_{free}	0.198 , 0.252 0.199 , 0.253	Depositor DCC
R_{free} test set	2533 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	35.7	Xtriage
Anisotropy	0.075	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 56.2	EDS
L-test for twinning ²	$< L > = 0.48$, $< L^2 > = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5163	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.99% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [\(i\)](#)

5.1 Standard geometry [\(i\)](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PGO, POG, NA, ACT, DIF

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.00	5/4792 (0.1%)	1.00	10/6479 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	98	ARG	CB-CG	6.07	1.69	1.52
1	A	468	PRO	N-CD	5.65	1.55	1.47
1	A	148	TYR	CG-CD2	5.50	1.46	1.39
1	A	48	ASP	CB-CG	5.39	1.63	1.51
1	A	517	PRO	N-CD	5.12	1.55	1.47

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	114	PRO	CB-CA-C	7.25	130.13	112.00
1	A	555	LEU	CA-CB-CG	7.22	131.90	115.30
1	A	114	PRO	N-CA-C	-6.99	93.92	112.10
1	A	30	TYR	CB-CG-CD2	-6.63	117.02	121.00
1	A	481	LEU	CB-CG-CD2	-6.13	100.58	111.00
1	A	30	TYR	CB-CA-C	-5.88	98.64	110.40
1	A	336	ARG	CB-CG-CD	-5.79	96.54	111.60
1	A	30	TYR	CB-CG-CD1	5.76	124.46	121.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	144	ARG	CB-CG-CD	-5.42	97.50	111.60
1	A	279	CYS	CA-CB-SG	-5.23	104.59	114.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	502	PHE	Peptide

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4669	0	4584	80	0
2	A	38	0	20	8	0
3	A	24	0	18	0	0
4	A	36	0	48	2	0
5	A	10	0	16	1	0
6	A	1	0	0	0	0
7	A	385	0	0	8	0
All	All	5163	0	4686	81	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (81) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (\AA)	Clash overlap (\AA)
1:A:433:VAL:HG22	1:A:452[B]:TYR:CD2	1.54	1.39
1:A:433:VAL:HG22	1:A:452[B]:TYR:CE2	1.98	0.98
1:A:433:VAL:CG2	1:A:452[B]:TYR:CD2	2.47	0.97
1:A:433:VAL:CG2	1:A:452[B]:TYR:HD2	1.83	0.90
4:A:610:POG:H173	4:A:610:POG:H181	1.62	0.81
1:A:433:VAL:HG22	1:A:452[B]:TYR:HD2	0.96	0.80
1:A:433:VAL:CG2	1:A:452[B]:TYR:CE2	2.65	0.78

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:449:VAL:HG12	2:A:601:DIF:H10	1.64	0.78
1:A:283:ILE:H	1:A:283:ILE:HD12	1.51	0.74
1:A:37[B]:GLU:HG2	7:A:877:HOH:O	1.89	0.71
1:A:294:HIS:HD2	7:A:935:HOH:O	1.71	0.71
1:A:372:LYS:O	1:A:376:GLU:HG3	1.94	0.67
1:A:86:ASP:HA	1:A:89:ASP:HB2	1.78	0.66
1:A:449:VAL:HG12	2:A:601:DIF:C10	2.29	0.62
1:A:384:PRO:HB2	1:A:446:LEU:HD11	1.81	0.61
1:A:344:VAL:CG1	1:A:450:GLU:HG3	2.30	0.61
1:A:503:ASN:HB2	1:A:505:GLU:HG2	1.82	0.61
1:A:547:VAL:HG21	1:A:583:LEU:HG	1.83	0.61
1:A:194:ALA:HB1	1:A:455:VAL:HG13	1.82	0.61
1:A:27:PHE:HB3	1:A:39:HIS:HD2	1.65	0.59
1:A:432:LYS:HE2	7:A:812:HOH:O	2.04	0.57
1:A:449:VAL:CG1	2:A:601:DIF:C9	2.82	0.57
1:A:509:PHE:CE1	1:A:513:ILE:HB	2.40	0.56
1:A:77:LEU:HG	1:A:79:SER:HB2	1.88	0.56
1:A:405:ASN:HA	1:A:408:LEU:HD12	1.88	0.56
1:A:283:ILE:H	1:A:283:ILE:CD1	2.18	0.55
1:A:449:VAL:CG1	2:A:601:DIF:C10	2.84	0.55
1:A:344:VAL:HG13	1:A:450:GLU:HG3	1.87	0.55
1:A:436:LYS:HD3	1:A:452[A]:TYR:CE1	2.42	0.54
1:A:50:ALA:O	1:A:54:VAL:HG23	2.08	0.54
1:A:509:PHE:CZ	1:A:513:ILE:HB	2.44	0.53
1:A:59:ALA:HB3	1:A:62:CYS:SG	2.48	0.53
1:A:190:LEU:HB3	5:A:611:PGO:H12	1.89	0.53
1:A:82:ASP:HA	1:A:84:TYR:O	2.09	0.53
1:A:503:ASN:C	1:A:505:GLU:H	2.12	0.53
1:A:239:THR:O	1:A:243:LYS:HG3	2.09	0.52
1:A:77:LEU:HB3	1:A:80:LEU:HD22	1.90	0.52
1:A:504:ALA:O	1:A:507:PHE:HB3	2.10	0.52
1:A:113:PRO:C	1:A:114:PRO:O	2.42	0.51
1:A:550:GLU:OE2	1:A:578:SER:HB3	2.11	0.51
1:A:433:VAL:CG2	1:A:452[B]:TYR:HE2	2.20	0.51
1:A:449:VAL:CG1	2:A:601:DIF:H9	2.40	0.50
1:A:119:GLU:HA	1:A:119:GLU:OE1	2.11	0.50
1:A:545:LYS:HE2	2:A:602:DIF:CL2	2.50	0.49
1:A:283:ILE:HD12	1:A:283:ILE:N	2.24	0.48
1:A:433:VAL:HG23	1:A:452[B]:TYR:HE2	1.79	0.48
1:A:294:HIS:CD2	1:A:294:HIS:H	2.31	0.47
1:A:328:GLY:HA3	4:A:609:POG:H161	1.97	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:516:LEU:HB3	1:A:520:GLU:HB2	1.98	0.46
1:A:554:LEU:HD21	1:A:571:GLU:HB3	1.97	0.46
1:A:547:VAL:HG23	1:A:582:THR:HB	1.98	0.45
1:A:552:THR:HG22	7:A:975:HOH:O	2.15	0.45
1:A:449:VAL:HG11	2:A:601:DIF:C9	2.47	0.45
1:A:209:ARG:HG3	7:A:870:HOH:O	2.17	0.45
1:A:225:LYS:HE2	1:A:297:GLU:O	2.16	0.45
1:A:243:LYS:HB3	1:A:243:LYS:HE3	1.69	0.45
1:A:475:LYS:O	1:A:479:GLU:HB2	2.17	0.45
1:A:76:ALA:O	1:A:78:PRO:HD3	2.17	0.44
1:A:263:TYR:CE1	1:A:267:HIS:CE1	3.05	0.44
1:A:135:PHE:CZ	1:A:162:LYS:HB2	2.53	0.43
1:A:441:PRO:HD2	1:A:444:GLU:OE2	2.19	0.43
1:A:35:PRO:HB2	1:A:37[B]:GLU:HG3	1.99	0.43
1:A:79:SER:O	1:A:81:ARG:HG3	2.18	0.42
1:A:384:PRO:CB	1:A:446:LEU:HD11	2.47	0.42
1:A:358[B]:LYS:NZ	7:A:705:HOH:O	2.51	0.42
1:A:86:ASP:HA	1:A:89:ASP:CB	2.48	0.42
1:A:409:VAL:HG21	2:A:602:DIF:CL4	2.56	0.42
1:A:73:LYS:HD3	1:A:73:LYS:HA	1.87	0.42
1:A:547:VAL:CG2	1:A:582:THR:HB	2.50	0.42
1:A:482:VAL:HG22	7:A:927:HOH:O	2.20	0.42
1:A:1:GLU:O	1:A:3:HIS:HD2	2.04	0.41
1:A:408:LEU:HD22	1:A:530:VAL:HG23	2.02	0.41
1:A:442:GLU:HA	1:A:445:ARG:HB3	2.01	0.41
1:A:555:LEU:O	1:A:559:CYS:HB2	2.20	0.41
1:A:518:GLU:HA	1:A:521:ARG:HD3	2.03	0.41
1:A:39:HIS:O	1:A:43:VAL:HG23	2.21	0.41
1:A:384:PRO:HB2	1:A:446:LEU:CD1	2.48	0.41
1:A:548:VAL:HG12	7:A:713:HOH:O	2.20	0.40
1:A:168:CYS:HB2	1:A:178:LEU:HD13	2.03	0.40
1:A:274:HIS:ND1	1:A:294:HIS:CE1	2.90	0.40
1:A:536:LYS:HG3	1:A:583:LEU:HB3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	589/584 (101%)	561 (95%)	28 (5%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	517/510 (101%)	504 (98%)	13 (2%)	47 41

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	GLU
1	A	80	LEU
1	A	86	ASP
1	A	452[A]	TYR
1	A	452[B]	TYR
1	A	453	LEU
1	A	455	VAL
1	A	502	PHE
1	A	503	ASN
1	A	506	THR
1	A	509	PHE
1	A	555	LEU
1	A	559	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	39	HIS
1	A	294	HIS
1	A	318	ASN
1	A	378	GLN
1	A	397	GLN
1	A	503	ASN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [\(i\)](#)

Of 13 ligands modelled in this entry, 1 is monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	POG	A	609	-	16,21,28	0.69	0	19,25,34	1.32	2 (10%)
2	DIF	A	602	-	20,20,20	2.76	5 (25%)	27,27,27	1.44	5 (18%)
3	ACT	A	604	-	3,3,3	0.86	0	3,3,3	0.82	0
4	POG	A	610	-	10,13,28	0.96	0	11,15,34	1.56	2 (18%)
3	ACT	A	608	-	3,3,3	0.67	0	3,3,3	1.16	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	PGO	A	612	-	3,4,4	0.35	0	1,4,4	0.42	0
3	ACT	A	607	-	3,3,3	0.78	0	3,3,3	0.77	0
2	DIF	A	601	-	20,20,20	2.60	3 (15%)	27,27,27	1.76	5 (18%)
3	ACT	A	603	-	3,3,3	0.64	0	3,3,3	0.84	0
3	ACT	A	606	-	3,3,3	0.84	0	3,3,3	0.56	0
5	PGO	A	611	-	3,4,4	0.13	0	1,4,4	0.41	0
3	ACT	A	605	-	3,3,3	0.78	0	3,3,3	0.85	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	POG	A	609	-	-	9/24/24/32	-
2	DIF	A	602	-	-	0/8/8/8	0/2/2/2
4	POG	A	610	-	-	9/14/14/32	-
5	PGO	A	612	-	-	0/2/2/2	-
2	DIF	A	601	-	-	0/8/8/8	0/2/2/2
5	PGO	A	611	-	-	2/2/2/2	-

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	DIF	C8-C7	8.50	1.50	1.40
2	A	602	DIF	C8-C7	8.17	1.50	1.40
2	A	602	DIF	C3-C4	5.52	1.48	1.40
2	A	602	DIF	C3-C2	5.18	1.47	1.40
2	A	601	DIF	C3-C2	5.15	1.47	1.40
2	A	601	DIF	C3-C4	5.02	1.47	1.40
2	A	602	DIF	C4-CL4	2.72	1.80	1.73
2	A	602	DIF	C2-CL2	2.68	1.80	1.73

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	DIF	C9-C8-C7	-4.24	114.75	119.76
2	A	601	DIF	C7-C8-N1	3.93	121.72	118.60
4	A	609	POG	C15-O7-C14	3.44	120.16	115.02
2	A	602	DIF	C3-C2-CL2	-3.18	115.64	119.27
2	A	602	DIF	C1-C2-CL2	3.08	124.58	118.41

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	DIF	C12-C7-C8	2.90	121.19	118.32
4	A	610	POG	C15-O7-C14	2.86	119.29	115.02
4	A	609	POG	C13-O6-C12	-2.86	110.76	115.02
2	A	601	DIF	O2-C14-C13	-2.66	115.44	123.04
4	A	610	POG	O5-C11-C12	-2.62	104.83	111.78
2	A	602	DIF	C5-C4-CL4	2.62	123.65	118.41
2	A	601	DIF	C3-C2-CL2	-2.60	116.30	119.27
2	A	602	DIF	O2-C14-C13	-2.10	117.05	123.04
2	A	602	DIF	C3-C4-CL4	-2.09	116.89	119.27

There are no chirality outliers.

All (20) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	609	POG	O3-C7-C8-O4
4	A	609	POG	O3-C7-C8-C20
4	A	609	POG	O5-C10-C9-O4
4	A	609	POG	C19-C10-C9-O4
4	A	609	POG	O5-C11-C12-C18
4	A	610	POG	C15-C5-O2-C3
4	A	610	POG	O5-C11-C12-O6
4	A	610	POG	C14-C13-O6-C12
4	A	610	POG	O6-C13-C14-O7
4	A	610	POG	O6-C13-C14-C17
5	A	611	PGO	O1-C1-C2-O2
4	A	610	POG	C18-C12-O6-C13
4	A	610	POG	C17-C14-O7-C15
4	A	609	POG	C12-C11-O5-C10
4	A	609	POG	C5-C15-O7-C14
4	A	610	POG	C16-C5-O2-C3
4	A	609	POG	O5-C11-C12-O6
5	A	611	PGO	O1-C1-C2-C3
4	A	610	POG	O5-C11-C12-C18
4	A	609	POG	C17-C14-O7-C15

There are no ring outliers.

5 monomers are involved in 11 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	609	POG	1	0
2	A	602	DIF	2	0


Continued on next page...


Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	610	POG	1	0
2	A	601	DIF	6	0
5	A	611	PGO	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data i

6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	584/584 (100%)	0.59	63 (10%) 5 6	22, 43, 111, 141	0

All (63) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	568	PHE	14.5
1	A	506	THR	12.1
1	A	560	SER	12.1
1	A	566	ALA	11.6
1	A	559	CYS	10.0
1	A	570	VAL	10.0
1	A	509	PHE	9.4
1	A	510	HIS	9.0
1	A	507	PHE	8.9
1	A	555	LEU	8.8
1	A	569	ALA	8.3
1	A	563	ASP	7.7
1	A	504	ALA	7.0
1	A	508	THR	6.5
1	A	567	CYS	6.4
1	A	572	GLY	5.8
1	A	511	ALA	5.6
1	A	88	ALA	5.5
1	A	171	ALA	5.5
1	A	502	PHE	5.3
1	A	513	ILE	5.1
1	A	551	PHE	5.0
1	A	172	ALA	4.9
1	A	564	LYS	4.8
1	A	515	THR	4.7
1	A	446	LEU	4.7
1	A	565	GLU	4.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	176	ALA	4.2
1	A	562	GLU	4.1
1	A	558	CYS	3.8
1	A	557	LYS	3.8
1	A	514	CYS	3.8
1	A	503	ASN	3.7
1	A	553	ALA	3.6
1	A	80	LEU	3.6
1	A	173	ASP	3.5
1	A	94	LYS	3.5
1	A	573	PRO	3.5
1	A	576	VAL	3.4
1	A	87	VAL	3.4
1	A	554	LEU	3.3
1	A	561	ALA	3.3
1	A	85	GLY	3.1
1	A	574	LYS	3.0
1	A	92	GLU	2.9
1	A	2	ALA	2.9
1	A	83	THR	2.8
1	A	76	ALA	2.7
1	A	575	LEU	2.6
1	A	115	PHE	2.5
1	A	581	ALA	2.5
1	A	516	LEU	2.5
1	A	556	ASP	2.4
1	A	96	PRO	2.4
1	A	98	ARG	2.2
1	A	1	GLU	2.2
1	A	170	GLU	2.2
1	A	505	GLU	2.2
1	A	571	GLU	2.1
1	A	93	LYS	2.1
1	A	548	VAL	2.1
1	A	3	HIS	2.1
1	A	84	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)


There are no monosaccharides in this entry.


6.4 Ligands [\(i\)](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	DIF	A	602	19/19	0.72	0.21	63,73,78,89	0
3	ACT	A	606	4/4	0.78	0.20	55,65,69,72	0
3	ACT	A	608	4/4	0.78	0.23	60,61,62,63	0
5	PGO	A	612	5/5	0.81	0.37	73,75,77,79	0
3	ACT	A	607	4/4	0.84	0.18	63,67,69,72	0
4	POG	A	610	14/29	0.85	0.18	47,51,53,54	0
5	PGO	A	611	5/5	0.85	0.18	60,64,66,70	0
4	POG	A	609	22/29	0.85	0.18	51,65,72,74	0
3	ACT	A	605	4/4	0.87	0.20	56,62,67,68	0
2	DIF	A	601	19/19	0.95	0.12	43,46,52,53	0
3	ACT	A	604	4/4	0.96	0.16	29,30,31,31	0
3	ACT	A	603	4/4	0.97	0.14	36,37,40,41	0
6	NA	A	613	1/1	0.99	0.14	43,43,43,43	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.