



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 19, 2023 – 05:37 AM EDT

PDB ID : 5BU1
Title : Structure of the truncated C-terminal domain of lpg1496 from Legionella pneumophila
Authors : Wong, K.; Kozlov, G.; Gehring, K.; Montreal-Kingston Bacterial Structural Genomics Initiative (BSGI)
Deposited on : 2015-06-03
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3398 (1.60-1.60)
Clashscore	141614	3665 (1.60-1.60)
Ramachandran outliers	138981	3564 (1.60-1.60)
Sidechain outliers	138945	3563 (1.60-1.60)
RSRZ outliers	127900	3321 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	297	 78% 14% . .

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2727 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called LPG1496.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	284	2306	1482	388	424	12	0	11	0

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	292	MET	-	initiating methionine	UNP G8UY02
A	581	LEU	-	expression tag	UNP G8UY02
A	582	GLU	-	expression tag	UNP G8UY02
A	583	HIS	-	expression tag	UNP G8UY02
A	584	HIS	-	expression tag	UNP G8UY02
A	585	HIS	-	expression tag	UNP G8UY02
A	586	HIS	-	expression tag	UNP G8UY02
A	587	HIS	-	expression tag	UNP G8UY02
A	588	HIS	-	expression tag	UNP G8UY02

- Molecule 2 is MALONATE ION (three-letter code: MLI) (formula: $C_3H_2O_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			7	3	4		


- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	414	Total	O	0	0
			414	414		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: LPG1496

Chain A:  78% 14%



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	60.89Å 71.47Å 77.00Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.61 – 1.60 29.61 – 1.60	Depositor EDS
% Data completeness (in resolution range)	99.8 (29.61-1.60) 99.8 (29.61-1.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	7.29 (at 1.60Å)	Xtrriage
Refinement program	REFMAC 5.8.0069	Depositor
R, R_{free}	0.160 , 0.204 0.160 , 0.204	Depositor DCC
R_{free} test set	2264 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	15.0	Xtrriage
Anisotropy	0.008	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 47.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	2727	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.00% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: MLI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.26	8/2398 (0.3%)	1.60	30/3241 (0.9%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	363	ARG	CZ-NH2	-11.12	1.18	1.33
1	A	363	ARG	CD-NE	-8.25	1.32	1.46
1	A	561	GLN	CD-NE2	6.44	1.49	1.32
1	A	416	ARG	NE-CZ	5.66	1.40	1.33
1	A	450	GLU	CD-OE2	-5.57	1.19	1.25
1	A	450	GLU	CD-OE1	5.46	1.31	1.25
1	A	353	SER	CB-OG	5.38	1.49	1.42
1	A	412	SER	CA-CB	5.15	1.60	1.52

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	363	ARG	NE-CZ-NH1	-36.21	102.19	120.30
1	A	363	ARG	CD-NE-CZ	-17.60	98.95	123.60
1	A	528	ARG	NE-CZ-NH1	15.39	128.00	120.30
1	A	528	ARG	NE-CZ-NH2	-13.07	113.77	120.30
1	A	363	ARG	NH1-CZ-NH2	11.48	132.03	119.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	416	ARG	NE-CZ-NH1	9.85	125.22	120.30
1	A	373	ARG	NE-CZ-NH1	-9.31	115.64	120.30
1	A	562	ASP	CB-CG-OD2	8.66	126.09	118.30
1	A	545	ASP	CB-CG-OD1	7.92	125.43	118.30
1	A	528	ARG	CD-NE-CZ	7.88	134.63	123.60
1	A	480	TYR	CB-CG-CD1	-7.55	116.47	121.00
1	A	505	ASP	CB-CG-OD2	-7.34	111.69	118.30
1	A	541	ASP	CB-CG-OD1	7.03	124.63	118.30
1	A	523	LYS	CD-CE-NZ	-6.79	96.09	111.70
1	A	363	ARG	NE-CZ-NH2	6.53	123.57	120.30
1	A	448	PHE	CB-CG-CD2	-6.36	116.35	120.80
1	A	510	ARG	NE-CZ-NH1	6.30	123.45	120.30
1	A	397	LEU	CB-CG-CD1	5.89	121.02	111.00
1	A	373	ARG	NE-CZ-NH2	5.88	123.24	120.30
1	A	553	ARG	NE-CZ-NH1	-5.64	117.48	120.30
1	A	528	ARG	CG-CD-NE	-5.61	100.02	111.80
1	A	411	PHE	CZ-CE2-CD2	-5.56	113.42	120.10
1	A	411	PHE	CG-CD1-CE1	-5.54	114.70	120.80
1	A	397	LEU	CA-CB-CG	5.50	127.96	115.30
1	A	329	ASP	CB-CG-OD1	5.37	123.14	118.30
1	A	545	ASP	CB-CG-OD2	-5.30	113.53	118.30
1	A	448	PHE	CB-CG-CD1	5.25	124.48	120.80
1	A	363	ARG	CG-CD-NE	5.09	122.49	111.80
1	A	348	MET	CG-SD-CE	5.06	108.30	100.20
1	A	432	PHE	CD1-CG-CD2	5.04	124.85	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	416	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2306	0	2292	34	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	7	0	2	0	0
3	A	414	0	0	20	1
All	All	2727	0	2294	34	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:417:GLU:CD	3:A:703:HOH:O	1.80	1.15
1:A:363:ARG:NH1	3:A:702:HOH:O	1.97	0.94
1:A:363:ARG:CZ	3:A:702:HOH:O	2.21	0.89
1:A:310[B]:ASP:OD1	3:A:701:HOH:O	1.94	0.84
1:A:363:ARG:NH2	3:A:702:HOH:O	2.16	0.78
1:A:533:GLU:O	1:A:537:LYS:HG2	1.89	0.73
1:A:570[A]:LYS:HE3	3:A:1028:HOH:O	1.88	0.71
1:A:534:ASP:HA	1:A:537:LYS:HG3	1.75	0.68
1:A:326:LYS:HE3	1:A:352:TRP:CE2	2.35	0.61
1:A:370:HIS:HD2	1:A:507:ASP:OD2	1.83	0.61
1:A:537:LYS:HD2	3:A:858:HOH:O	2.02	0.59
1:A:570[A]:LYS:HE2	3:A:909:HOH:O	2.02	0.58
1:A:504:HIS:CE1	3:A:815:HOH:O	2.56	0.58
1:A:433[A]:LYS:HG3	3:A:936:HOH:O	2.03	0.57
1:A:533:GLU:O	1:A:537:LYS:CG	2.54	0.56
1:A:433[A]:LYS:HG2	1:A:439:GLN:O	2.07	0.55
1:A:463:ASN:ND2	3:A:710:HOH:O	2.42	0.52
1:A:433[B]:LYS:HG2	1:A:438:GLY:HA2	1.90	0.52
1:A:307:LYS:HG3	3:A:880:HOH:O	2.11	0.51
1:A:350:GLN:OE1	3:A:704:HOH:O	2.20	0.50
1:A:417:GLU:OE2	3:A:703:HOH:O	2.10	0.49
1:A:442[A]:LYS:NZ	3:A:707:HOH:O	2.44	0.49
1:A:433[A]:LYS:CG	3:A:936:HOH:O	2.62	0.48
1:A:528:ARG:CD	3:A:1006:HOH:O	2.62	0.47
1:A:417:GLU:OE1	3:A:703:HOH:O	2.08	0.47
1:A:505:ASP:HB3	1:A:521:LYS:HD2	1.99	0.45
1:A:404:LYS:HZ3	1:A:477:GLU:CD	2.20	0.45
1:A:528:ARG:HD3	3:A:1006:HOH:O	2.16	0.45
1:A:547:ILE:HG23	1:A:552:ASP:HB3	1.99	0.44
1:A:490:PRO:HB2	1:A:492[A]:ILE:HG23	2.01	0.42
1:A:537:LYS:HE3	1:A:537:LYS:HB3	1.68	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:325[B]:LEU:HD12	3:A:703:HOH:O	2.20	0.41
1:A:442[A]:LYS:NZ	1:A:450:GLU:OE2	2.51	0.40
1:A:555:MET:HB2	1:A:555:MET:HE2	1.79	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:803:HOH:O	3:A:1013:HOH:O[2_554]	1.86	0.34

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	293/297 (99%)	290 (99%)	3 (1%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	249/252 (99%)	240 (96%)	9 (4%)	35 12

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	325[A]	LEU
1	A	325[B]	LEU
1	A	397	LEU
1	A	419	ASP
1	A	492[A]	ILE
1	A	492[B]	ILE
1	A	520	LEU
1	A	528	ARG
1	A	537	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	350	GLN
1	A	355	ASN
1	A	370	HIS
1	A	439	GLN
1	A	463	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	MLI	A	601	-	6,6,6	1.19	0	7,7,7	1.42	1 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MLI	A	601	-	-	0/4/4/4	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	MLI	O9-C3-C1	2.21	121.60	114.54

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	284/297 (95%)	-0.31	1 (0%) 92 92	9, 15, 26, 35	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	440	TYR	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	MLI	A	601	7/7	0.91	0.19	31,34,40,40	0

6.5 Other polymers [i](#)

There are no such residues in this entry.