



Full wwPDB X-ray Structure Validation Report ⓘ

May 15, 2020 – 02:40 am BST

PDB ID : 2BWJ
Title : Structure of adenylate kinase 5
Authors : Bunkoczi, G.; Filippakopoulos, P.; Fedorov, O.; Jansson, A.; Longman, E.; Ugochukwu, E.; Knapp, S.; von Delft, F.; Arrowsmith, C.; Edwards, A.; Sundstrom, M.; Weigelt, J.
Deposited on : 2005-07-14
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	199	91% 7% ..
1	B	199	90% 6% ..
1	C	199	91% 6% .
1	D	199	90% 6% .
1	E	199	85% 10% . 5%
1	F	199	91% 6% ..

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 9288 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called ADENYLATE KINASE 5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	196	1495	950	246	289	10	0	0	0
1	B	195	1490	942	242	296	10	0	2	0
1	C	193	1465	926	239	289	11	0	1	0
1	D	192	1366	857	228	271	10	0	0	0
1	E	190	1429	909	229	281	10	0	2	0
1	F	194	1482	930	247	293	12	0	1	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	135	MET	ARG	engineered mutation	UNP Q9Y6K8
B	135	MET	ARG	engineered mutation	UNP Q9Y6K8
C	135	MET	ARG	engineered mutation	UNP Q9Y6K8
D	135	MET	ARG	engineered mutation	UNP Q9Y6K8
E	135	MET	ARG	engineered mutation	UNP Q9Y6K8
F	135	MET	ARG	engineered mutation	UNP Q9Y6K8

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		
2	D	1	Total	O	S	0	0
			5	4	1		
2	E	1	Total	O	S	0	0
			5	4	1		
2	F	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Cl	0	0
			1	1		
3	A	1	Total	Cl	0	0
			1	1		
3	F	2	Total	Cl	0	0
			2	2		
3	E	1	Total	Cl	0	0
			1	1		

- Molecule 4 is ADENOSINE MONOPHOSPHATE (three-letter code: AMP) (formula: C₁₀H₁₄N₅O₇P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	B	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
4	C	1	Total	C	N	O		0	0
			19	10	5	4			
4	F	1	Total	C	N	O		0	0
			19	10	5	4			

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	125	Total	O	0	0
			125	125		
5	B	102	Total	O	0	0
			102	102		
5	C	86	Total	O	0	0
			86	86		
5	D	8	Total	O	0	0
			8	8		
5	E	40	Total	O	0	0
			40	40		
5	F	99	Total	O	0	0
			99	99		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: ADENYLATE KINASE 5

Chain A:  91% 7% ..



- Molecule 1: ADENYLATE KINASE 5

Chain B:  90% 6% ..



- Molecule 1: ADENYLATE KINASE 5

Chain C:  91% 6% .



- Molecule 1: ADENYLATE KINASE 5

Chain D:  90% 6% .



- Molecule 1: ADENYLATE KINASE 5

Chain E:  85% 10% • 5%



- Molecule 1: ADENYLATE KINASE 5

Chain F:  91% 6% ..



4 Data and refinement statistics

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, α , β , γ	180.38Å 180.38Å 121.70Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 2.30 43.32 – 2.30	Depositor EDS
% Data completeness (in resolution range)	88.9 (50.00-2.30) 88.9 (43.32-2.30)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.10 (at 2.29Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.172 , 0.213 0.181 , 0.219	Depositor DCC
R_{free} test set	4454 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	44.6	Xtrriage
Anisotropy	0.112	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 56.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.031 for h,-h-k,-l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	9288	wwPDB-VP
Average B, all atoms (Å ²)	61.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.22% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: AMP, SO4, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.67	1/1516 (0.1%)	0.75	2/2042 (0.1%)
1	B	0.65	1/1523 (0.1%)	0.79	3/2056 (0.1%)
1	C	0.58	0/1488	0.65	0/2010
1	D	0.43	0/1383	0.54	0/1877
1	E	0.54	1/1455 (0.1%)	0.66	0/1969
1	F	0.60	0/1505	0.71	1/2028 (0.0%)
All	All	0.58	3/8870 (0.0%)	0.69	6/11982 (0.1%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	120	CYS	CB-SG	-6.36	1.71	1.82
1	B	123	CYS	CB-SG	-5.64	1.72	1.81
1	A	120	CYS	CB-SG	-5.50	1.72	1.81

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	100	ARG	NE-CZ-NH1	7.89	124.24	120.30
1	A	158	ARG	NE-CZ-NH2	-6.39	117.10	120.30
1	B	100	ARG	NE-CZ-NH2	-5.92	117.34	120.30
1	B	66	ARG	NE-CZ-NH1	5.28	122.94	120.30
1	F	100	ARG	NE-CZ-NH2	5.20	122.90	120.30
1	A	158	ARG	NE-CZ-NH1	5.07	122.84	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1495	0	1488	5	0
1	B	1490	0	1458	9	0
1	C	1465	0	1442	7	0
1	D	1366	0	1276	5	0
1	E	1429	0	1393	7	0
1	F	1482	0	1468	8	0
2	A	10	0	0	0	0
2	B	5	0	0	0	0
2	C	5	0	0	0	0
2	D	5	0	0	0	0
2	E	5	0	0	0	0
2	F	5	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	E	1	0	0	0	0
3	F	2	0	0	0	0
4	B	23	0	12	2	0
4	C	19	0	12	0	0
4	F	19	0	12	1	0
5	A	125	0	0	0	0
5	B	102	0	0	2	0
5	C	86	0	0	0	0
5	D	8	0	0	0	0
5	E	40	0	0	0	0
5	F	99	0	0	2	0
All	All	9288	0	8561	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:172:GLN:HG2	1:F:196:ILE:HD11	1.72	0.70
1:A:64:MET:HE3	5:B:2039:HOH:O	1.97	0.64

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:117:LEU:HD13	1:C:196:ILE:HD12	1.84	0.59
1:A:95:ILE:HG22	1:A:98:TYR:HB3	1.84	0.58
1:C:145:THR:HG23	1:F:55:GLU:HG2	1.86	0.58
1:C:117:LEU:HD13	1:C:196:ILE:CD1	2.34	0.57
1:D:65:GLU:O	1:E:42:THR:OG1	2.19	0.55
4:F:1202:AMP:N3	5:F:2098:HOH:O	2.33	0.55
1:E:80:LYS:O	1:E:84:VAL:HG13	2.07	0.54
1:E:11:CYS:O	1:E:116:GLN:NE2	2.42	0.53
1:B:176:ILE:CD1	1:B:189:LEU:HD23	2.41	0.50
1:F:109:GLY:HA2	1:F:113:GLY:O	2.12	0.49
1:E:119:ILE:HG23	1:E:176:ILE:HD13	1.95	0.49
1:A:71:PRO:HB2	1:A:74:ILE:HD13	1.96	0.47
1:C:145:THR:HG23	1:F:55:GLU:CG	2.45	0.46
1:F:104:GLN:HG3	5:F:2057:HOH:O	2.15	0.46
1:B:166:TYR:CZ	1:B:170:LYS:HD2	2.50	0.46
1:B:176:ILE:HD13	1:B:189:LEU:HD23	1.98	0.45
1:A:119:ILE:HG23	1:A:176:ILE:HD13	1.99	0.45
1:E:30:LYS:HB3	1:E:186:PHE:CE1	2.52	0.45
1:C:31:LEU:HD22	1:C:193:ILE:CD1	2.47	0.44
1:E:119:ILE:HG21	1:E:189:LEU:CD2	2.47	0.44
1:A:100:ARG:HG2	1:A:104:GLN:OE1	2.18	0.44
1:D:28:CYS:HB3	1:D:39:HIS:CE1	2.53	0.43
4:B:1201:AMP:H2'	4:B:1201:AMP:O2P	2.17	0.43
1:E:131:ARG:O	1:E:135:MET:HG2	2.18	0.43
1:C:95:ILE:HG22	1:C:98:TYR:HB3	1.99	0.43
1:C:181:THR:HG22	1:C:184:ASP:OD2	2.19	0.42
4:B:1201:AMP:O5'	4:B:1201:AMP:C8	2.73	0.42
1:F:119:ILE:HG21	1:F:189:LEU:CD2	2.49	0.42
1:B:119:ILE:HG23	1:B:176:ILE:HD12	2.02	0.41
1:D:80:LYS:O	1:D:84:VAL:HG23	2.21	0.41
1:B:9:ARG:NH1	1:B:109:GLY:O	2.53	0.41
1:F:71:PRO:HB2	1:F:74:ILE:HD13	2.02	0.41
1:D:196:ILE:N	1:D:196:ILE:HD12	2.36	0.41
1:D:181:THR:O	1:D:185:VAL:HG23	2.21	0.41
1:B:66:ARG:HD3	5:B:2042:HOH:O	2.21	0.41
1:B:90:THR:HG22	1:B:92:GLY:H	1.84	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	194/199 (98%)	192 (99%)	2 (1%)	0	100	100
1	B	195/199 (98%)	193 (99%)	2 (1%)	0	100	100
1	C	192/199 (96%)	189 (98%)	3 (2%)	0	100	100
1	D	190/199 (96%)	184 (97%)	6 (3%)	0	100	100
1	E	190/199 (96%)	188 (99%)	2 (1%)	0	100	100
1	F	193/199 (97%)	191 (99%)	2 (1%)	0	100	100
All	All	1154/1194 (97%)	1137 (98%)	17 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	156/171 (91%)	152 (97%)	4 (3%)	46	63
1	B	157/171 (92%)	150 (96%)	7 (4%)	27	39
1	C	155/171 (91%)	153 (99%)	2 (1%)	69	82
1	D	132/171 (77%)	128 (97%)	4 (3%)	41	57
1	E	147/171 (86%)	137 (93%)	10 (7%)	16	21
1	F	158/171 (92%)	156 (99%)	2 (1%)	69	82
All	All	905/1026 (88%)	876 (97%)	29 (3%)	40	54

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	122	ASP
1	A	144	ASP
1	A	174	HIS
1	A	176	ILE
1	B	9	ARG
1	B	11	CYS
1	B	90	THR
1	B	100	ARG
1	B	101[A]	GLU
1	B	101[B]	GLU
1	B	188	GLN
1	C	47	ARG
1	C	144	ASP
1	D	13	ILE
1	D	26	THR
1	D	69	LEU
1	D	137	ARG
1	E	8	LEU
1	E	11	CYS
1	E	26	THR
1	E	86	SER
1	E	121	MET
1	E	126	ASP
1	E	135	MET
1	E	137	ARG
1	E	142	VAL
1	E	187	LEU
1	F	100	ARG
1	F	122	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 15 ligands modelled in this entry, 5 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	AMP	F	1202	-	18,21,25	1.15	2 (11%)	18,31,38	1.66	4 (22%)
2	SO4	A	1200	-	4,4,4	0.12	0	6,6,6	0.27	0
2	SO4	C	1198	-	4,4,4	0.16	0	6,6,6	0.44	0
2	SO4	E	1198	-	4,4,4	0.17	0	6,6,6	0.63	0
4	AMP	C	1199	-	18,21,25	1.14	3 (16%)	18,31,38	1.76	5 (27%)
4	AMP	B	1201	-	22,25,25	1.17	2 (9%)	25,38,38	1.90	8 (32%)
2	SO4	D	1197	-	4,4,4	0.13	0	6,6,6	0.53	0
2	SO4	F	1199	-	4,4,4	0.12	0	6,6,6	0.66	0
2	SO4	A	1199	-	4,4,4	0.33	0	6,6,6	0.93	0
2	SO4	B	1199	-	4,4,4	0.22	0	6,6,6	1.27	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	AMP	F	1202	-	-	2/2/22/26	0/3/3/3
4	AMP	B	1201	-	-	4/6/26/26	0/3/3/3
4	AMP	C	1199	-	-	0/2/22/26	0/3/3/3

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	1201	AMP	O4'-C1'	2.90	1.45	1.41

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	F	1202	AMP	O4'-C1'	2.56	1.44	1.41
4	B	1201	AMP	C5-C4	2.54	1.47	1.40
4	C	1199	AMP	C5-C4	2.52	1.47	1.40
4	F	1202	AMP	C5-C4	2.44	1.47	1.40
4	C	1199	AMP	O4'-C1'	2.31	1.44	1.41
4	C	1199	AMP	C2-N3	2.03	1.35	1.32

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	1201	AMP	N3-C2-N1	-4.53	121.60	128.68
4	F	1202	AMP	N3-C2-N1	-4.21	122.09	128.68
4	C	1199	AMP	N3-C2-N1	-3.98	122.45	128.68
4	B	1201	AMP	C3'-C2'-C1'	3.68	106.51	100.98
4	B	1201	AMP	C1'-N9-C4	-3.63	120.26	126.64
4	F	1202	AMP	C3'-C2'-C1'	3.18	105.76	100.98
4	C	1199	AMP	C3'-C2'-C1'	3.15	105.72	100.98
4	C	1199	AMP	C1'-N9-C4	-3.06	121.27	126.64
4	C	1199	AMP	C4-C5-N7	-2.82	106.46	109.40
4	B	1201	AMP	O4'-C1'-C2'	-2.77	102.88	106.93
4	F	1202	AMP	C4-C5-N7	-2.77	106.52	109.40
4	B	1201	AMP	C4-C5-N7	-2.46	106.84	109.40
4	B	1201	AMP	C2-N1-C6	2.29	122.68	118.75
4	B	1201	AMP	P-O5'-C5'	2.23	124.43	118.30
4	C	1199	AMP	C2'-C3'-C4'	2.21	106.93	102.64
4	F	1202	AMP	C2-N1-C6	2.09	122.32	118.75
4	B	1201	AMP	O3'-C3'-C4'	-2.02	105.20	111.05

There are no chirality outliers.

All (6) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	B	1201	AMP	C5'-O5'-P-O1P
4	B	1201	AMP	C5'-O5'-P-O2P
4	B	1201	AMP	C5'-O5'-P-O3P
4	F	1202	AMP	O4'-C4'-C5'-O5'
4	B	1201	AMP	C4'-C5'-O5'-P
4	F	1202	AMP	C3'-C4'-C5'-O5'


There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	F	1202	AMP	1	0
4	B	1201	AMP	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.




6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around AMP C 1199:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around AMP F 1202:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [\(i\)](#)

Unable to reproduce the depositors R factor - this section is therefore empty.