

wwPDB X-ray Structure Validation Summary Report (i)

Dec 20, 2023 – 12:41 pm GMT

PDB ID : 8BZ6

Title : Crystal structure of the L. monocytogenes RmlT in complex with UDP-glucose

Authors: Cereija, T.B.; Morais-Cabral, J.H.

Deposited on : 2022-12-14

Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.4, CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.36

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},{\rm resolution\ range}(\mathring{\rm A})) \end{array}$
R_{free}	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	624	91%	6%	
1			13%	076	
1	В	624	86%	11%	•
1	C	624	92%	5%	-
1	D	624	88%	9%	-
1	Е	624	88%	9%	-



Continued from previous page...

Mol	Chain	Length	Quality of chain		
4	Б	00.4	9%		_
1	F'	624	88%	9%	•



2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 30490 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Glycosyltransferase.


Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	A	606	Total	С	N	О	S	0	1	0
1	Λ	000	4915	3132	817	954	12	U	1	
1	В	605	Total	С	N	О	S	0	0	0
1	Ъ	005	4897	3121	815	949	12	U	0	
1	С	606	Total	С	N	О	S	0	1	0
1		000	4914	3131	817	954	12	U	1	
1	D	606	Total	С	N	О	S	0	2	0
1	D	000	4922	3135	819	956	12	U		
1	Е	606	Total	С	N	О	S	0	0	0
1	l L	000	4906	3127	816	951	12	U	0	
1	F	606	Total	С	N	О	S	0	0	0
1	I'	000	4906	3127	816	951	12	U	U	

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	expression tag	UNP A0A401AAP7
В	0	GLY	-	expression tag	UNP A0A401AAP7
С	0	GLY	-	expression tag	UNP A0A401AAP7
D	0	GLY	-	expression tag	UNP A0A401AAP7
Е	0	GLY	-	expression tag	UNP A0A401AAP7
F	0	GLY	-	expression tag	UNP A0A401AAP7

• Molecule 2 is URIDINE-5'-DIPHOSPHATE-GLUCOSE (three-letter code: UPG) (formula: $C_{15}H_{24}N_2O_{17}P_2$) (labeled as "Ligand of Interest" by depositor).






Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf
2	Λ	1	Total	С	N	О	Р	0	0
	A	1	36	15	2	17	2	0	U
2	В	1	Total	С	N	О	Р	0	0
2	Б	1	36	15	2	17	2	U	0
2	C	1	Total	С	N	О	Р	0	0
2		1	36	15	2	17	2	0	
2	D	1	Total	С	N	О	Р	0	0
	D	1	36	15	2	17	2	U	U
2	E	1	Total	С	N	Ο	Р	0	0
	Ľ	1	36	15	2	17	2	U	U
2	F	1	Total	С	N	О	Р	0	0
	I.	1	36	15	2	17	2	U	

 \bullet Molecule 3 is ACETATE ION (three-letter code: ACT) (formula: $\mathrm{C_2H_3O_2}).$





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	В	1	Total C O 4 2 2	0	0
3	В	1	Total C O 4 2 2	0	0
3	В	1	Total C O 4 2 2	0	0
3	В	1	Total C O 4 2 2	0	0
3	В	1	Total C O 4 2 2	0	0
3	С	1	Total C O 4 2 2	0	0
3	С	1	Total C O 4 2 2	0	0
3	С	1	Total C O 4 2 2	0	0



 $Continued\ from\ previous\ page...$

Mol	Chain	$oxed{ \mathbf{Residues} }$	Atoms	ZeroOcc	AltConf
3	С	1	Total C O 4 2 2	0	0
3	С	1	Total C O 4 2 2	0	0
3	С	1	Total C O 4 2 2	0	0
3	D	1	Total C O 4 2 2	0	0
3	D	1	Total C O 4 2 2	0	0
3	D	1	Total C O 4 2 2	0	0
3	D	1	Total C O 4 2 2	0	0
3	D	1	Total C O 4 2 2	0	0
3	D	1	Total C O 4 2 2	0	0
3	E	1	Total C O 4 2 2	0	0
3	Е	1	Total C O 4 2 2	0	0
3	Е	1	Total C O 4 2 2	0	0
3	Е	1	Total C O 4 2 2	0	0
3	Е	1	Total C O 4 2 2	0	0
3	Е	1	Total C O 4 2 2	0	0
3	F	1	Total C O 4 2 2	0	0
3	F	1	Total C O 4 2 2	0	0
3	F	1	Total C O 4 2 2	0	0
3	F	1	Total C O 4 2 2	0	0

• Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Mg 1 1	0	0
4	В	1	Total Mg 1 1	0	0
4	С	1	Total Mg 1 1	0	0
4	D	1	Total Mg 1 1	0	0
4	Е	1	Total Mg 1 1	0	0
4	F	1	Total Mg 1 1	0	0

• Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	В	1	Total Cl 1 1	0	0
5	F	1	Total Cl 1 1	0	0

• Molecule 6 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	156	Total O 156 156	0	0
6	В	79	Total O 79 79	0	0
6	С	115	Total O 115 115	0	0
6	D	100	Total O 100 100	0	0
6	E	133	Total O 133 133	0	0
6	F	91	Total O 91 91	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Glycosyltransferase








1495 K501 K501 1504 L558 K566 K566 K581 K581 L583 K588 K589 K588 L606 L606 L606 L606 L606


• Molecule 1: Glycosyltransferase

Chain E: 88% 9% .



• Molecule 1: Glycosyltransferase

Chain F: 88% 9% •





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	85.52Å 291.06Å 89.82Å	Donositon
a, b, c, α , β , γ	90.00° 100.08° 90.00°	Depositor
Resolution (Å)	44.22 - 2.40	Depositor
Resolution (A)	49.34 - 2.40	EDS
% Data completeness	99.1 (44.22-2.40)	Depositor
(in resolution range)	99.1 (49.34-2.40)	EDS
R_{merge}	0.10	Depositor
R_{sym}	0.10	Depositor
$< I/\sigma(I) > 1$	1.55 (at 2.39Å)	Xtriage
Refinement program	PHENIX 1.19.2_4158	Depositor
D D	0.195 , 0.235	Depositor
R, R_{free}	0.192 , 0.232	DCC
R_{free} test set	8374 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	56.0	Xtriage
Anisotropy	0.200	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.32, 42.6	EDS
L-test for twinning ²	$< L >=0.50, < L^2>=0.34$	Xtriage
Estimated twinning fraction	0.013 for l,-k,h	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	30490	wwPDB-VP
Average B, all atoms (Å ²)	69.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 2.96% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: UPG, CL, MG, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
IVIOI		RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.28	0/5016	0.49	0/6774	
1	В	0.27	0/4998	0.49	0/6751	
1	С	0.27	0/5015	0.49	0/6773	
1	D	0.28	0/5023	0.49	0/6784	
1	Е	0.28	0/5007	0.50	0/6762	
1	F	0.27	0/5007	0.48	0/6762	
All	All	0.28	0/30066	0.49	0/40606	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4915	0	4849	22	0
1	В	4897	0	4833	39	0
1	С	4914	0	4847	21	0
1	D	4922	0	4852	37	0
1	Е	4906	0	4844	33	0
1	F	4906	0	4844	33	0
2	A	36	0	22	1	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	В	36	0	22	0	0
2	С	36	0	22	0	0
2	D	36	0	22	0	0
2	Ε	36	0	22	0	0
2	F	36	0	22	1	0
3	A	24	0	18	0	0
3	В	20	0	15	1	0
3	С	24	0	18	1	0
3	D	24	0	18	1	0
3	Ε	24	0	18	1	0
3	F	16	0	12	1	0
4	A	1	0	0	0	0
4	В	1	0	0	0	0
4	С	1	0	0	0	0
4	D	1	0	0	0	0
4	Е	1	0	0	0	0
4	F	1	0	0	0	0
5	В	1	0	0	0	0
5	F	1	0	0	0	0
6	A	156	0	0	1	0
6	В	79	0	0	3	0
6	С	115	0	0	1	0
6	D	100	0	0	2	0
6	Е	133	0	0	2	0
6	F	91	0	0	3	0
All	All	30490	0	29300	184	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 184 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$\begin{array}{c} \text{Clash} \\ \text{overlap } (\text{\AA}) \end{array}$
1:B:582:LEU:HB2	1:B:610:ILE:HD13	1.69	0.74
1:D:43:ASP:OD2	1:D:76:ARG:NH1	2.20	0.72
1:F:521:VAL:HG22	1:F:522:ARG:HG3	1.71	0.71
1:F:383:MET:HG3	1:F:406:LEU:HD11	1.72	0.70
1:C:233:SER:HA	1:C:236:ARG:HG3	1.72	0.69

There are no symmetry-related clashes.



5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured Allowed		Outliers Per		ercentiles	
1	A	605/624~(97%)	582 (96%)	23 (4%)	0	100	100	
1	В	603/624 (97%)	569 (94%)	34 (6%)	0	100	100	
1	С	605/624 (97%)	587 (97%)	18 (3%)	0	100	100	
1	D	606/624 (97%)	588 (97%)	18 (3%)	0	100	100	
1	E	604/624 (97%)	586 (97%)	18 (3%)	0	100	100	
1	F	604/624 (97%)	582 (96%)	22 (4%)	0	100	100	
All	All	3627/3744 (97%)	3494 (96%)	133 (4%)	0	100	100	

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric Outliers		Percentiles		
1	A	538/553 (97%)	535 (99%)	3 (1%)	86	94	
1	В	536/553~(97%)	533 (99%)	3 (1%)	86	94	
1	С	538/553 (97%)	536 (100%)	2 (0%)	91	96	
1	D	539/553~(98%)	534 (99%)	5 (1%)	78	90	
1	E	537/553 (97%)	536 (100%)	1 (0%)	93	97	
1	F	537/553 (97%)	535 (100%)	2 (0%)	91	96	
All	All	3225/3318 (97%)	3209 (100%)	16 (0%)	88	95	

5 of 16 residues with a non-rotameric sidechain are listed below:



Mol	Chain	Res	Type
1	F	111	HIS
1	Е	111	HIS
1	D	35	LEU
1	D	599	LYS
1	С	572	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 47 ligands modelled in this entry, 8 are monoatomic - leaving 39 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Во	Bond lengths			Bond angles		
MIOI	туре				Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
3	ACT	Е	708	-	3,3,3	1.03	0	3,3,3	0.83	0	
3	ACT	A	702	-	3,3,3	1.24	0	3,3,3	1.46	0	
3	ACT	F	706	-	3,3,3	1.06	0	3,3,3	0.67	0	
3	ACT	F	702	-	3,3,3	1.11	0	3,3,3	1.41	0	
2	UPG	A	701	4	35,38,38	0.37	0	53,58,58	0.48	0	
3	ACT	D	706	-	3,3,3	1.09	0	3,3,3	0.80	0	
2	UPG	F	701	4	35,38,38	0.41	0	53,58,58	0.61	0	



Mal	Trino	Chain	Dag	T inle	Во	ond leng	ths	В	ond ang	gles
Mol	Type	Chain	Res	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	UPG	Е	701	4	35,38,38	0.37	0	53,58,58	0.46	0
2	UPG	В	701	4	35,38,38	0.38	0	53,58,58	0.47	0
3	ACT	С	707	-	3,3,3	1.14	0	3,3,3	0.85	0
2	UPG	D	701	4	35,38,38	0.37	0	53,58,58	0.55	0
3	ACT	С	702	-	3,3,3	1.28	0	3,3,3	1.45	0
3	ACT	С	708	-	3,3,3	1.23	1 (33%)	3,3,3	0.81	0
3	ACT	D	707	-	3,3,3	1.12	0	3,3,3	0.80	0
3	ACT	D	708	-	3,3,3	0.96	0	3,3,3	0.93	0
2	UPG	С	701	4	35,38,38	0.38	0	53,58,58	0.50	0
3	ACT	A	705	-	3,3,3	1.09	0	3,3,3	0.79	0
3	ACT	В	706	-	3,3,3	1.03	0	3,3,3	0.81	0
3	ACT	Е	707	-	3,3,3	1.12	0	3,3,3	0.81	0
3	ACT	С	705	-	3,3,3	1.03	0	3,3,3	0.91	0
3	ACT	F	704	-	3,3,3	1.07	0	3,3,3	0.85	0
3	ACT	D	705	-	3,3,3	0.99	0	3,3,3	0.94	0
3	ACT	С	704	-	3,3,3	1.06	0	3,3,3	0.80	0
3	ACT	С	706	-	3,3,3	1.07	0	3,3,3	0.86	0
3	ACT	D	704	-	3,3,3	0.98	0	3,3,3	0.71	0
3	ACT	Е	704	-	3,3,3	1.10	0	3,3,3	0.74	0
3	ACT	A	704	_	3,3,3	1.15	0	3,3,3	0.90	0
3	ACT	E	706	_	3,3,3	1.04	0	3,3,3	0.67	0
3	ACT	A	706	-	3,3,3	1.04	0	3,3,3	0.73	0
3	ACT	В	705	_	3,3,3	1.16	0	3,3,3	0.86	0
3	ACT	D	702	_	3,3,3	1.18	0	3,3,3	1.49	0
3	ACT	Е	705	-	3,3,3	0.98	0	3,3,3	0.77	0
3	ACT	F	705	_	3,3,3	0.96	0	3,3,3	0.82	0
3	ACT	В	707	-	3,3,3	0.98	0	3,3,3	0.81	0
3	ACT	Е	702	-	3,3,3	1.09	0	3,3,3	1.50	0
3	ACT	В	703	-	3,3,3	1.00	0	3,3,3	0.77	0
3	ACT	A	708	-	3,3,3	1.14	0	3,3,3	0.74	0
3	ACT	A	707	-	3,3,3	1.14	0	3,3,3	0.66	0
3	ACT	В	704	-	3,3,3	1.01	0	3,3,3	0.83	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	UPG	D	701	4	-	6/23/59/59	0/3/3/3
2	UPG	F	701	4	-	4/23/59/59	0/3/3/3
2	UPG	A	701	4	-	4/23/59/59	0/3/3/3



Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	UPG	С	701	4	-	4/23/59/59	0/3/3/3
2	UPG	E	701	4	-	6/23/59/59	0/3/3/3
2	UPG	В	701	4	-	4/23/59/59	0/3/3/3

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\operatorname{\AA})$	$\operatorname{Ideal}({ ext{A}})$
3	С	708	ACT	OXT-C	-2.04	1.21	1.30

There are no bond angle outliers.

There are no chirality outliers.

5 of 28 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	В	701	UPG	C5C-O5C-PA-O1A
2	D	701	UPG	C5C-O5C-PA-O1A
2	D	701	UPG	C5C-O5C-PA-O2A
2	F	701	UPG	O5'-C1'-O3B-PB
2	Е	701	UPG	C1'-O3B-PB-O3A

There are no ring outliers.


7 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	701	UPG	1	0
2	F	701	UPG	1	0
3	D	708	ACT	1	0
3	Е	707	ACT	1	0
3	С	704	ACT	1	0
3	F	705	ACT	1	0
3	В	707	ACT	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the




average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.






















5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\#\mathrm{RSRZ}{>}2$	$OWAB(Å^2)$	Q < 0.9
1	A	606/624 (97%)	0.24	10 (1%) 70 68	41, 61, 103, 161	0
1	В	605/624 (96%)	0.71	78 (12%) 3 3	43, 72, 148, 196	0
1	С	606/624 (97%)	0.31	13 (2%) 63 61	45, 63, 93, 189	0
1	D	606/624 (97%)	0.23	11 (1%) 68 66	46, 61, 89, 148	0
1	E	$606/624 \ (97\%)$	0.15	7 (1%) 79 77	40, 61, 91, 147	0
1	F	606/624 (97%)	0.62	54 (8%) 9 9	43, 75, 129, 190	0
All	All	3635/3744 (97%)	0.38	173 (4%) 30 29	40, 64, 114, 196	0

The worst 5 of 173 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	191	GLY	15.1
1	В	604	TYR	9.5
1	В	595	LEU	9.3
1	F	573	ALA	7.9
1	В	572	ASP	7.2

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.



6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
4	MG	В	702	1/1	0.87	0.08	66,66,66,66	0
3	ACT	D	704	4/4	0.91	0.14	64,69,69,71	0
3	ACT	В	704	4/4	0.91	0.16	72,75,76,81	0
4	MG	F	703	1/1	0.91	0.08	74,74,74,74	0
2	UPG	F	701	36/36	0.92	0.16	69,82,92,95	0
3	ACT	F	705	4/4	0.92	0.27	74,75,77,79	0
3	ACT	В	703	4/4	0.93	0.12	81,82,83,87	0
4	MG	A	703	1/1	0.93	0.09	65,65,65,65	0
2	UPG	В	701	36/36	0.93	0.18	65,72,77,79	0
3	ACT	A	705	4/4	0.93	0.19	56,60,64,65	0
5	CL	В	708	1/1	0.93	0.08	70,70,70,70	0
3	ACT	С	706	4/4	0.94	0.19	49,54,56,59	0
2	UPG	A	701	36/36	0.94	0.18	70,78,86,87	0
2	UPG	Е	701	36/36	0.94	0.20	56,65,72,74	0
3	ACT	F	706	4/4	0.94	0.11	63,64,65,66	0
3	ACT	Е	705	4/4	0.95	0.17	57,59,59,63	0
2	UPG	D	701	36/36	0.95	0.16	56,62,69,73	0
3	ACT	С	708	4/4	0.95	0.23	56,56,58,65	0
3	ACT	D	702	4/4	0.95	0.23	56,59,60,69	0
3	ACT	В	707	4/4	0.95	0.20	67,71,73,74	0
4	MG	Е	703	1/1	0.95	0.10	56,56,56,56	0
3	ACT	D	706	4/4	0.95	0.20	54,54,58,59	0
3	ACT	Е	704	4/4	0.95	0.18	60,60,61,62	0
3	ACT	F	704	4/4	0.96	0.23	58,59,61,67	0
4	MG	D	703	1/1	0.96	0.07	65,65,65,65	0
2	UPG	С	701	36/36	0.96	0.15	59,68,77,81	0
3	ACT	С	705	4/4	0.96	0.21	57,59,59,67	0
3	ACT	A	707	4/4	0.96	0.18	52,54,54,57	0
3	ACT	С	707	4/4	0.97	0.26	51,51,53,56	0
3	ACT	D	708	4/4	0.97	0.19	62,64,65,69	0
3	ACT	A	706	4/4	0.97	0.15	54,55,58,65	0
3	ACT	A	708	4/4	0.97	0.24	62,63,63,65	0
3	ACT	Е	706	4/4	0.97	0.18	56,56,60,68	0
3	ACT	В	706	4/4	0.97	0.20	62,65,65,69	0
3	ACT	D	705	4/4	0.97	0.34	60,63,65,70	0
3	ACT	Е	702	4/4	0.98	0.29	43,43,46,52	0
3	ACT	В	705	4/4	0.98	0.29	52,52,57,59	0



Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}(\mathring{\mathbf{A}}^2)$	Q<0.9
3	ACT	С	702	4/4	0.98	0.18	56,64,64,68	0
3	ACT	С	704	4/4	0.98	0.19	62,62,65,65	0
4	MG	С	703	1/1	0.98	0.16	58,58,58,58	0
3	ACT	E	707	4/4	0.98	0.25	51,52,56,59	0
3	ACT	Ε	708	4/4	0.98	0.24	58,60,62,66	0
3	ACT	F	702	4/4	0.98	0.30	51,51,54,58	0
3	ACT	A	702	4/4	0.98	0.28	44,49,49,59	0
5	CL	F	707	1/1	0.98	0.12	60,60,60,60	0
3	ACT	D	707	4/4	0.99	0.24	50,51,52,52	0
3	ACT	A	704	4/4	0.99	0.29	54,56,56,60	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around UPG B 701:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around UPG A 701:


 $2 \mathrm{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)













Electron density around UPG D 701: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${ m mF}_o{ m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive) Electron density around UPG C 701: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.

