



Full wwPDB X-ray Structure Validation Report ⓘ

May 13, 2020 – 01:39 am BST

PDB ID : 4C0H
Title : Extended interface between Pcf11p and Clp1p and structural basis for ATP loss in Gly135Arg point mutant
Authors : Dupin, A.F.; Fribourg, S.
Deposited on : 2013-08-02
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtrriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	445	<p>3% 85% 9% 5%</p>
1	B	445	<p>3% 85% 9% . .</p>
2	C	110	<p>3% 20% . . 75%</p>
2	D	110	<p>2% 23% 6% . 70%</p>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 7336 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MRNA CLEAVAGE AND POLYADENYLATION FACTOR CLP1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	422	3391	2180	569	630	12	0	1	0
1	B	427	3421	2198	574	637	12	0	1	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	135	ARG	GLY	engineered mutation	UNP Q08685
B	135	ARG	GLY	engineered mutation	UNP Q08685

- Molecule 2 is a protein called PCF11P.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	C	28	231	141	40	50	0	0	0
2	D	33	268	165	47	56	0	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	8	Total 8 O 8	0	0
3	B	14	Total 14 O 14	0	0
3	C	1	Total 1 O 1	0	0
3	D	2	Total 2 O 2	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: MRNA CLEAVAGE AND POLYADENYLATION FACTOR CLP1



- Molecule 1: MRNA CLEAVAGE AND POLYADENYLATION FACTOR CLP1



- Molecule 2: PCF11P



- Molecule 2: PCF11P



GLY	SER	GLN	ASN	THR	ASP	ALA	ASN	THR	GLY	SER	S462	S463	S464	S465	S466	S467	L468	S469	THR	THR	THR	THR	THR	THR	THR	ARG	TYR	K475	N476	VAL	VAL	R480	VAL	PRO	GLU	GLU	THR	THR	S487	S487	S489	THR	LYS	HIS	LYS	LYS	LYS	ASN	ASN	ASP	TYR	THR	THR	THR	PRO	HIS	HIS	ALA	ALA	ASN	LYS	LYS	ILE	ILE	ASP	LYS	LYS	SER	SER	ALA	LEU	ASN	ASN	ILE	ILE	HIS	HIS	ALA	ALA	ASP	GLU	ASN	ASN	ASP	ASP
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	------	------	------	------	------	------	------	------	-----	-----	-----	-----	-----	-----	-----	-----	-----	------	------	-----	-----	------	-----	-----	-----	-----	-----	-----	------	------	------	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

GLU	GLY	SER	VAL	ASP	ASN	THR	LEU	GLY	SER	ASP	ARG	SER	ASN	GLU	GLU	LEU	GLU	ILE	THR	THR	THR	THR	THR	GLY	LYS	TYR	VAL	VAL	VAL	PRO	GLU	GLU	THR	THR	SER	GLN	GLN	ASP	MET	ALA	ALA	PHE	LYS	LYS
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	89.39Å 95.61Å 182.06Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.80 – 2.70 46.24 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.7 (47.80-2.70) 99.7 (46.24-2.70)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$	-	Xtrriage
Refinement program	BUSTER 2.10.0	Depositor
R, R_{free}	0.191 , 0.223 0.201 , 0.229	Depositor DCC
R_{free} test set	2188 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	55.4	Xtrriage
Anisotropy	0.048	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 52.7	EDS
L-test for twinning ¹	$\langle L \rangle = 0.53$, $\langle L^2 \rangle = 0.37$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7336	wwPDB-VP
Average B, all atoms (Å ²)	59.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.41% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.48	0/3467	0.74	1/4713 (0.0%)
1	B	0.51	0/3498	0.76	0/4758
2	C	0.50	0/235	0.70	0/317
2	D	0.62	0/272	0.75	0/366
All	All	0.50	0/7472	0.75	1/10154 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	310	LEU	N-CA-CB	5.00	120.41	110.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3391	0	3430	21	0
1	B	3421	0	3459	26	0
2	C	231	0	200	10	0
2	D	268	0	244	5	0
3	A	8	0	0	0	0
3	B	14	0	0	0	0
3	C	1	0	0	0	0
3	D	2	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	7336	0	7333	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:343:PRO:O	2:C:467:ASN:HA	1.70	0.91
1:B:168:THR:HG23	2:D:483:TYR:OH	1.75	0.85
1:A:345:ALA:HB3	2:C:466:SER:HA	1.62	0.80
1:A:127:VAL:HG22	1:A:274:VAL:HG11	1.66	0.78
2:C:467:ASN:ND2	2:C:467:ASN:H	1.87	0.72
1:B:348:VAL:HG11	1:B:431:LEU:HD22	1.70	0.72
2:D:466:SER:OG	2:D:467:ASN:N	2.22	0.71
1:B:412:THR:HG21	1:B:422:ARG:HE	1.56	0.70
1:B:135:ARG:HH22	1:B:253:PRO:C	1.99	0.65
1:A:53:VAL:HG13	1:A:75:ILE:HG23	1.82	0.62
2:C:467:ASN:HD22	2:C:467:ASN:H	1.47	0.62
1:B:124:PRO:O	1:B:247:GLY:HA3	2.00	0.60
1:A:348:VAL:HG11	1:A:431:LEU:HD22	1.84	0.59
1:B:137:THR:HG23	1:B:138:SER:H	1.70	0.57
2:C:465:ASN:ND2	2:C:466:SER:OG	2.37	0.56
1:A:371:LEU:HG	1:A:401:ILE:HD12	1.91	0.53
1:B:412:THR:HG21	1:B:422:ARG:NE	2.22	0.53
1:B:134:THR:O	1:B:138:SER:HB3	2.09	0.53
1:B:243:VAL:O	1:B:246:SER:O	2.27	0.52
1:A:357:LYS:HB2	1:A:374:VAL:HG22	1.92	0.52
1:A:55:ILE:HD12	1:A:60:LEU:HD21	1.92	0.51
1:B:55:ILE:HD12	1:B:60:LEU:HD21	1.91	0.51
1:B:128:ILE:CG2	1:B:135:ARG:HB2	2.40	0.51
1:B:371:LEU:HG	1:B:401:ILE:HD12	1.92	0.50
1:A:135:ARG:HB3	1:A:280:LEU:HD22	1.94	0.50
2:D:465:ASN:O	2:D:466:SER:HB3	2.12	0.49
1:B:137:THR:O	1:B:138:SER:C	2.50	0.49
1:B:109:LEU:HD21	1:B:278:LEU:HD21	1.95	0.49
1:A:53:VAL:CG1	1:A:75:ILE:HG23	2.42	0.49
1:B:148:LEU:HD21	1:B:180:ASP:C	2.33	0.49
1:B:168:THR:HG22	1:B:169:VAL:H	1.78	0.48
1:B:357:LYS:HB2	1:B:374:VAL:HG22	1.95	0.48
2:C:467:ASN:N	2:C:467:ASN:ND2	2.60	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:426:PRO:HB2	2:C:480:ARG:HB3	1.96	0.47
1:B:253:PRO:HG2	1:B:258:LEU:HD21	1.95	0.47
1:B:168:THR:HG23	2:D:483:TYR:HH	1.77	0.46
1:B:139:LEU:HD22	1:B:280:LEU:HD11	1.98	0.45
1:A:418:ARG:O	1:A:420:LYS:N	2.50	0.45
1:B:137:THR:CG2	1:B:138:SER:H	2.29	0.45
2:C:467:ASN:N	2:C:467:ASN:HD22	2.15	0.45
1:B:40:ALA:HB3	1:B:41:GLU:OE1	2.19	0.43
2:C:465:ASN:HA	2:C:466:SER:HA	1.73	0.43
1:B:418:ARG:O	1:B:420:LYS:N	2.50	0.43
1:A:135:ARG:HH21	1:A:254:SER:HA	1.83	0.43
1:B:136:LYS:HD3	1:B:164:GLN:HG2	2.00	0.43
1:A:377:THR:HG22	1:A:379:SER:H	1.83	0.42
1:A:135:ARG:HG3	1:A:136:LYS:N	2.35	0.41
1:B:427:VAL:HG22	2:D:480:ARG:O	2.20	0.41
1:A:102:THR:OG1	1:A:310:LEU:HB3	2.20	0.41
1:B:377:THR:HG22	1:B:379:SER:H	1.83	0.41
1:A:107:TYR:CD1	1:A:107:TYR:N	2.88	0.41
1:A:102:THR:HG21	1:A:310:LEU:HA	2.03	0.41
1:A:215:ILE:HG22	1:A:219:LYS:HD3	2.03	0.41
1:A:278:LEU:HB3	1:A:307:ILE:CD1	2.51	0.41
1:A:345:ALA:H	2:C:466:SER:HB3	1.86	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	419/445 (94%)	395 (94%)	21 (5%)	3 (1%)	22 46
1	B	426/445 (96%)	408 (96%)	15 (4%)	3 (1%)	22 46
2	C	24/110 (22%)	20 (83%)	3 (12%)	1 (4%)	3 5

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	D	29/110 (26%)	25 (86%)	2 (7%)	2 (7%)	1	1
All	All	898/1110 (81%)	848 (94%)	41 (5%)	9 (1%)	15	37

All (9) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	309	LYS
1	A	419	ARG
1	B	138	SER
1	B	419	ARG
2	D	464	SER
2	C	465	ASN
2	D	466	SER
1	B	40	ALA
1	A	138	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	383/401 (96%)	369 (96%)	14 (4%)	34	63
1	B	386/401 (96%)	368 (95%)	18 (5%)	26	54
2	C	26/98 (26%)	22 (85%)	4 (15%)	2	7
2	D	30/98 (31%)	28 (93%)	2 (7%)	16	37
All	All	825/998 (83%)	787 (95%)	38 (5%)	27	54

All (38) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	39	LYS
1	A	109	LEU
1	A	112	MET
1	A	118	MET
1	A	127	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	224	GLU
1	A	260	GLU
1	A	283	GLU
1	A	310	LEU
1	A	323	SER
1	A	380	ASN
1	A	394	ARG
1	A	418	ARG
1	A	422	ARG
1	B	41	GLU
1	B	53	VAL
1	B	64	ASP
1	B	109	LEU
1	B	114[A]	GLU
1	B	114[B]	GLU
1	B	136	LYS
1	B	137	THR
1	B	168	THR
1	B	219	LYS
1	B	224	GLU
1	B	232	VAL
1	B	245	ARG
1	B	323	SER
1	B	364	ASN
1	B	380	ASN
1	B	418	ARG
1	B	430	ARG
2	C	464	SER
2	C	465	ASN
2	C	467	ASN
2	C	487	SER
2	D	476	ASN
2	D	487	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
2	C	465	ASN
2	C	467	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	422/445 (94%)	0.05	15 (3%) 42 42	32, 62, 102, 122	0
1	B	427/445 (95%)	-0.07	12 (2%) 53 54	27, 49, 90, 113	0
2	C	28/110 (25%)	0.33	3 (10%) 6 4	32, 49, 102, 111	0
2	D	33/110 (30%)	0.02	2 (6%) 21 20	27, 40, 99, 105	0
All	All	910/1110 (81%)	0.00	32 (3%) 44 44	27, 55, 98, 122	0

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	C	466	SER	6.4
1	A	284	THR	4.1
1	B	137	THR	3.8
1	B	284	THR	3.5
2	D	469	ASN	2.9
1	A	417	LYS	2.9
1	B	418	ARG	2.8
1	A	283	GLU	2.7
1	B	93	ASN	2.6
1	B	135	ARG	2.5
1	A	362	PHE	2.4
2	C	464	SER	2.4
2	D	463	ILE	2.4
1	B	414	VAL	2.4
2	C	465	ASN	2.4
1	B	315	ALA	2.4
1	A	85	TRP	2.4
1	A	136	LYS	2.3
1	A	93	ASN	2.3
1	A	281	CYS	2.3
1	A	415	ASN	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	394	ARG	2.3
1	B	42	GLY	2.3
1	B	350	TYR	2.3
1	A	197	ALA	2.2
1	A	105	TYR	2.2
1	A	137	THR	2.2
1	A	316	VAL	2.1
1	B	286	PRO	2.1
1	B	40	ALA	2.1
1	A	286	PRO	2.0
1	B	376	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.