



Full wwPDB X-ray Structure Validation Report i

Dec 2, 2023 – 03:35 pm GMT

PDB ID : 2C1Y
Title : Structure of PDI-related Chaperone, Wind mutant-Y55K
Authors : Sevvana, M.; Ma, Q.; Barnewitz, K.; Guo, C.; Soling, H.-D.; Ferrari, D.M.; Sheldrick, G.M.
Deposited on : 2005-09-22
Resolution : 2.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The types of validation reports are described at
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1377 (2.26-2.26)
Clashscore	141614	1487 (2.26-2.26)
Ramachandran outliers	138981	1449 (2.26-2.26)
Sidechain outliers	138945	1450 (2.26-2.26)
RSRZ outliers	127900	1356 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



2 Entry composition [\(i\)](#)

There are 2 unique types of molecules in this entry. The entry contains 2704 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called WINDBEUTEL PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	231	Total	C	N	O	S	0	0	0
			1720	1108	282	326	4			

1	B	128	Total	C	N	O	S	0	0	0
			929	596	153	178	2			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	55	LYS	TYR	engineered mutation	UNP O44342
B	55	LYS	TYR	engineered mutation	UNP O44342


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	35	Total O 35 35	0	0
2	B	20	Total O 20 20	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: WINDBEUTEL PROTEIN



- Molecule 1: WINDBEUTEL PROTEIN



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	35.42Å 119.09Å 64.03Å 90.00° 100.53° 90.00°	Depositor
Resolution (Å)	20.00 – 2.25 59.55 – 2.25	Depositor EDS
% Data completeness (in resolution range)	99.9 (20.00-2.25) 97.7 (59.55-2.25)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ ¹	3.01 (at 2.25Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R , R_{free}	0.223 , 0.292 0.226 , 0.294	Depositor DCC
R_{free} test set	1220 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	45.6	Xtriage
Anisotropy	0.357	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 42.1	EDS
L-test for twinning ²	$< L > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2704	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.20% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [\(i\)](#)

5.1 Standard geometry [\(i\)](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.52	17/1751 (1.0%)	1.37	19/2377 (0.8%)
1	B	1.58	7/950 (0.7%)	1.39	11/1293 (0.9%)
All	All	1.54	24/2701 (0.9%)	1.37	30/3670 (0.8%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (24) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	173	GLU	CG-CD	8.75	1.65	1.51
1	A	32	GLU	CB-CG	7.88	1.67	1.52
1	A	81	VAL	CB-CG1	7.43	1.68	1.52
1	B	40	GLU	CB-CG	7.23	1.65	1.52
1	A	143	TYR	CD2-CE2	7.01	1.49	1.39
1	A	184	ASP	CB-CG	6.98	1.66	1.51
1	A	173	GLU	CB-CG	6.75	1.65	1.52
1	A	203	GLU	CG-CD	6.59	1.61	1.51
1	A	246	LYS	C-O	6.55	1.35	1.23
1	A	119	VAL	CB-CG2	-6.14	1.40	1.52
1	A	118	TYR	C-O	6.04	1.34	1.23
1	A	149	CYS	CB-SG	6.02	1.92	1.82
1	A	79	ALA	CA-CB	5.96	1.65	1.52
1	A	194	TYR	CD1-CE1	5.80	1.48	1.39
1	B	81	VAL	CB-CG2	5.72	1.64	1.52
1	B	134	ALA	CA-CB	5.71	1.64	1.52
1	B	53	TYR	N-CA	5.41	1.57	1.46
1	A	241	VAL	CA-CB	5.25	1.65	1.54

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	180	GLU	CG-CD	5.23	1.59	1.51
1	A	62	PHE	CD2-CE2	5.21	1.49	1.39
1	A	97	ARG	NE-CZ	5.14	1.39	1.33
1	B	144	ILE	CA-C	5.13	1.66	1.52
1	B	40	GLU	CD-OE1	5.11	1.31	1.25
1	B	136	VAL	CA-CB	5.04	1.65	1.54

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	41	ARG	NE-CZ-NH2	-12.79	113.91	120.30
1	A	218	ARG	NE-CZ-NH2	-9.50	115.55	120.30
1	A	41	ARG	NE-CZ-NH1	8.02	124.31	120.30
1	A	50	ASP	CB-CG-OD1	7.58	125.13	118.30
1	A	184	ASP	CB-CG-OD1	7.35	124.92	118.30
1	B	77	LEU	CA-CB-CG	6.92	131.22	115.30
1	A	209	LEU	CA-CB-CG	6.87	131.10	115.30
1	A	41	ARG	CG-CD-NE	-6.87	97.38	111.80
1	A	130	ASP	CB-CG-OD1	6.57	124.22	118.30
1	A	119	VAL	CG1-CB-CG2	-6.52	100.47	110.90
1	B	29	ASP	CB-CG-OD1	6.50	124.15	118.30
1	A	50	ASP	CB-CG-OD2	-6.50	112.45	118.30
1	A	116	ASP	CB-CG-OD1	6.40	124.06	118.30
1	B	76	LEU	CA-CB-CG	6.33	129.87	115.30
1	A	252	PRO	N-CA-C	6.13	128.05	112.10
1	A	130	ASP	CB-CG-OD2	-6.11	112.80	118.30
1	B	22	VAL	N-CA-C	5.92	126.99	111.00
1	B	29	ASP	CB-CG-OD2	-5.89	113.00	118.30
1	A	132	LEU	CA-CB-CG	5.68	128.37	115.30
1	B	50	ASP	CB-CG-OD2	-5.56	113.30	118.30
1	B	31	ASP	CB-CG-OD1	5.55	123.30	118.30
1	A	165	ASP	CB-CG-OD1	5.50	123.25	118.30
1	A	86	TYR	N-CA-C	-5.44	96.31	111.00
1	B	81	VAL	N-CA-C	-5.37	96.49	111.00
1	B	144	ILE	CB-CA-C	5.37	122.35	111.60
1	A	110	LEU	CA-CB-CG	5.26	127.40	115.30
1	B	77	LEU	CB-CG-CD2	5.19	119.82	111.00
1	A	173	GLU	CA-CB-CG	5.11	124.64	113.40
1	B	130	ASP	CB-CG-OD2	-5.08	113.72	118.30
1	A	41	ARG	CB-CA-C	-5.08	100.24	110.40

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	251	ALA	Peptide
1	A	85	ASP	Peptide

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1720	0	1611	43	0
1	B	929	0	847	18	0
2	A	35	0	0	7	0
2	B	20	0	0	2	0
All	All	2704	0	2458	60	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:252:PRO:CB	2:A:2034:HOH:O	1.64	1.32
1:A:210:GLU:O	1:A:214:LYS:HG3	1.69	0.92
1:A:97:ARG:CG	2:A:2014:HOH:O	2.18	0.90
1:B:129:LEU:HD22	1:B:133:LYS:HE3	1.54	0.89
1:A:252:PRO:N	2:A:2033:HOH:O	2.12	0.81
1:A:97:ARG:NE	2:A:2014:HOH:O	2.12	0.79
1:A:97:ARG:CD	2:A:2014:HOH:O	2.35	0.75
1:A:163:ILE:HG12	1:A:168:GLN:HG3	1.70	0.73
1:A:159:ASN:O	1:A:163:ILE:HG23	1.89	0.72
1:B:133:LYS:NZ	1:B:145:GLY:O	2.23	0.70
1:A:97:ARG:HG2	2:A:2014:HOH:O	1.88	0.69
1:A:234:ARG:O	1:A:238:ILE:HG13	1.94	0.68
1:B:129:LEU:CD2	1:B:133:LYS:HE3	2.24	0.68
1:A:23:THR:O	1:A:24:CYS:O	2.14	0.64
1:A:112:LYS:HE3	1:A:139:ASN:O	1.97	0.64
1:A:53:TYR:N	1:A:54:PRO:HD3	2.12	0.64

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:36:GLU:O	1:A:40:GLU:HG3	1.97	0.64
1:B:149:CYS:C	2:B:2020:HOH:O	2.39	0.60
1:A:110:LEU:HB3	1:A:119:VAL:HG12	1.82	0.60
1:B:69:ALA:O	1:B:73:THR:HG22	2.01	0.60
1:A:204:VAL:O	1:A:204:VAL:HG12	2.02	0.59
1:B:44:TYR:CD2	1:B:142:LEU:HD11	2.37	0.58
1:B:63:THR:O	1:B:67:LYS:HG3	2.04	0.58
1:A:100:VAL:HG23	1:A:100:VAL:O	2.05	0.57
1:A:179:GLN:HG3	1:A:191:ALA:HB1	1.87	0.56
1:B:48:LYS:HG3	1:B:106:PRO:HB2	1.87	0.56
1:A:53:TYR:H	1:A:54:PRO:HD3	1.70	0.55
1:B:97:ARG:HD3	1:B:98:TYR:CE2	2.43	0.53
1:A:207:ASP:OD1	1:A:207:ASP:N	2.38	0.53
1:A:100:VAL:O	1:A:100:VAL:CG2	2.58	0.52
1:B:46:VAL:HG23	1:B:76:LEU:HD11	1.92	0.51
1:B:110:LEU:C	1:B:110:LEU:HD23	2.31	0.51
1:A:69:ALA:O	1:A:73:THR:HG22	2.12	0.50
1:A:169:LEU:HG	1:A:202:HIS:CE1	2.46	0.50
1:A:27:CYS:HB3	1:A:80:THR:HG23	1.94	0.50
1:A:46:VAL:HG23	1:A:76:LEU:HD11	1.95	0.49
1:B:102:ASP:HB3	1:B:105:PHE:CE2	2.48	0.48
1:A:129:LEU:HD22	1:A:133:LYS:HE3	1.95	0.48
1:B:44:TYR:HD2	1:B:142:LEU:HD11	1.79	0.48
1:B:73:THR:HG23	2:B:2009:HOH:O	2.14	0.48
1:A:179:GLN:HG3	1:A:191:ALA:CB	2.44	0.47
1:A:248:THR:HG23	1:A:248:THR:O	2.14	0.47
1:A:102:ASP:HA	1:A:105:PHE:CE2	2.49	0.47
1:A:187:GLN:HE21	1:A:190:ASN:HD22	1.61	0.47
1:A:212:GLU:HA	1:A:212:GLU:OE1	2.16	0.46
1:A:97:ARG:NH2	1:A:116:ASP:OD1	2.49	0.45
1:A:163:ILE:HG12	1:A:168:GLN:CG	2.45	0.45
1:B:71:LYS:HE3	1:B:71:LYS:HB2	1.29	0.45
1:A:204:VAL:O	1:A:204:VAL:CG1	2.66	0.44
1:A:23:THR:O	1:A:23:THR:OG1	2.34	0.44
1:A:184:ASP:OD2	1:A:186:GLU:HB3	2.18	0.43
1:A:97:ARG:NH1	2:A:2015:HOH:O	2.52	0.43
1:A:121:LEU:O	1:A:122:PRO:C	2.55	0.43
1:A:102:ASP:OD2	1:A:102:ASP:N	2.43	0.43
1:A:134:ALA:HA	1:A:137:SER:OG	2.19	0.42
1:B:49:PHE:O	1:B:106:PRO:HA	2.20	0.42
1:A:85:ASP:HB3	1:A:89:LEU:HD23	2.02	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:64:ALA:O	1:B:68:SER:HB3	2.20	0.41
1:A:41:ARG:HD2	1:B:41:ARG:O	2.21	0.40
1:A:187:GLN:NE2	1:A:190:ASN:HD22	2.18	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	229/248 (92%)	211 (92%)	13 (6%)	5 (2%)	6 3
1	B	126/248 (51%)	116 (92%)	8 (6%)	2 (2%)	9 5
All	All	355/496 (72%)	327 (92%)	21 (6%)	7 (2%)	7 4

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	24	CYS
1	A	104	ASN
1	A	252	PRO
1	B	144	ILE
1	A	185	PRO
1	B	146	ARG
1	A	53	TYR

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	164/218 (75%)	145 (88%)	19 (12%)	5 3
1	B	90/218 (41%)	81 (90%)	9 (10%)	7 5
All	All	254/436 (58%)	226 (89%)	28 (11%)	6 4

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	23	THR
1	A	55	LYS
1	A	84	LYS
1	A	85	ASP
1	A	89	LEU
1	A	97	ARG
1	A	102	ASP
1	A	126	ASP
1	A	129	LEU
1	A	130	ASP
1	A	132	LEU
1	A	159	ASN
1	A	163	ILE
1	A	184	ASP
1	A	185	PRO
1	A	187	GLN
1	A	207	ASP
1	A	248	THR
1	A	250	THR
1	B	24	CYS
1	B	71	LYS
1	B	77	LEU
1	B	102	ASP
1	B	105	PHE
1	B	112	LYS
1	B	129	LEU
1	B	130	ASP
1	B	132	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	124	HIS
1	A	187	GLN
1	A	202	HIS

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [\(i\)](#)

There are no ligands in this entry.

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data i

6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	231/248 (93%)	0.06	3 (1%) 77 79	30, 47, 75, 96	0
1	B	128/248 (51%)	0.10	0 100 100	29, 46, 69, 75	0
All	All	359/496 (72%)	0.08	3 (0%) 86 87	29, 47, 73, 96	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	209	LEU	2.9
1	A	247	VAL	2.7
1	A	202	HIS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains i

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates i

There are no monosaccharides in this entry.

6.4 Ligands i

There are no ligands in this entry.

6.5 Other polymers i

There are no such residues in this entry.