



# Full wwPDB X-ray Structure Validation Report i

Mar 4, 2024 – 01:45 AM EST

PDB ID : 1C29

Title : CRYSTAL STRUCTURE OF THE COMPLEX OF BACTERIAL TRYPTOPHAN SYNTHASE WITH THE TRANSITION STATE ANALOGUE INHIBITOR 4-(2-HYDROXYPHENYLTHIO)-1-BUTENYLPHOSPHONIC ACID

Authors : Sachpatzidis, A.; Dealwis, C.; Lubetsky, J.B.; Liang, P.H.; Anderson, K.S.; Lolis, E.

Deposited on : 1999-07-23

Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the i symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## X-RAY DIFFRACTION


The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
 Validation Pipeline (wwPDB-VP) : 2.36

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	PLP	B	900	-	X	-	-

## 2 Entry composition [\(i\)](#)

There are 6 unique types of molecules in this entry. The entry contains 5120 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called TRYPTOPHAN SYNTHASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	264	Total	C 1992	N 1264	O 348	S 373	7	0	0

- Molecule 2 is a protein called TRYPTOPHAN SYNTHASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	387	Total	C 2934	N 1843	O 516	S 556	19	0	0

- Molecule 3 is 4-(2-HYDROXYPHENYLTHIO)-1-BUTENYLPHOSPHONIC ACID (three-letter code: HE1) (formula: C<sub>10</sub>H<sub>13</sub>O<sub>4</sub>PS).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C 16	O 10	P 4	S 1	0	0

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total Na 1 1	0	0

- Molecule 5 is PYRIDOXAL-5'-PHOSPHATE (three-letter code: PLP) (formula: C<sub>8</sub>H<sub>10</sub>NO<sub>6</sub>P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total C N O P 15 8 1 5 1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	68	Total O 68 68	0	0
6	B	94	Total O 94 94	0	0


### 3 Residue-property plots [\(i\)](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: TRYPTOPHAN SYNTHASE



- Molecule 2: TRYPTOPHAN SYNTHASE



## 4 Data and refinement statistics i

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	182.29Å    58.87Å    67.40Å 90.00°    94.40°    90.00°	Depositor
Resolution (Å)	30.00 – 2.30 42.22 – 2.30	Depositor EDS
% Data completeness (in resolution range)	91.0 (30.00-2.30) 92.1 (42.22-2.30)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) >$ <sup>1</sup>	4.94 (at 2.29Å)	Xtriage
Refinement program	X-PLOR 3.851	Depositor
$R$ , $R_{free}$	0.208 , 0.275 0.206 , 0.275	Depositor DCC
$R_{free}$ test set	2978 reflections (9.98%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	14.3	Xtriage
Anisotropy	0.545	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 45.8	EDS
L-test for twinning <sup>2</sup>	$<  L  > = 0.49$ , $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	5120	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	15.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.98% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $< |L| >$ ,  $< L^2 >$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: HE1, PLP, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.44	0/2031	0.71	0/2757
2	B	0.47	0/2992	0.78	2/4042 (0.0%)
All	All	0.46	0/5023	0.75	2/6799 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	151	LEU	CA-CB-CG	5.33	127.56	115.30
2	B	287	MET	N-CA-C	-5.14	97.12	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1992	0	1995	49	0
2	B	2934	0	2902	78	0
3	A	16	0	11	0	0
4	B	1	0	0	0	0
5	B	15	0	7	0	0
6	A	68	0	0	1	0
6	B	94	0	0	10	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	5120	0	4915	124	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (124) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:182:VAL:HG11	2:B:175:ARG:HG2	1.49	0.93
1:A:133:VAL:H	2:B:19:GLN:HE22	1.28	0.82
2:B:99:LYS:HE3	2:B:126:LEU:HB3	1.68	0.73
1:A:140:ARG:HD2	1:A:169:TYR:HB3	1.69	0.73
1:A:109:ASN:HD22	1:A:109:ASN:H	1.37	0.72
2:B:288:GLN:HE21	2:B:292:GLY:HA2	1.56	0.70
1:A:185:ALA:HB3	1:A:239:LYS:NZ	2.06	0.70
2:B:112:ALA:O	2:B:142:GLN:HG2	1.92	0.70
2:B:182:GLU:HB2	6:B:1095:HOH:O	1.92	0.69
2:B:89:ASN:HD22	2:B:205:GLN:NE2	1.91	0.69
2:B:89:ASN:HD22	2:B:205:GLN:HE22	1.43	0.67
2:B:81:LEU:HD13	2:B:88:THR:HB	1.79	0.65
2:B:327:ILE:HG23	2:B:331:GLU:HB3	1.79	0.65
1:A:211:GLY:O	1:A:212:PHE:HB2	1.96	0.64
1:A:26:GLY:HA3	1:A:76:VAL:HG21	1.83	0.61
1:A:53:PRO:HA	1:A:68:ASN:HD22	1.64	0.61
2:B:89:ASN:HB2	2:B:205:GLN:HE22	1.65	0.61
1:A:137:ALA:HB3	1:A:138:PRO:HD3	1.83	0.60
2:B:76:LYS:HD3	2:B:212:THR:OG1	2.01	0.59
2:B:202:ARG:HD3	2:B:312:GLN:HE22	1.66	0.59
2:B:216:ILE:HG21	2:B:224:PRO:HD3	1.84	0.59
2:B:90:GLN:HG3	2:B:201:VAL:HG13	1.85	0.58
2:B:360:LYS:O	2:B:364:GLU:HB2	2.04	0.58
1:A:220:VAL:O	1:A:224:VAL:HG23	2.04	0.58
1:A:185:ALA:HB3	1:A:239:LYS:HZ1	1.69	0.57
2:B:99:LYS:HZ3	2:B:128:LEU:HD21	1.70	0.56
2:B:271:LEU:HD12	2:B:309:VAL:HG21	1.87	0.56
1:A:86:ALA:HB2	1:A:121:VAL:HG22	1.87	0.56
2:B:87:LYS:HG2	6:B:1047:HOH:O	2.05	0.56
2:B:16:TYR:O	2:B:281:GLY:HA2	2.05	0.55
2:B:9:PHE:CD2	2:B:312:GLN:HG3	2.40	0.55
2:B:313:HIS:HD2	2:B:324:TYR:OH	1.88	0.55
2:B:211:GLU:O	2:B:215:GLN:HB2	2.07	0.54

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:202:ARG:HH21	2:B:312:GLN:HE22	1.54	0.54
2:B:236:ASN:HD22	2:B:375:ASN:HD22	1.56	0.53
2:B:99:LYS:NZ	2:B:128:LEU:HD21	2.24	0.52
2:B:115:HIS:CE1	2:B:189:GLY:HA2	2.45	0.52
1:A:188:ARG:HG2	1:A:219:GLN:NE2	2.25	0.52
2:B:27:GLN:HE22	2:B:103:LYS:NZ	2.08	0.52
1:A:3:ARG:HG3	1:A:94:THR:O	2.10	0.52
2:B:94:GLN:HE21	2:B:185:HIS:CE1	2.28	0.52
2:B:89:ASN:HB2	2:B:205:GLN:NE2	2.25	0.52
1:A:22:PHE:HD2	1:A:49:GLU:HB3	1.75	0.51
1:A:185:ALA:HB3	1:A:239:LYS:HZ2	1.73	0.51
2:B:202:ARG:HH21	2:B:312:GLN:NE2	2.08	0.51
2:B:195:HIS:HD2	6:B:1006:HOH:O	1.93	0.51
1:A:178:SER:HA	1:A:193:LEU:HD21	1.93	0.51
2:B:147:PHE:O	2:B:151:LEU:HB2	2.12	0.51
1:A:64:ILE:HG12	6:A:1031:HOH:O	2.11	0.50
1:A:165:GLN:HG2	1:A:169:TYR:CE1	2.46	0.50
1:A:153:ILE:HG23	1:A:177:LEU:HD13	1.93	0.50
2:B:53:ALA:HB1	2:B:81:LEU:HD22	1.93	0.50
1:A:163:LEU:HD21	1:A:196:LEU:HD22	1.92	0.49
2:B:108:ALA:HB3	2:B:132:ILE:HG12	1.94	0.49
2:B:131:ARG:HB2	6:B:1132:HOH:O	2.13	0.48
1:A:85:LEU:HD21	1:A:99:LEU:HD11	1.94	0.48
2:B:346:ILE:HG21	2:B:384:ILE:HD11	1.94	0.48
1:A:44:GLY:O	1:A:263:LYS:HD3	2.12	0.48
2:B:65:ILE:HD12	2:B:339:LEU:HD12	1.95	0.48
2:B:20:ILE:HG23	2:B:21:LEU:HD22	1.96	0.48
1:A:53:PRO:HA	1:A:68:ASN:ND2	2.27	0.48
2:B:49:LEU:O	2:B:55:ARG:HB2	2.13	0.48
1:A:57:PRO:HA	1:A:102:TYR:CZ	2.48	0.47
2:B:85:ALA:HB3	2:B:114:GLN:NE2	2.29	0.47
2:B:99:LYS:HZ1	2:B:128:LEU:HG	1.79	0.47
2:B:109:GLU:HG3	2:B:170:CYS:SG	2.54	0.47
2:B:134:MET:O	2:B:158:PRO:HA	2.14	0.47
2:B:76:LYS:HE3	2:B:215:GLN:HE22	1.79	0.47
2:B:329:ASP:O	2:B:333:LEU:HD22	2.15	0.47
2:B:109:GLU:O	2:B:115:HIS:HD2	1.98	0.46
1:A:133:VAL:N	2:B:19:GLN:HE22	2.04	0.46
2:B:234:GLY:O	2:B:313:HIS:HE1	1.98	0.46
2:B:185:HIS:HD2	6:B:1084:HOH:O	1.99	0.46
2:B:293:GLN:HA	2:B:293:GLN:OE1	2.15	0.46

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:61:LYS:HG3	2:B:74:TYR:CZ	2.51	0.46
2:B:87:LYS:HD2	2:B:114:GLN:HE21	1.81	0.46
2:B:90:GLN:CG	2:B:201:VAL:HG13	2.46	0.46
1:A:179:ARG:HG3	1:A:189:GLY:HA2	1.98	0.45
2:B:345:ILE:O	2:B:347:PRO:HD3	2.17	0.45
1:A:60:ASP:HB3	1:A:64:ILE:HB	2.00	0.44
1:A:20:VAL:O	1:A:232:ILE:HA	2.18	0.44
1:A:58:LEU:H	1:A:58:LEU:CD2	2.31	0.44
1:A:164:ARG:HG2	1:A:164:ARG:HH21	1.82	0.44
2:B:94:GLN:HE21	2:B:185:HIS:HE1	1.65	0.44
2:B:87:LYS:HD3	2:B:115:HIS:HA	1.99	0.44
2:B:105:GLU:HB3	2:B:184:ALA:HB2	2.00	0.44
2:B:313:HIS:CD2	2:B:324:TYR:OH	2.71	0.43
2:B:207:MET:O	2:B:211:GLU:HB2	2.19	0.43
1:A:150:PRO:HD2	1:A:171:ARG:HB2	2.01	0.43
1:A:52:VAL:HG22	1:A:101:MET:HE3	2.01	0.43
2:B:376:LEU:HD12	2:B:376:LEU:HA	1.90	0.43
2:B:142:GLN:HG3	6:B:1067:HOH:O	2.18	0.43
2:B:272:LYS:HD2	2:B:272:LYS:HA	1.70	0.42
1:A:153:ILE:HD12	6:B:1064:HOH:O	2.17	0.42
2:B:163:SER:O	2:B:165:THR:HG23	2.19	0.42
2:B:272:LYS:HE3	6:B:1001:HOH:O	2.19	0.42
2:B:110:THR:HG23	2:B:134:MET:HG3	2.01	0.42
1:A:141:GLN:HB3	1:A:145:ARG:NH1	2.35	0.42
1:A:165:GLN:HG2	1:A:169:TYR:HE1	1.85	0.42
2:B:144:PRO:HD2	6:B:1067:HOH:O	2.20	0.42
2:B:76:LYS:HA	2:B:76:LYS:HD2	1.83	0.42
2:B:71:THR:HB	2:B:362:MET:HG3	2.01	0.41
1:A:50:LEU:HD23	1:A:85:LEU:HD11	2.02	0.41
1:A:249:LYS:HE3	1:A:249:LYS:HB2	1.91	0.41
2:B:246:ASN:HB2	6:B:1139:HOH:O	2.20	0.41
1:A:149:ALA:HB1	1:A:171:ARG:HB3	2.03	0.41
2:B:94:GLN:NE2	2:B:185:HIS:HE1	2.18	0.41
1:A:211:GLY:O	1:A:212:PHE:CB	2.68	0.41
1:A:140:ARG:O	1:A:144:LEU:HD22	2.21	0.41
2:B:71:THR:HG23	2:B:372:LEU:HD22	2.03	0.41
2:B:100:ARG:HH21	2:B:100:ARG:HG3	1.86	0.41
2:B:229:ALA:HB1	2:B:236:ASN:HD21	1.86	0.41
1:A:14:ARG:HB3	1:A:16:GLU:HB2	2.03	0.41
1:A:85:LEU:HB3	1:A:121:VAL:CG1	2.51	0.41
1:A:20:VAL:HG22	1:A:47:ALA:HB3	2.02	0.40

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:220:GLU:HG2	2:B:371:LEU:HD22	2.04	0.40
2:B:78:GLU:HB2	2:B:376:LEU:O	2.22	0.40
2:B:90:GLN:OE1	2:B:94:GLN:OE1	2.39	0.40
1:A:54:PHE:H	1:A:68:ASN:ND2	2.19	0.40
1:A:70:ARG:NH2	1:A:242:GLU:HG3	2.36	0.40
1:A:109:ASN:HD22	1:A:109:ASN:N	2.12	0.40
2:B:59:LEU:HD22	2:B:215:GLN:HB3	2.03	0.40
1:A:248:PRO:O	1:A:252:LEU:HG	2.21	0.40
1:A:58:LEU:HD23	1:A:102:TYR:CE1	2.57	0.40

There are no symmetry-related clashes.

### 5.3 Torsion angles [\(i\)](#)

#### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	260/268 (97%)	252 (97%)	7 (3%)	1 (0%)	34 42
2	B	385/397 (97%)	369 (96%)	16 (4%)	0	100 100
All	All	645/665 (97%)	621 (96%)	23 (4%)	1 (0%)	47 58

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	212	PHE

#### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	206/208 (99%)	176 (85%)	30 (15%)	3   3
2	B	303/311 (97%)	259 (86%)	44 (14%)	3   3
All	All	509/519 (98%)	435 (86%)	74 (14%)	3   3

All (74) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	GLU
1	A	5	GLU
1	A	11	LEU
1	A	31	GLU
1	A	32	GLN
1	A	40	LEU
1	A	50	LEU
1	A	52	VAL
1	A	58	LEU
1	A	65	GLN
1	A	72	PHE
1	A	84	MET
1	A	85	LEU
1	A	87	LEU
1	A	99	LEU
1	A	101	MET
1	A	105	LEU
1	A	109	ASN
1	A	120	GLN
1	A	121	VAL
1	A	144	LEU
1	A	171	ARG
1	A	180	SER
1	A	188	ARG
1	A	193	LEU
1	A	201	LYS
1	A	209	LEU
1	A	220	VAL
1	A	255	LEU
1	A	263	LYS
2	B	15	MET
2	B	21	LEU
2	B	30	GLU
2	B	34	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	B	37	LYS
2	B	40	GLU
2	B	42	GLN
2	B	48	LEU
2	B	49	LEU
2	B	50	LYS
2	B	55	ARG
2	B	65	ILE
2	B	70	ARG
2	B	76	LYS
2	B	77	ARG
2	B	81	LEU
2	B	90	GLN
2	B	101	MET
2	B	103	LYS
2	B	109	GLU
2	B	114	GLN
2	B	143	SER
2	B	152	MET
2	B	166	LEU
2	B	188	LEU
2	B	205	GLN
2	B	207	MET
2	B	215	GLN
2	B	217	LEU
2	B	219	LYS
2	B	236	ASN
2	B	263	GLU
2	B	266	GLU
2	B	272	LYS
2	B	295	GLU
2	B	297	SER
2	B	333	LEU
2	B	343	GLU
2	B	355	LEU
2	B	359	LEU
2	B	365	GLN
2	B	370	GLN
2	B	372	LEU
2	B	382	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (22) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	6	ASN
1	A	68	ASN
1	A	109	ASN
1	A	120	GLN
1	A	244	ASN
2	B	19	GLN
2	B	27	GLN
2	B	64	ASN
2	B	82	HIS
2	B	90	GLN
2	B	114	GLN
2	B	142	GLN
2	B	171	ASN
2	B	185	HIS
2	B	195	HIS
2	B	205	GLN
2	B	215	GLN
2	B	236	ASN
2	B	288	GLN
2	B	312	GLN
2	B	313	HIS
2	B	370	GLN

### 5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [\(i\)](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	PLP	B	900	2	15,15,16	2.15	7 (46%)	20,22,23	4.74	13 (65%)
3	HE1	A	270	-	15,16,16	3.00	4 (26%)	16,21,21	1.36	3 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	PLP	B	900	2	-	0/6/6/8	0/1/1/1
3	HE1	A	270	-	-	3/6/9/9	0/1/1/1

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	270	HE1	P1-O3	7.67	1.61	1.48
3	A	270	HE1	P1-C1	-5.42	1.69	1.76
3	A	270	HE1	P1-O1	4.75	1.60	1.54
3	A	270	HE1	P1-O2	4.19	1.60	1.54
5	B	900	PLP	C4A-C4	-4.05	1.43	1.51
5	B	900	PLP	O4P-C5A	-3.26	1.32	1.45
5	B	900	PLP	C3-C2	-3.05	1.37	1.40
5	B	900	PLP	P-O3P	-2.61	1.44	1.54
5	B	900	PLP	P-O4P	-2.44	1.52	1.60
5	B	900	PLP	C2-N1	2.38	1.38	1.33
5	B	900	PLP	P-O2P	-2.22	1.46	1.54

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	900	PLP	C4A-C4-C5	14.60	135.98	120.94
5	B	900	PLP	C4A-C4-C3	-8.99	105.26	120.50
5	B	900	PLP	C6-C5-C4	5.35	122.37	118.16
5	B	900	PLP	C5A-C5-C6	-4.86	111.38	119.37
5	B	900	PLP	O4P-C5A-C5	-4.60	100.58	109.35
5	B	900	PLP	O3P-P-O4P	-4.45	94.89	106.73

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	270	HE1	C4-C3-S1	-3.56	104.06	112.28
5	B	900	PLP	C3-C4-C5	-3.39	115.08	118.74
5	B	900	PLP	O2P-P-O4P	-3.30	97.95	106.73
5	B	900	PLP	O3P-P-O1P	2.67	121.13	110.68
5	B	900	PLP	O4P-P-O1P	2.52	113.54	106.47
5	B	900	PLP	C2A-C2-N1	2.51	122.57	117.67
3	A	270	HE1	C3-S1-C8	2.28	107.82	103.04
5	B	900	PLP	C3-C2-N1	-2.27	117.84	120.77
3	A	270	HE1	C9-C8-C7	2.21	121.19	119.15
5	B	900	PLP	C5-C6-N1	-2.10	120.31	123.82

There are no chirality outliers.


All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	270	HE1	C2-C1-P1-O3
3	A	270	HE1	C9-C8-S1-C3
3	A	270	HE1	C7-C8-S1-C3

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data i

### 6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	264/268 (98%)	-0.07	3 (1%) 80 85	6, 16, 35, 54	0
2	B	387/397 (97%)	-0.20	0 100 100	4, 12, 25, 38	0
All	All	651/665 (97%)	-0.15	3 (0%) 91 94	4, 13, 29, 54	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	187	ASN	2.2
1	A	249	LYS	2.1
1	A	189	GLY	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains i

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates i

There are no monosaccharides in this entry.

### 6.4 Ligands i

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	HE1	A	270	16/16	0.96	0.15	11,16,18,21	0
4	NA	B	1168	1/1	0.97	0.10	18,18,18,18	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	PLP	B	900	15/16	0.97	0.17	11,14,18,19	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.