Full wwPDB X-ray Structure Validation Report (i)

Aug 23, 2023 – 02:47 PM EDT

PDB ID	:	3C4W
Title	:	Crystal Structure of G protein coupled receptor kinase 1 bound to ATP and magnesium chloride at 2.7A
Authors	:	Singh, P.; Tesmer, J.J.G.
Deposited on	:	2008-01-30
Resolution	:	2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.35
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	543	3% 80% 10% • 9%
1	B	543	6% 82% 12% • •

2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 8390 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Rhodopsin kinase.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
1	A	494	Total C N O S 3989 2558 690 722 19	0	3	0
1	B	519	Total C N O P S 4167 2666 721 760 1 19	0	2	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	536	VAL	-	expression tag	UNP P28327
A	537	ASP	-	expression tag	UNP P28327
A	538	HIS	-	expression tag	UNP P28327
A	539	HIS	-	expression tag	UNP P28327
A	540	HIS	-	expression tag	UNP P28327
A	541	HIS	-	expression tag	UNP P28327
A	542	HIS	-	expression tag	UNP P28327
A	543	HIS	-	expression tag	UNP P28327
B	536	VAL	-	expression tag	UNP P28327
B	537	ASP	-	expression tag	UNP P28327
B	538	HIS	-	expression tag	UNP P28327
B	539	HIS	-	expression tag	UNP P28327
B	540	HIS	-	expression tag	UNP P28327
B	541	HIS	-	expression tag	UNP P28327
B	542	HIS	-	expression tag	UNP P28327
B	543	HIS	-	expression tag	UNP P28327

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Mg 1 1	0	0
2	B	2	Total Mg 2 2	0	0

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Cl		
3	A	3	3	3	0	0

- Molecule 4 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: $C_{10}H_{16}N_5O_{13}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
4	A	1	31	10	5	13	3	0	0
4	B	1	31	10	5	13	3	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
5	A	109	109	109	0	0
5	B	57	57	57	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Rhodopsin kinase

Chain A:



• Molecule 1: Rhodopsin kinase

Chain B:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α, β, γ	202.89Å 55.05Å 122.70Å 90.00° 100.82° 90.00°	Depositor
Resolution (Å)	19.98 – 2.70 19.97 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.8 (19.98-2.70) 99.8 (19.97-2.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	3.20 (at 2.71Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.192 , (Not available) 0.194 , 0.207	Depositor DCC
R_{free} test set	1859 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å^2)	48.1	Xtriage
Anisotropy	0.078	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.34 , 49.6	EDS
L-test for twinning[2]	$< \|L\| > = 0.50$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.94	EDS
Total number of atoms	8390	wwPDB-VP
Average B, all atoms (Å^2)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.64% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, CL, ATP, TPO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.60	1/4084 (0.0%)	0.68	0/5509
1	B	0.53	0/4253	0.65	1/5736 (0.0%)
All	All	0.57	1/8337 (0.0%)	0.66	1/11245 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	443	CYS	CB-SG	-6.04	1.72	1.82

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	B	447	ARG	NE-CZ-NH2	-6.57	117.01	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3989	0	3948	40	0
1	B	4167	0	4116	37	0
2	A	1	0	0	0	0
2	B	2	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	3	0	0	0	0
4	A	31	0	12	0	0
4	B	31	0	12	1	0
5	A	109	0	0	0	0
5	B	57	0	0	0	0
All	All	8390	0	8088	75	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (75) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:81:GLN:HE21	1:A:81:GLN:H	1.14	0.93
1:B:423:PHE:CE2	1:B:427:LEU:HD11	2.27	0.68
1:A:88:LEU:HD12	1:A:110:LEU:HD11	1.78	0.64
1:A:157:ALA:HB3	1:A:158:PRO:HD3	1.79	0.64
1:A:297:GLN:OE1	1:A:328:VAL:HG22	1.98	0.62
1:B:344:THR:HG23	1:B:345:LYS:HG3	1.81	0.62
1:A:110:LEU:HD12	1:A:114:TYR:CE1	2.35	0.62
1:A:88:LEU:HD23	1:A:150:VAL:CG2	2.30	0.60
1:A:522:THR:HG23	1:A:524:VAL:HG23	1.83	0.60
1:A:138:ALA:HA	1:A:142:LEU:HD22	1.85	0.59
1:B:193:LEU:HD13	4:B:562:ATP:C2	2.39	0.58
1:A:405:VAL:O	1:A:430:LYS:HE3	2.04	0.57
1:A:56:LEU:HA	1:A:61:MET:HE2	1.87	0.56
1:A:166:LEU:HD22	1:B:166:LEU:HD22	1.86	0.56
1:B:446:LEU:C	1:B:446:LEU:HD23	2.26	0.56
1:A:142:LEU:HD12	1:A:142:LEU:O	2.06	0.56
1:A:81:GLN:H	1:A:81:GLN:NE2	1.96	0.56
1:B:279:VAL:HG21	1:B:290:ARG:HH12	1.71	0.55
1:A:359:GLU:HG2	1:A:432:PRO:HG3	1.88	0.55
1:B:111:ARG:HG2	1:B:115:LEU:HD12	1.89	0.54
1:B:447:ARG:HG2	1:B:457:TRP:CZ2	2.43	0.53
1:A:88:LEU:HD23	1:A:150:VAL:HG21	1.90	0.53
1:B:157:ALA:HB3	1:B:158:PRO:HD3	1.92	0.52
1:B:142:LEU:HD23	1:B:143:PHE:CE1	2.45	0.51
1:A:360:LEU:C	1:A:360:LEU:HD23	2.30	0.51
1:B:292:ILE:HD13	1:B:451:LEU:O	2.11	0.50
1:A:288:GLU:HB3	1:A:289:PRO:HD3	1.93	0.50
1:B:59:GLU:HG3	1:B:63:LEU:HD12	1.94	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:322:LEU:HG	1:B:328:VAL:HG12	1.93	0.49
1:A:110:LEU:HD12	1:A:114:TYR:HE1	1.73	0.49
1:A:262:LEU:HD21	1:A:506:PHE:CZ	2.47	0.49
1:B:395:LYS:HA	1:B:395:LYS:HE2	1.94	0.49
1:B:142:LEU:HD23	1:B:143:PHE:CZ	2.48	0.48
1:B:396:VAL:HG13	1:B:396:VAL:O	2.13	0.48
1:B:447:ARG:HG2	1:B:457:TRP:CH2	2.49	0.48
1:A:445:GLY:O	1:A:448:THR:HB	2.13	0.48
1:B:238:ILE:O	1:B:242:VAL:HG13	2.14	0.48
1:B:491:LYS:O	1:B:493:VAL:HG23	2.14	0.48
1:A:221:LYS:HB3	1:A:490:VAL:CG2	2.44	0.47
1:A:77[B]:ARG:HG3	1:A:86:LEU:HD21	1.98	0.46
1:A:99:ASP:O	1:A:103:ARG:HG3	2.16	0.46
1:A:111:ARG:HG2	1:A:115:LEU:HD12	1.97	0.45
1:B:8:TPO:O2P	1:B:93:GLU:OE1	2.34	0.45
1:B:66:PRO:O	1:B:70:ARG:HB2	2.16	0.45
1:B:518:GLU:O	1:B:522:THR:HB	2.16	0.45
1:A:322:LEU:HD12	1:A:328:VAL:HG12	1.98	0.45
1:A:110:LEU:HD23	1:A:143:PHE:CZ	2.52	0.45
1:A:532:ARG:HH21	1:B:530:VAL:HG12	1.82	0.45
1:A:522:THR:CG2	1:A:524:VAL:HG23	2.45	0.45
1:B:76:LEU:HD22	1:B:82:HIS:HB3	1.98	0.44
1:B:447:ARG:HD3	1:B:457:TRP:CE3	2.52	0.44
1:B:271:ASP:HB3	1:B:321:LEU:HD23	2.00	0.43
1:B:230:GLN:O	1:B:234:VAL:HG23	2.18	0.43
1:B:276:ILE:HD13	1:B:381:GLU:HG2	1.99	0.43
1:A:446:LEU:C	1:A:446:LEU:HD23	2.39	0.43
1:B:127:ASP:HB2	1:B:130:THR:H	1.84	0.43
1:A:216:LYS:HE2	1:A:218:LEU:HD21	2.00	0.43
1:A:206:MET:SD	1:A:208:ALA:HB3	2.59	0.43
1:A:405:VAL:O	1:A:430:LYS:CE	2.66	0.43
1:B:192:VAL:HG22	1:B:202:PHE:CE2	2.54	0.42
1:B:144:GLN:N	1:B:144:GLN:HE21	2.18	0.42
1:B:144:GLN:NE2	1:B:144:GLN:CA	2.83	0.42
1:A:518:GLU:O	1:A:522:THR:HB	2.19	0.42
1:A:110:LEU:HD23	1:A:143:PHE:CE2	2.55	0.41
1:B:267:MET:HE2	1:B:329:ARG:HB2	2.01	0.41
1:A:199:GLY:HA2	1:A:487:PHE:CE1	2.55	0.41
1:A:77[B]:ARG:HG3	1:A:86:LEU:CD2	2.50	0.41
1:A:44:LEU:HD23	1:A:44:LEU:HA	1.90	0.41
1:A:184:GLU:HB2	1:A:256:THR:HG22	2.03	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:236:LYS:NZ	1:A:508:SER:OG	2.51	0.41
1:A:357:ALA:HB1	1:A:359:GLU:OE2	2.21	0.41
1:B:9:VAL:O	1:B:520:ILE:HG21	2.21	0.41
1:B:389:PHE:CD1	1:B:405:VAL:HG22	2.56	0.40
1:B:224:LYS:NZ	1:B:499:ASP:OD2	2.47	0.40
1:B:359:GLU:HG2	1:B:432:PRO:HG3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	493/543 (91%)	473 (96%)	20 (4%)	0	100	100
1	B	514/543 (95%)	494 (96%)	17 (3%)	3 (1%)	25	50
All	All	1007/1086 (93%)	967 (96%)	37 (4%)	3 (0%)	41	66

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	496	GLU
1	B	490	VAL
1	B	492	GLY

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	414/448 (92%)	399 (96%)	15 (4%)	35	64
1	B	432/448 (96%)	416 (96%)	16 (4%)	34	63
All	All	846/896 (94%)	815 (96%)	31 (4%)	35	63

All (31) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	50	LEU
1	A	53	SER
1	A	77[A]	ARG
1	A	77[B]	ARG
1	A	81	GLN
1	A	105	GLN
1	A	118	GLN
1	A	206	MET
1	A	217	LYS
1	A	226	ARG
1	A	386	ARG
1	A	397	GLU
1	A	415	LYS
1	A	448	THR
1	A	522	THR
1	B	13	SER
1	B	45	SER
1	B	50	LEU
1	B	105	GLN
1	B	120	GLN
1	B	140	ASP
1	B	144	GLN
1	B	206	MET
1	B	242	VAL
1	B	344	THR
1	B	395	LYS
1	B	407	GLU
1	B	448	THR
1	B	496	GLU
1	B	522	THR
1	B	532	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	81	GLN
1	A	87	GLN
1	A	105	GLN
1	A	118	GLN
1	A	309	ASN
1	A	429	GLN
1	B	73	GLN
1	B	87	GLN
1	B	108	GLN
1	B	120	GLN
1	B	144	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
1	TPO	B	8	1	8,10,11	0.90	0	10,14,16	1.04	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TPO	B	8	1	-	0/9/11/13	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	8	TPO	1	0

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 8 ligands modelled in this entry, 6 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
4	ATP	A	562	2	26,33,33	0.97	1 (3%)	31,52,52	1.36	2 (6%)
4	ATP	B	562	2	26,33,33	0.92	0	31,52,52	1.51	6 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ATP	A	562	2	-	2/18/38/38	0/3/3/3
4	ATP	B	562	2	-	0/18/38/38	0/3/3/3

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	562	ATP	C5-C4	2.19	1.46	1.40

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
4	A	562	ATP	N3-C2-N1	-4.09	122.28	128.68
4	B	562	ATP	N3-C2-N1	-3.99	122.44	128.68
4	B	562	ATP	PB-O3B-PG	-3.06	122.32	132.83
4	B	562	ATP	C1'-N9-C4	-2.91	121.53	126.64
4	A	562	ATP	C1'-N9-C4	-2.87	121.59	126.64
4	B	562	ATP	O2B-PB-O1B	2.66	125.38	112.24
4	B	562	ATP	PA-O3A-PB	-2.40	124.60	132.83
4	B	562	ATP	C4-C5-N7	-2.02	107.29	109.40

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	562	ATP	C5'-O5'-PA-O1A
4	A	562	ATP	C5'-O5'-PA-O3A

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	562	ATP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand ATP A 562



Ligand ATP B 562



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	494/543 (90%)	0.06	17 (3%)	45	45	11, 24, 49, 68	0
1	B	518/543 (95%)	0.20	35 (6%)	17	15	11, 24, 49, 60	0
All	All	1012/1086 (93%)	0.13	52 (5%)	28	26	11, 24, 49, 68	0

All (52) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	474	ARG	5.8
1	A	395	LYS	5.0
1	A	140	ASP	4.9
1	B	279	VAL	4.0
1	B	395	LYS	3.8
1	B	282	ASP	3.8
1	A	33	ARG	3.7
1	B	479	LYS	3.7
1	B	491	LYS	3.6
1	B	341	ALA	3.6
1	B	21	SER	3.5
1	B	33	ARG	3.4
1	B	24	ALA	3.4
1	B	414	ASP	3.2
1	B	120	GLN	3.1
1	B	498	ALA	3.0
1	B	480	ASN	2.9
1	B	283	ASN	2.9
1	A	138	ALA	2.9
1	A	101	ALA	2.9
1	B	129	GLU	2.7
1	B	478	ALA	2.7
1	A	483	ASP	2.7
1	B	230	GLN	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	399	LYS	2.6
1	A	129	GLU	2.6
1	A	49	ALA	2.6
1	A	472	ASP	2.5
1	A	48	GLU	2.5
1	A	141	GLY	2.5
1	A	393	GLY	2.4
1	A	283	ASN	2.4
1	B	492	GLY	2.4
1	B	501	GLU	2.4
1	B	105	GLN	2.3
1	B	347	LYS	2.3
1	A	486	ALA	2.3
1	B	100	ASP	2.3
1	B	489	THR	2.2
1	A	137	GLY	2.2
1	B	533	PRO	2.2
1	B	53	SER	2.2
1	B	391	ALA	2.2
1	B	464	MET	2.2
1	B	197	GLY	2.1
1	B	14	ALA	2.1
1	A	280	ASP	2.1
1	B	49	ALA	2.1
1	B	7	GLU	2.1
1	B	472	ASP	2.1
1	A	139	GLY	2.1
1	B	393	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
1	TPO	B	8	11/12	0.96	0.13	22,31,38,40	0

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	MG	B	563	1/1	0.78	0.15	49,49,49,49	0
2	MG	A	564	1/1	0.94	0.06	18,18,18,18	0
2	MG	B	564	1/1	0.95	0.10	24,24,24,24	0
3	CL	A	565	1/1	0.95	0.20	21,21,21,21	0
3	CL	A	567	1/1	0.97	0.23	16,16,16,16	0
4	ATP	B	562	31/31	0.97	0.13	6,17,23,25	0
4	ATP	A	562	31/31	0.98	0.09	6,16,19,20	0
3	CL	A	566	1/1	0.98	0.29	23,23,23,23	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around ATP B 562:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around ATP A 562:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.