



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 19, 2023 – 12:09 AM EDT

PDB ID : 5C6R  
Title : Crystal structure of PH domain of ASAP1  
Authors : Xia, D.; Tang, W.K.  
Deposited on : 2015-06-23  
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 2.35.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	150	
1	B	150	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	PGE	A	502	-	-	X	-

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 1838 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Arf-GAP.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	110	894	561	165	165	3	0	1	0
1	B	108	870	545	160	162	3	0	0	0

There are 46 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	302	MET	-	expression tag	UNP Q9QWY8
A	303	GLY	-	expression tag	UNP Q9QWY8
A	304	HIS	-	expression tag	UNP Q9QWY8
A	305	HIS	-	expression tag	UNP Q9QWY8
A	306	HIS	-	expression tag	UNP Q9QWY8
A	307	HIS	-	expression tag	UNP Q9QWY8
A	308	HIS	-	expression tag	UNP Q9QWY8
A	309	HIS	-	expression tag	UNP Q9QWY8
A	310	HIS	-	expression tag	UNP Q9QWY8
A	311	HIS	-	expression tag	UNP Q9QWY8
A	312	HIS	-	expression tag	UNP Q9QWY8
A	313	HIS	-	expression tag	UNP Q9QWY8
A	314	SER	-	expression tag	UNP Q9QWY8
A	315	SER	-	expression tag	UNP Q9QWY8
A	316	GLY	-	expression tag	UNP Q9QWY8
A	317	HIS	-	expression tag	UNP Q9QWY8
A	318	ILE	-	expression tag	UNP Q9QWY8
A	319	ASP	-	expression tag	UNP Q9QWY8
A	320	ASP	-	expression tag	UNP Q9QWY8
A	321	ASP	-	expression tag	UNP Q9QWY8
A	322	LYS	-	expression tag	UNP Q9QWY8
A	323	HIS	-	expression tag	UNP Q9QWY8
A	324	MET	-	expression tag	UNP Q9QWY8
B	302	MET	-	expression tag	UNP Q9QWY8
B	303	GLY	-	expression tag	UNP Q9QWY8

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	304	HIS	-	expression tag	UNP Q9QWY8
B	305	HIS	-	expression tag	UNP Q9QWY8
B	306	HIS	-	expression tag	UNP Q9QWY8
B	307	HIS	-	expression tag	UNP Q9QWY8
B	308	HIS	-	expression tag	UNP Q9QWY8
B	309	HIS	-	expression tag	UNP Q9QWY8
B	310	HIS	-	expression tag	UNP Q9QWY8
B	311	HIS	-	expression tag	UNP Q9QWY8
B	312	HIS	-	expression tag	UNP Q9QWY8
B	313	HIS	-	expression tag	UNP Q9QWY8
B	314	SER	-	expression tag	UNP Q9QWY8
B	315	SER	-	expression tag	UNP Q9QWY8
B	316	GLY	-	expression tag	UNP Q9QWY8
B	317	HIS	-	expression tag	UNP Q9QWY8
B	318	ILE	-	expression tag	UNP Q9QWY8
B	319	ASP	-	expression tag	UNP Q9QWY8
B	320	ASP	-	expression tag	UNP Q9QWY8
B	321	ASP	-	expression tag	UNP Q9QWY8
B	322	LYS	-	expression tag	UNP Q9QWY8
B	323	HIS	-	expression tag	UNP Q9QWY8
B	324	MET	-	expression tag	UNP Q9QWY8

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	P		
2	A	1	5	4	1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	P		
2	B	1	5	4	1	0	0

- Molecule 3 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula: C<sub>6</sub>H<sub>14</sub>O<sub>4</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
3	A	1	10	6	4	0	0
3	A	1	10	6	4	0	0
3	A	1	10	6	4	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	20	20	20	0	0
4	B	14	14	14	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Arf-GAP



- Molecule 1: Arf-GAP



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 43	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	48.12Å 48.12Å 111.33Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.12 – 1.80 34.03 – 1.80	Depositor EDS
% Data completeness (in resolution range)	95.3 (48.12-1.80) 95.4 (34.03-1.80)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.75 (at 1.81Å)	Xtrriage
Refinement program	REFMAC 5.8.0107	Depositor
R, $R_{free}$	0.208 , 0.260 0.214 , 0.256	Depositor DCC
$R_{free}$ test set	1165 reflections (5.21%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	28.7	Xtrriage
Anisotropy	0.367	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.42 , 62.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.066 for h,-k,-l	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	1838	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	48.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.29% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PGE, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.90	1/913 (0.1%)	0.97	1/1224 (0.1%)
1	B	0.92	0/885	1.02	1/1186 (0.1%)
All	All	0.91	1/1798 (0.1%)	0.99	2/2410 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	337	GLU	CG-CD	5.96	1.60	1.51

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	337	GLU	OE1-CD-OE2	-5.76	116.39	123.30
1	B	346	LEU	CA-CB-CG	5.24	127.36	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	894	0	902	13	0
1	B	870	0	876	13	0
2	A	5	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	5	0	0	0	0
3	A	30	0	42	7	0
4	A	20	0	0	4	0
4	B	14	0	0	1	0
All	All	1838	0	1820	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:386:LEU:HD11	1:A:433:ALA:HB3	1.70	0.72
3:A:502:PGE:H4	4:A:601:HOH:O	1.93	0.68
1:B:386:LEU:HD11	1:B:433:ALA:HB3	1.75	0.68
3:A:502:PGE:C4	4:A:601:HOH:O	2.47	0.61
3:A:502:PGE:C6	1:B:412:GLN:HE22	2.13	0.61
1:A:376:SER:HA	1:A:378:ARG:N	2.17	0.59
3:A:502:PGE:H6	1:B:412:GLN:HE22	1.67	0.59
1:B:414:GLU:HG3	4:B:601:HOH:O	2.02	0.59
1:A:351:ASP:N	1:A:351:ASP:OD1	2.38	0.55
1:A:333:GLN:O	1:A:431:GLU:HG3	2.07	0.55
3:A:502:PGE:H6	4:A:601:HOH:O	2.08	0.52
1:A:386:LEU:HD12	1:A:386:LEU:N	2.25	0.51
3:A:502:PGE:H4	1:B:412:GLN:OE1	2.10	0.50
1:A:376:SER:HA	1:A:377:ASN:C	2.32	0.50
1:A:386:LEU:HD11	1:A:433:ALA:CB	2.41	0.49
1:B:393:ASN:HB3	1:B:399:SER:HB2	1.96	0.48
1:B:386:LEU:HD11	1:B:433:ALA:CB	2.46	0.45
1:A:435:THR:O	1:A:439:ARG:HG2	2.16	0.44
1:B:407:ARG:HD2	1:B:409:TYR:CZ	2.52	0.44
1:B:403:ILE:CD1	1:B:408:THR:HG23	2.48	0.43
1:A:386:LEU:N	1:A:386:LEU:CD1	2.82	0.43
1:A:407:ARG:HD2	1:A:409:TYR:CZ	2.53	0.43
3:A:502:PGE:H62	1:B:412:GLN:HE22	1.80	0.43
1:B:376:SER:HA	1:B:378:ARG:H	1.84	0.43
1:A:403:ILE:CD1	1:A:408:THR:HG23	2.50	0.42
1:A:336:LYS:HG2	4:A:612:HOH:O	2.18	0.42
1:B:403:ILE:HD13	1:B:408:THR:HG23	2.02	0.41
1:A:439:ARG:HA	1:A:439:ARG:HD3	1.87	0.41
1:B:427:THR:O	1:B:431:GLU:HG3	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	109/150 (73%)	105 (96%)	3 (3%)	1 (1%)	17	6
1	B	106/150 (71%)	105 (99%)	1 (1%)	0	100	100
All	All	215/300 (72%)	210 (98%)	4 (2%)	1 (0%)	29	15

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	377	ASN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	97/131 (74%)	93 (96%)	4 (4%)	30	16
1	B	94/131 (72%)	89 (95%)	5 (5%)	22	9
All	All	191/262 (73%)	182 (95%)	9 (5%)	26	12

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	351	ASP
1	A	375	THR

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	402	LEU
1	A	407	ARG
1	B	351	ASP
1	B	375	THR
1	B	378	ARG
1	B	407	ARG
1	B	438	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	412	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	PGE	A	502	-	9,9,9	0.28	0	8,8,8	1.12	1 (12%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	PO4	B	501	-	4,4,4	0.73	0	6,6,6	0.78	0
3	PGE	A	503	-	9,9,9	0.41	0	8,8,8	0.48	0
3	PGE	A	504	-	9,9,9	0.38	0	8,8,8	0.63	0
2	PO4	A	501	-	4,4,4	1.00	0	6,6,6	0.91	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PGE	A	504	-	-	4/7/7/7	-
3	PGE	A	502	-	-	3/7/7/7	-
3	PGE	A	503	-	-	4/7/7/7	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	502	PGE	O3-C4-C3	2.52	121.77	110.39

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	504	PGE	O3-C5-C6-O4
3	A	502	PGE	O2-C3-C4-O3
3	A	503	PGE	O3-C5-C6-O4
3	A	502	PGE	O1-C1-C2-O2
3	A	504	PGE	C6-C5-O3-C4
3	A	503	PGE	O2-C3-C4-O3
3	A	503	PGE	C3-C4-O3-C5
3	A	503	PGE	C4-C3-O2-C2
3	A	502	PGE	C3-C4-O3-C5
3	A	504	PGE	O2-C3-C4-O3
3	A	504	PGE	C1-C2-O2-C3

There are no ring outliers.

1 monomer is involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	502	PGE	7	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	110/150 (73%)	0.97	17 (15%) 2   1	23, 43, 97, 113	0
1	B	108/150 (72%)	1.13	15 (13%) 2   2	25, 42, 87, 109	0
All	All	218/300 (72%)	1.05	32 (14%) 2   1	23, 42, 97, 113	0

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	394	ALA	11.0
1	B	377	ASN	10.7
1	B	438	PHE	10.2
1	A	353	ILE	10.0
1	B	376	SER	9.2
1	A	354	ARG	7.1
1	B	375	THR	6.3
1	A	438	PHE	6.2
1	A	352	GLY	5.5
1	B	333	GLN	5.3
1	A	374	ALA	4.5
1	A	377	ASN	4.0
1	B	405	HIS	3.9
1	B	439	ARG	3.9
1	B	334	GLY	3.8
1	B	374	ALA	3.7
1	B	406	ASN	3.6
1	B	396	ASP	3.6
1	A	351	ASP	3.5
1	B	351	ASP	3.4
1	A	440	GLY	3.2
1	B	397	LYS	3.1
1	A	355	LYS	3.1
1	A	356	VAL	2.9

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	439	ARG	2.9
1	B	378	ARG	2.7
1	A	376	SER	2.6
1	A	331	GLN	2.4
1	A	375	THR	2.4
1	A	357	TRP	2.3
1	A	405[A]	HIS	2.0
1	A	395	GLU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	PGE	A	504	10/10	0.86	0.17	64,66,78,86	0
3	PGE	A	503	10/10	0.89	0.15	51,56,67,70	0
3	PGE	A	502	10/10	0.90	0.19	37,45,53,57	0
2	PO4	B	501	5/5	0.97	0.07	60,64,66,67	0
2	PO4	A	501	5/5	0.97	0.10	66,67,69,74	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.