

Nov 20, 2022 – 02:52 pm GMT

PDB ID	:	2C7D
EMDB ID	:	EMD-1181
Title	:	Fitted coordinates for GroEL-ADP7-GroES Cryo-EM complex (EMD-1181)
Authors	:	Ranson, N.A.; Clare, D.K.; Farr, G.W.; Houldershaw, D.; Horwich, A.L.;
		Saibil, H.R.
Deposited on	:	2005-11-22
Resolution	:	8.70  Å(reported)
Based on initial model	:	1AON

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/EMValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

:	0.0.1. dev 43
:	4.02b-467
:	20191225.v01 (using entries in the PDB archive December 25th 2019)
:	1.9.9
:	Engh & Huber (2001)
:	Parkinson et al. (1996)
:	2.31.2
	: : : : :

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $ELECTRON\ MICROSCOPY$ 

The reported resolution of this entry is 8.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Matria	Whole archive	EM structures	
Metric	$(\# {\rm Entries})$	$(\# {\rm Entries})$	
Clashscore	158937	4297	
Ramachandran outliers	154571	4023	
Sidechain outliers	154315	3826	

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion < 40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain					
1	А	547	8%	5% •				
1	В	547	90%	6% •				
1	С	547	9%	• •				
1	D	547	91%	5% •				
1	Е	547	90%	6% •				
1	F	547	8%	5% •				
1	G	547	8%	• •				
1	Н	547	20%	5% •				



Mol	Chain	Length	Quality of chain	
1	Ι	547	20%	
1	J	547	91%	5% •
1	Κ	547	90%	6% ·
1	L	547	90%	5% • •
1	М	547	91%	5% •
1	Ν	547	90%	5% •
2	Ο	97	82%	11% • •
2	Р	97	82%	12% • •
2	Q	97	81%	13% • •
2	R	97	80%	14% • •
2	S	97	77%	18% • •
2	Т	97	8%	13% • •
2	U	97	77%	16% • •



## 2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 57946 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms					AltConf	Trace		
1	Δ	525	Total	С	Ν	0	S	0	1		
	A	525	3805	2397	622	766	20	0	1		
1	Р	525	Total	С	Ν	0	S	0	1		
	D	525	3805	2397	622	766	20	0	1		
1	C	525	Total	С	Ν	0	$\mathbf{S}$	0	1		
		525	3805	2397	622	766	20	0	1		
1	П	525	Total	С	Ν	0	$\mathbf{S}$	0	1		
1	D	525	3805	2397	622	766	20	0	T		
1	F	525	Total	С	Ν	0	$\mathbf{S}$	0	1		
1		525	3805	2397	622	766	20	0	L		
1	F	F	F	525	Total	С	Ν	0	$\mathbf{S}$	0	1
1	Ľ	525	3805	2397	622	766	20	0	1		
1	G	525	Total	С	Ν	Ο	$\mathbf{S}$	0	1		
L	I G	525	3805	2397	622	766	20	0	1		
1	н	523	Total	С	Ν	Ο	$\mathbf{S}$	0	1		
1	11	525	3793	2389	620	764	20	0	1		
1	т	523	Total	$\mathbf{C}$	Ν	Ο	$\mathbf{S}$	0	1		
1	T	525	3793	2389	620	764	20	0	T		
1	Т	523	Total	$\mathbf{C}$	Ν	Ο	$\mathbf{S}$	0	1		
	0	020	3793	2389	620	764	20	0	1		
1	K	523	Total	$\mathbf{C}$	Ν	Ο	$\mathbf{S}$	0	1		
1	11	525	3793	2389	620	764	20	0	T		
1	L	523	Total	$\mathbf{C}$	Ν	Ο	$\mathbf{S}$	0	1		
1		525	3793	2389	620	764	20	0	T		
1	M 5	523	Total	$\mathbf{C}$	Ν	Ο	$\mathbf{S}$	0	1		
		020	3793	2389	620	764	20	0	1		
1	N	523	Total	$\mathbf{C}$	Ν	Ο	$\mathbf{S}$	0	1		
	020	3793	2389	620	764	20	0	L			

• Molecule 1 is a protein called 60 KDA CHAPERONIN.

• Molecule 2 is a protein called 10 KDA CHAPERONIN MOLECULE: GROES, PROTEIN CPN10, GROES PROTEIN.



Mol	Chain	Residues		At	oms			AltConf	Trace
0	2 0	03	Total	С	Ν	0	S	0	1
	0	90	680	432	112	135	1	0	
9	P	03	Total	С	Ν	0	S	0	1
	1	90	680	432	112	135	1	0	1
2	0	93	Total	С	Ν	Ο	S	0	1
	2 Q		680	432	112	135	1	0	Ţ
2	0 D	93	Total	С	Ν	Ο	$\mathbf{S}$	0	1
2	10		680	432	112	135	1		
2	S	03	Total	С	Ν	Ο	$\mathbf{S}$	0	1
2	0	30	680	432	112	135	1	0	L
0	Т	03	Total	С	Ν	Ο	S	0	1
	1	90	680	432	112	135	1	0	1
0	9 U	03	Total	С	Ν	0	S	0	1
	U	90	680	432	112	135	1		



## 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: 60 KDA CHAPERONIN















 $\bullet$  Molecule 2: 10 KDA CHAPERONIN MOLECULE: GROES, PROTEIN CPN10, GROES PROTEIN



• Molecule 2: 10 KDA CHAPERONIN MOLECULE: GROES, PROTEIN CPN10, GROES PROTEIN





 $\bullet$  Molecule 2: 10 KDA CHAPERONIN MOLECULE: GROES, PROTEIN CPN10, GROES PROTEIN



 $\bullet$  Molecule 2: 10 KDA CHAPERONIN MOLECULE: GROES, PROTEIN CPN10, GROES PROTEIN



 $\bullet$  Molecule 2: 10 KDA CHAPERONIN MOLECULE: GROES, PROTEIN CPN10, GROES PROTEIN



 $\bullet$  Molecule 2: 10 KDA CHAPERONIN MOLECULE: GROES, PROTEIN CPN10, GROES PROTEIN



# 4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C7	Depositor
Number of particles used	7591	Depositor
Resolution determination method	Not provided	
CTF correction method	FULL CORRECTION ON 2D CLASS AV-	Depositor
	ERAGES	
Microscope	FEI TECNAI F20	Depositor
Voltage (kV)	200	Depositor
Electron dose $(e^-/\text{\AA}^2)$	15	Depositor
Minimum defocus (nm)	1400	Depositor
Maximum defocus (nm)	3300	Depositor
Magnification	50000	Depositor
Image detector	KODAK SO-163 FILM	Depositor
Maximum map value	7.869	Depositor
Minimum map value	-4.594	Depositor
Average map value	0.197	Depositor
Map value standard deviation	0.750	Depositor
Recommended contour level	1.9	Depositor
Map size (Å)	268.8, 268.8, 268.8	wwPDB
Map dimensions	192, 192, 192	wwPDB
Map angles (°)	90, 90, 90	wwPDB
Pixel spacing (Å)	1.4, 1.4, 1.4	Depositor



# 5 Model quality (i)

## 5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		Bond	lengths	Bond angles		
	Ullalli	RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	А	0.60	0/3833	0.91	0/5165	
1	В	0.60	0/3833	0.92	0/5165	
1	С	0.60	0/3833	0.91	0/5165	
1	D	0.60	0/3833	0.89	0/5165	
1	Е	0.60	0/3833	0.92	0/5165	
1	F	0.59	0/3833	0.91	0/5165	
1	G	0.60	0/3833	0.90	0/5165	
1	Н	0.60	0/3820	0.91	1/5146~(0.0%)	
1	Ι	0.60	0/3820	0.92	0/5146	
1	J	0.60	0/3820	0.90	0/5146	
1	Κ	0.60	0/3820	0.91	0/5146	
1	L	0.60	0/3820	0.92	0/5146	
1	М	0.61	0/3820	0.91	0/5146	
1	N	0.60	0/3820	0.92	2/5146~(0.0%)	
2	0	0.64	0/684	0.99	1/918~(0.1%)	
2	Р	0.63	0/684	1.01	0/918	
2	Q	0.64	0/684	1.01	1/918~(0.1%)	
2	R	0.64	0/684	1.03	0/918	
2	S	0.63	0/684	1.02	0/918	
2	Т	0.62	0/684	1.04	1/918~(0.1%)	
2	U	0.63	0/684	1.00	0/918	
All	All	0.60	0/58359	0.92	6/78603~(0.0%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	А	2	3
1	В	2	3
1	С	2	3
1	D	2	3



Mol	Chain	#Chirality outliers	#Planarity outliers
1	Ε	2	3
1	F	2	3
1	G	2	5
1	J	0	1
1	Κ	0	1
1	L	0	4
1	Ν	0	3
2	Р	0	1
2	Т	0	1
2	U	0	1
All	All	14	35

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms		$Observed(^{o})$	$Ideal(^{o})$
2	Т	50	GLU	C-N-CA	5.40	135.21	121.70
1	Ν	186	GLU	OE1-CD-OE2	-5.34	116.89	123.30
1	Н	404	ARG	CD-NE-CZ	5.18	130.86	123.60
1	Ν	506	TYR	CB-CG-CD2	-5.05	117.97	121.00
2	0	79	ASP	C-N-CA	5.05	134.32	121.70
2	Q	14	ARG	CD-NE-CZ	5.00	130.60	123.60

All (14) chirality outliers are listed below:

Mol	Chain	$\mathbf{Res}$	Type	Atom
1	А	136	VAL	CA
1	А	137	PRO	CA
1	В	136	VAL	CA
1	В	137	PRO	CA
1	С	136	VAL	CA
1	С	137	PRO	CA
1	D	136	VAL	CA
1	D	137	PRO	CA
1	Е	136	VAL	CA
1	Е	137	PRO	CA
1	F	136	VAL	CA
1	F	137	PRO	CA
1	G	136	VAL	CA
1	G	137	PRO	CA

All (35) planarity outliers are listed below:



Mol	Chain	Res	Type	Group
1	А	136	VAL	Peptide
1	А	4	LYS	Peptide
1	А	8	PHE	Peptide
1	В	136	VAL	Peptide
1	В	218	PRO	Peptide
1	В	4	LYS	Peptide
1	С	136	VAL	Peptide
1	С	4	LYS	Peptide
1	С	8	PHE	Peptide
1	D	136	VAL	Peptide
1	D	195	PHE	Peptide
1	D	4	LYS	Peptide
1	Е	136	VAL	Peptide
1	Е	284	ARG	Peptide
1	Е	4	LYS	Peptide
1	F	136	VAL	Peptide
1	F	172	GLU	Peptide
1	F	4	LYS	Peptide
1	G	136	VAL	Peptide
1	G	139	SER	Peptide
1	G	218	PRO	Peptide
1	G	307	MET	Peptide
1	G	4	LYS	Peptide
1	J	355	GLU	Peptide
1	K	136	VAL	Peptide
1	L	139	SER	Peptide
1	L	178	GLU	Peptide
1	L	218	PRO	Peptide
1	L	4	LYS	Peptide
1	Ν	218	PRO	Peptide
1	N	4	LYS	Peptide
1	N	63	GLU	Peptide
2	Р	19	THR	Peptide
2	Т	93	ALA	Peptide
2	U	93	ALA	Peptide

## 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	3805	0	3881	1	0
1	В	3805	0	3881	2	0
1	С	3805	0	3881	1	0
1	D	3805	0	3881	3	0
1	Е	3805	0	3881	3	0
1	F	3805	0	3881	2	0
1	G	3805	0	3881	3	0
1	Н	3793	0	3869	2	0
1	Ι	3793	0	3869	1	0
1	J	3793	0	3869	1	0
1	Κ	3793	0	3869	3	0
1	L	3793	0	3869	2	0
1	М	3793	0	3869	2	0
1	Ν	3793	0	3869	1	0
2	0	680	0	703	2	0
2	Р	680	0	703	1	0
2	Q	680	0	703	1	0
2	R	680	0	703	3	0
2	S	680	0	703	3	0
2	Т	680	0	703	3	0
2	U	680	0	703	2	0
All	All	57946	0	59171	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:J:314:LEU:H	1:J:314:LEU:HD12	1.69	0.56
2:R:3:ILE:HA	2:S:95:VAL:N	2.23	0.53
1:D:385:THR:HG21	1:E:510:VAL:HG23	1.92	0.52
1:B:387:VAL:HA	1:B:390:LYS:HE3	1.91	0.52
1:E:7:LYS:HE2	1:E:66:PHE:CE2	2.45	0.52
1:F:385:THR:HG21	1:G:510:VAL:HG23	1.92	0.50
1:G:7:LYS:HE2	1:G:66:PHE:CE2	2.47	0.50
1:E:7:LYS:HE2	1:E:66:PHE:CD2	2.47	0.49
1:I:7:LYS:HE2	1:I:66:PHE:CE2	2.47	0.49
1:M:149:THR:HG22	1:M:156:GLU:HA	1.95	0.48
1:K:190:VAL:HG22	1:K:191:GLU:H	1.79	0.47
1:N:489:ILE:HD13	1:N:494:LEU:HD21	1.95	0.47
1:B:7:LYS:HE2	1:B:66:PHE:CE2	2.49	0.47



EMD-1181,	2C7D
-----------	------

Atom 1	Atom 2	Interatomic	Clash			
Atom-1	Atom-2	distance (Å)	overlap (Å)			
1:H:7:LYS:HE2	1:H:66:PHE:CE2	2.49	0.47			
1:A:7:LYS:HE2	1:A:66:PHE:CE2	2.50	0.47			
2:O:60:LYS:N	2:O:60:LYS:HE3	2.31	0.46			
1:M:314:LEU:H	1:M:314:LEU:HD12	1.80	0.46			
2:Q:3:ILE:HA	2:R:94:ILE:O	2.17	0.45			
1:F:7:LYS:HE2	1:F:66:PHE:CE2	2.51	0.45			
2:T:3:ILE:HA	2:U:95:VAL:N	2.31	0.44			
1:L:7:LYS:HE2	1:L:66:PHE:CE2	2.51	0.44			
2:T:3:ILE:HA	2:U:94:ILE:O	2.16	0.44			
2:R:3:ILE:HA	2:S:94:ILE:O	2.18	0.43			
1:G:146:GLN:HE21	1:G:494:LEU:HD21	1.82	0.43			
1:D:7:LYS:HE2	1:D:66:PHE:CE2	2.55	0.42			
1:K:7:LYS:HE2	1:K:66:PHE:CE2	2.54	0.42			
2:O:3:ILE:HA	2:P:95:VAL:N	2.33	0.42			
1:H:190:VAL:HG22	1:H:191:GLU:H	1.85	0.42			
1:L:24:ALA:HB3	1:L:97:GLN:HE21	1.85	0.42			
1:C:7:LYS:HE2	1:C:66:PHE:CE2	2.56	0.41			
2:S:3:ILE:HA	2:T:95:VAL:N	2.36	0.41			
1:K:149:THR:HG22	1:K:156:GLU:HA	2.03	0.40			
1:D:57:ALA:O	1:D:75:LYS:HE2	2.21	0.40			

There are no symmetry-related clashes.

## 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	А	523/547~(96%)	488 (93%)	28~(5%)	7 (1%)	12	48
1	В	523/547~(96%)	490 (94%)	25~(5%)	8 (2%)	10	46
1	С	523/547~(96%)	486 (93%)	30 (6%)	7 (1%)	12	48
1	D	523/547~(96%)	491 (94%)	28 (5%)	4 (1%)	19	60



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	Е	523/547~(96%)	483 (92%)	32~(6%)	8 (2%)	10	46
1	F	523/547~(96%)	485 (93%)	30 (6%)	8 (2%)	10	46
1	G	523/547~(96%)	488 (93%)	31 (6%)	4 (1%)	19	60
1	Н	521/547~(95%)	476 (91%)	38 (7%)	7 (1%)	12	48
1	Ι	521/547~(95%)	486 (93%)	29 (6%)	6 (1%)	13	50
1	J	521/547~(95%)	489 (94%)	23 (4%)	9 (2%)	9	42
1	К	521/547~(95%)	477 (92%)	33 (6%)	11 (2%)	7	36
1	L	521/547~(95%)	473 (91%)	39 (8%)	9(2%)	9	42
1	М	521/547~(95%)	486 (93%)	29 (6%)	6 (1%)	13	50
1	Ν	521/547~(95%)	478 (92%)	36 (7%)	7 (1%)	12	48
2	Ο	91/97~(94%)	65 (71%)	21 (23%)	5~(6%)	2	19
2	Р	91/97~(94%)	70 (77%)	16 (18%)	5~(6%)	2	19
2	Q	91/97~(94%)	70 (77%)	16 (18%)	5~(6%)	2	19
2	R	91/97~(94%)	70 (77%)	18 (20%)	3(3%)	4	26
2	S	91/97~(94%)	70 (77%)	15 (16%)	6 (7%)	1	16
2	Т	91/97~(94%)	71 (78%)	15 (16%)	5 (6%)	2	19
2	U	91/97~(94%)	70 (77%)	15 (16%)	6 (7%)	1	16
All	All	7945/8337~(95%)	7262 (91%)	547 (7%)	136 (2%)	13	42

All (136) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	А	137	PRO
1	В	137	PRO
1	С	137	PRO
1	С	483	GLU
1	D	137	PRO
1	Е	137	PRO
1	Е	197	ARG
1	Е	285	ARG
1	F	137	PRO
1	G	137	PRO
1	Н	217	SER
1	Н	524	LEU
1	Ι	154	SER
1	Ι	217	SER



Mol	Chain	Res	Type
1	J	217	SER
1	K	154	SER
1	К	217	SER
1	К	334	ASP
1	L	154	SER
1	М	154	SER
1	Ν	137	PRO
1	Ν	483	GLU
2	0	80	ASN
2	Р	32	ALA
2	Р	80	ASN
2	Q	21	SER
2	R	32	ALA
2	R	51	ASN
2	S	73	VAL
2	Т	16	GLU
2	Т	35	SER
2	Т	51	ASN
2	U	73	VAL
1	А	154	SER
1	В	153	ASN
1	В	310	GLU
1	С	9	GLY
1	D	334	ASP
1	Е	139	SER
1	Е	483	GLU
1	Ι	139	SER
1	Ι	483	GLU
1	J	154	SER
1	J	268	ARG
1	J	334	ASP
1	K	210	THR
1	L	217	SER
1	L	233	MET
1	L	409	GLU
1	L	483	GLU
1	М	217	SER
1	N	63	GLU
1	N	$15\overline{4}$	SER
2	Q	80	ASN
2	Q	94	ILE
2	S	7	HIS



Mol	Chain	Res	Type
2	S	18	GLU
2	Т	7	HIS
2	U	94	ILE
1	А	153	ASN
1	В	373	ALA
1	D	153	ASN
1	F	139	SER
1	F	334	ASP
1	G	174	VAL
1	Ι	43	SER
1	J	483	GLU
1	K	139	SER
1	Κ	409	GLU
1	М	139	SER
2	0	7	HIS
2	0	18	GLU
2	Р	7	HIS
2	Q	18	GLU
2	S	35	SER
2	U	35	SER
1	А	45	GLY
1	А	525	PRO
1	В	483	GLU
1	С	10	ASN
1	D	217	SER
1	Е	336	VAL
1	F	153	ASN
1	F	196	ASP
1	G	154	SER
1	Н	139	SER
1	J	210	THR
1	J	474	GLY
1	K	190	VAL
1	K	281	PHE
1	L	45	GLY
1	N	62	LEU
1	N	217	SER
2	Ο	8	ASP
2	Р	94	ILE
2	Q	7	HIS
2	R	7	HIS
2	S	8	ASP



Mol	Chain	Res	Type
2	Т	94	ILE
2	U	7	HIS
2	U	30	SER
1	А	310	GLU
1	В	231	ARG
1	С	217	SER
1	С	373	ALA
1	Е	217	SER
1	F	217	SER
1	Н	29	VAL
1	Н	327	LYS
1	K	136	VAL
1	L	357	THR
1	М	209	GLU
1	Ν	209	GLU
2	U	32	ALA
1	В	525	PRO
1	Е	337	GLY
1	F	395	ARG
1	G	483	GLU
1	Н	210	THR
1	J	45	GLY
1	L	169	VAL
1	М	210	THR
1	А	205	ILE
1	F	45	GLY
1	Н	137	PRO
1	K	336	VAL
2	0	94	ILE
1	K	137	PRO
1	L	336	VAL
1	С	205	ILE
1	J	190	VAL
2	Р	48	ILE
1	В	205	ILE
1	Ι	137	PRO
2	S	72	GLY
1	М	137	PRO



#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the side chain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perc	entiles
1	А	404/414~(98%)	386~(96%)	18 (4%)	27	52
1	В	404/414~(98%)	386~(96%)	18 (4%)	27	52
1	С	404/414~(98%)	392~(97%)	12 (3%)	41	63
1	D	404/414~(98%)	389~(96%)	15~(4%)	34	58
1	Ε	404/414~(98%)	383~(95%)	21 (5%)	23	48
1	F	404/414 (98%)	390 (96%)	14 (4%)	36	59
1	G	404/414~(98%)	391~(97%)	13 (3%)	39	61
1	Н	403/414~(97%)	387~(96%)	16 (4%)	31	55
1	Ι	403/414 (97%)	387~(96%)	16 (4%)	31	55
1	J	403/414~(97%)	387~(96%)	16 (4%)	31	55
1	Κ	403/414 (97%)	386 (96%)	17 (4%)	30	54
1	L	403/414 (97%)	385~(96%)	18 (4%)	27	52
1	М	403/414 (97%)	385~(96%)	18 (4%)	27	52
1	Ν	403/414~(97%)	384 (95%)	19 (5%)	26	51
2	Ο	76/80~(95%)	69 (91%)	7 (9%)	9	29
2	Р	76/80~(95%)	69~(91%)	7 (9%)	9	29
2	Q	76/80~(95%)	68~(90%)	8 (10%)	7	24
2	R	76/80~(95%)	65~(86%)	11 (14%)	3	15
2	S	76/80~(95%)	66 (87%)	10 (13%)	4	18
2	Т	76/80~(95%)	70 (92%)	6 (8%)	12	35
2	U	76/80~(95%)	65~(86%)	11 (14%)	3	15
All	All	6181/6356~(97%)	5890 (95%)	291 (5%)	30	51

All (291) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	11	ASP



Mol	Chain	Res	Type
1	А	52	ASP
1	А	74	VAL
1	А	97	GLN
1	А	111	MET
1	А	136	VAL
1	А	157	THR
1	А	168	LYS
1	А	184	GLN
1	А	197	ARG
1	А	225	LYS
1	А	232	GLU
1	А	237	LEU
1	А	257	GLU
1	А	345	ARG
1	А	351	GLN
1	А	391	GLU
1	А	400	LEU
1	В	11	ASP
1	В	18	ARG
1	В	111	MET
1	В	136	VAL
1	В	157	THR
1	В	168	LYS
1	В	179	ASP
1	В	184	GLN
1	В	197	ARG
1	В	206	ASN
1	В	217	SER
1	В	225	LYS
1	В	228	SER
1	В	237	LEU
1	В	257	GLU
1	В	268	ARG
1	В	345	ARG
1	В	391	GLU
1	С	23	LEU
1	С	25	ASP
1	С	97	GLN
1	С	136	VAL
1	С	157	THR
1	С	184	GLN
1	С	206	ASN



Mol	Chain	Res	Type
1	С	225	LYS
1	С	257	GLU
1	С	268	ARG
1	С	345	ARG
1	С	391	GLU
1	D	23	LEU
1	D	52	ASP
1	D	111	MET
1	D	136	VAL
1	D	157	THR
1	D	176	THR
1	D	184	GLN
1	D	204	PHE
1	D	225	LYS
1	D	237	LEU
1	D	257	GLU
1	D	268	ARG
1	D	345	ARG
1	D	386	GLU
1	D	397	GLU
1	Е	23	LEU
1	Е	74	VAL
1	Е	77	VAL
1	Е	111	MET
1	Е	136	VAL
1	Е	157	THR
1	Ε	179	ASP
1	Ε	184	GLN
1	Е	197	ARG
1	Ε	204	PHE
1	Ε	206	ASN
1	E	225	LYS
1	E	228	SER
1	E	237	LEU
1	E	257	GLU
1	E	268	ARG
1	Е	345	ARG
1	Е	376	VAL
1	E	391	GLU
1	Е	463	SER
1	E	473	ASP
1	F	11	ASP



Mol	Chain	Res	Type
1	F	44	PHE
1	F	52	ASP
1	F	97	GLN
1	F	111	MET
1	F	136	VAL
1	F	197	ARG
1	F	225	LYS
1	F	228	SER
1	F	237	LEU
1	F	257	GLU
1	F	345	ARG
1	F	417	VAL
1	F	494	LEU
1	G	111	MET
1	G	136	VAL
1	G	157	THR
1	G	168	LYS
1	G	179	ASP
1	G	184	GLN
1	G	206	ASN
1	G	237	LEU
1	G	257	GLU
1	G	268	ARG
1	G	345	ARG
1	G	391	GLU
1	G	494	LEU
1	Н	52	ASP
1	Н	77	VAL
1	Н	111	MET
1	Н	132	LYS
1	Н	153	ASN
1	Н	196	ASP
1	Н	233	MET
1	Н	255	GLU
1	Н	268	ARG
1	H	313	THR
1	H	328	ASP
1	Н	329	THR
1	Η	389	MET
1	H	398	ASP
1	Н	404	ARG
1	Н	495	ASP



Mol	Chain	Res	Type
1	Ι	59	GLU
1	Ι	74	VAL
1	Ι	77	VAL
1	Ι	111	MET
1	Ι	132	LYS
1	Ι	169	VAL
1	Ι	187	LEU
1	Ι	230	ILE
1	Ι	233	MET
1	Ι	268	ARG
1	Ι	284	ARG
1	Ι	308	GLU
1	Ι	328	ASP
1	Ι	357	THR
1	Ι	391	GLU
1	Ι	495	ASP
1	J	25	ASP
1	J	54	VAL
1	J	87	ASP
1	J	97	GLN
1	J	111	MET
1	J	196	ASP
1	J	228	SER
1	J	233	MET
1	J	268	ARG
1	J	284	ARG
1	J	338	GLU
1	J	357	THR
1	J	398	ASP
1	J	428	ASP
1	J	495	ASP
1	J	499	VAL
1	Κ	43	SER
1	K	52	ASP
1	K	77	VAL
1	K	97	GLN
1	Κ	111	MET
1	K	140	ASP
1	K	172	GLU
1	Κ	230	ILE
1	K	233	MET
1	Κ	268	ARG



Mol	Chain	Res	Type
1	K	272	LYS
1	K	284	ARG
1	К	325	ILE
1	K	326	ASN
1	К	364	LYS
1	K	389	MET
1	K	398	ASP
1	L	25	ASP
1	L	52	ASP
1	L	77	VAL
1	L	97	GLN
1	L	111	MET
1	L	132	LYS
1	L	153	ASN
1	L	172	GLU
1	L	197	ARG
1	L	233	MET
1	L	265	ASN
1	L	326	ASN
1	L	331	THR
1	L	357	THR
1	L	364	LYS
1	L	422	VAL
1	L	424	SER
1	L	499	VAL
1	М	23	LEU
1	М	52	ASP
1	М	74	VAL
1	М	77	VAL
1	М	139	SER
1	М	146	GLN
1	М	161	LEU
1	М	169	VAL
1	М	233	MET
1	Μ	283	ASP
1	М	284	ARG
1	М	308	GLU
1	М	355	GLU
1	М	357	THR
1	М	359	ASP
1	М	398	ASP
1	М	428	ASP



Mol	Chain	Res	Type
1	М	495	ASP
1	Ν	52	ASP
1	N	79	SER
1	Ν	97	GLN
1	Ν	111	MET
1	Ν	121	ASP
1	Ν	132	LYS
1	Ν	137	PRO
1	N	172	GLU
1	N	197	ARG
1	N	233	MET
1	N	253	ASP
1	Ν	268	ARG
1	Ν	284	ARG
1	N	308	GLU
1	N	357	THR
1	N	389	MET
1	Ν	398	ASP
1	N	422	VAL
1	Ν	473	ASP
2	0	6	LEU
2	0	8	ASP
2	0	30	SER
2	0	51	ASN
2	0	55	LYS
2	0	60	LYS
2	0	87	SER
2	Р	6	LEU
2	Р	20	LYS
2	Р	55	LYS
2	P	60	LYS
2	P	86	MET
2	Р	87	SER
2	P	94	ILE
2	Q	6	LEU
2	Q	16	GLU
2	Q	37	ARG
2	Q	55	LYS
2	Q	60	LYS
2	Q	68	ASN
2	Q	86	MET
2	Q	94	ILE



Mol	Chain	Res	Type
2	R	6	LEU
2	R	9	ARG
2	R	20	LYS
2	R	34	LYS
2	R	51	ASN
2	R	55	LYS
2	R	58	ASP
2	R	60	LYS
2	R	81	GLU
2	R	86	MET
2	R	87	SER
2	S	6	LEU
2	S	20	LYS
2	S	30	SER
2	S	37	ARG
2	S	58	ASP
2	S	60	LYS
2	S	68	ASN
2	S	81	GLU
2	S	86	MET
2	S	94	ILE
2	Т	6	LEU
2	Т	20	LYS
2	Т	30	SER
2	Т	51	ASN
2	Т	55	LYS
2	Т	60	LYS
2	U	6	LEU
2	U	9	ARG
2	U	15	LYS
2	U	20	LYS
2	U	30	SER
2	U	34	LYS
2	U	51	ASN
2	U	55	LYS
2	U	60	LYS
2	U	68	ASN
2	U	86	MET

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such side chains are listed below:



Mol	Chain	Res	Type
1	С	184	GLN
1	Ι	194	GLN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry (i)

There are no ligands in this entry.

## 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-1181. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

## 6.1 Orthogonal projections (i)

#### 6.1.1 Primary map



The images above show the map projected in three orthogonal directions.

## 6.2 Central slices (i)

#### 6.2.1 Primary map



X Index: 96



Y Index: 96



Z Index: 96



The images above show central slices of the map in three orthogonal directions.

## 6.3 Largest variance slices (i)

#### 6.3.1 Primary map



X Index: 73

Y Index: 118

Z Index: 89

The images above show the largest variance slices of the map in three orthogonal directions.

## 6.4 Orthogonal surface views (i)

#### 6.4.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 1.9. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.



## 6.5 Mask visualisation (i)

This section was not generated. No masks/segmentation were deposited.



## 7 Map analysis (i)

This section contains the results of statistical analysis of the map.

## 7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



## 7.2 Volume estimate (i)



The volume at the recommended contour level is 965  $\rm nm^3;$  this corresponds to an approximate mass of 872 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



## 7.3 Rotationally averaged power spectrum (i)



\*Reported resolution corresponds to spatial frequency of 0.115  ${\rm \AA^{-1}}$ 



# 8 Fourier-Shell correlation (i)

This section was not generated. No FSC curve or half-maps provided.



## 9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-1181 and PDB model 2C7D. Per-residue inclusion information can be found in section 3 on page 6.

## 9.1 Map-model overlay (i)



The images above show the 3D surface view of the map at the recommended contour level 1.9 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.



## 9.2 Q-score mapped to coordinate model (i)



The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

#### 9.3 Atom inclusion mapped to coordinate model (i)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (1.9).



## 9.4 Atom inclusion (i)



At the recommended contour level, 85% of all backbone atoms, 74% of all non-hydrogen atoms, are inside the map.



1.0

0.0 <0.0

## 9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (1.9) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	$\mathbf{Q} ext{-score}$
All	0.7434	0.0780
А	0.7745	0.0840
В	0.7735	0.0840
С	0.7900	0.0870
D	0.7806	0.0870
E	0.7926	0.0910
F	0.7827	0.0880
G	0.7827	0.0850
Η	0.6855	0.0620
Ι	0.6910	0.0710
J	0.7008	0.0670
Κ	0.7045	0.0640
L	0.7013	0.0650
М	0.6950	0.0670
Ν	0.6968	0.0710
О	0.7809	0.0910
Р	0.7882	0.1000
Q	0.8015	0.0990
R	0.7971	0.0950
S	0.7809	0.0950
Т	0.7868	0.0930
U	0.7809	0.0930

