Full wwPDB X-ray Structure Validation Report (i)

Dec 4, 2023 – 04:54 am GMT

PDB ID	:	2C7O
Title	:	HhaI DNA methyltransferase complex with 13mer oligonucleotide containing 2-aminopurine adjacent to the target base (PCGC:GMGC) and SAH
Authors	:	Daujotyte, D.; Grazulis, S.
Deposited on	:	2005-11-25
Resolution	:	1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	6847 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	327	86% 14%
2	C	13	69% 23% 8%
3	D	13	38% 46% 8% 8%

2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 3520 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MODIFICATION METHYLASE HHAI.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	327	2624	1677	443	491	13	0	4	0

- Molecule 2 is a DNA chain called 5'-D(*T*GP*GP*AP*TP*GP*(5CM)*GP*CP*TP*GP*AP *C)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	C	13	267	128	50	77	12	0	0	0

- Molecule 3 is a DNA chain called 5'-D(*T*GP*TP*CP*AP*(2PR)*CP*GP*CP*AP*TP*CP *C)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
3	D	12	242	115	44	71	12	0	0	0

- Molecule 4 is S-ADENOSYL-L-HOMOCYSTEINE (three-letter code: SAH) (formula: $C_{14}H_{20}N_6O_5S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
4	A	1	26	14	6	5	1	0	0

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	O	S			
5	A	1	5	4	1		0	0
5	A	1	Total 5	O 4	S 1		0	0
5	A	1	Total 5	O 4	S 1		0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
5	A	1	5	4	1	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
6	A	295	295	295	0	0
			Total	O		
6	C	28	28	28	0	0
			Total	O		
6	D	18	18	18	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: MODIFICATION METHYLASE HHAI

Chain A: 86% 14%



• Molecule 2: 5'-D(*T*GP*GP*AP*TP*GP*(5CM)*GP*CP*TP*GP*AP *C)-3'

Chain C: 69% 23% 8%

• Molecule 3: 5'-D(*T*GP*TP*CP*AP*(2PR)*CP*GP*CP*AP*TP*CP *C)-3'

Chain D: 38% 46% 8% 8%

4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α, β, γ	95.18Å 95.18Å 312.55Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	23.08 – 1.90	Depositor
% Data completeness (in resolution range)	99.3 (23.08-1.90)	Depositor
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.194 , 0.215	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3520	wwPDB-VP
Average B, all atoms (Å^2)	18.0	wwPDB-VP

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, 5CM, SAH, 2PR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.35	0/2679	0.60	1/3615 (0.0%)
2	C	0.36	0/276	0.73	0/423
3	D	0.34	0/245	0.87	0/372
All	All	0.35	0/3200	0.64	1/4410 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	D	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	78	GLY	N-CA-C	-6.16	97.71	113.10

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	D	432	DC	Sidechain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2624	0	2584	38	0
2	C	267	0	150	5	0
3	D	242	0	135	10	0
4	A	26	0	19	0	0
5	A	20	0	0	0	0
6	A	295	0	0	4	0
6	C	28	0	0	0	0
6	D	18	0	0	0	0
All	All	3520	0	2888	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:275:HIS:HD2	1:A:277:ARG:H	1.25	0.83
3:D:423:DT:H2"	3:D:424:DC:H5'	1.59	0.82
2:C:408:DG:H2"	2:C:409:DC:C5	2.19	0.78
1:A:81:CYS:SG	3:D:427:DC:C6	2.80	0.73
1:A:82:GLN:H	1:A:82:GLN:HE21	1.40	0.70
1:A:256:GLY:HA3	2:C:406:DG:O6	1.93	0.68
1:A:275:HIS:CD2	1:A:277:ARG:H	2.10	0.68
3:D:423:DT:H2"	3:D:424:DC:C5'	2.23	0.68
1:A:127:HIS:HD2	1:A:132[A]:THR:OG1	1.80	0.64
1:A:221:GLN:HG3	6:A:2173:HOH:O	1.97	0.63
1:A:42:ASP:O	1:A:46:GLN:HG3	2.03	0.58
1:A:127:HIS:HD2	1:A:132[B]:THR:HG21	1.69	0.58
1:A:25:ARG:HH12	1:A:55:GLU:CD	2.08	0.57
1:A:169:ILE:HG13	1:A:180:PHE:CZ	2.40	0.56
1:A:81:CYS:SG	3:D:427:DC:H2'	2.46	0.56
1:A:306:VAL:HG21	1:A:311[B]:LEU:HD11	1.86	0.56
3:D:425:DA:H2"	3:D:426:2PR:C8	2.37	0.54
1:A:93:PHE:CE2	1:A:132[B]:THR:HG22	2.43	0.53
3:D:425:DA:H2"	3:D:426:2PR:H8	1.91	0.52
1:A:156:ASP:HB3	1:A:185:PRO:HB3	1.92	0.52
1:A:81:CYS:HG	3:D:427:DC:H2'	1.75	0.52
2:C:407:5CM:H2"	2:C:408:DG:C8	2.45	0.52
1:A:127:HIS:HD2	1:A:132[B]:THR:CG2	2.24	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:408:DG:H2"	2:C:409:DC:C6	2.46	0.50
1:A:137:LYS:HE2	6:A:2107:HOH:O	2.10	0.50
1:A:297:GLN:O	1:A:301:GLN:HG3	2.13	0.49
1:A:155:LEU:HD12	1:A:155:LEU:C	2.33	0.49
1:A:15:ILE:HG22	1:A:37:TYR:HB3	1.95	0.48
1:A:125:ALA:HB2	1:A:166:ILE:HD12	1.94	0.48
1:A:132[B]:THR:HG23	6:A:2094:HOH:O	2.14	0.47
1:A:209:ARG:HB3	1:A:211:ASP:OD1	2.15	0.46
3:D:422:DG:C2'	3:D:423:DT:H71	2.46	0.46
2:C:406:DG:H1'	2:C:407:5CM:H5"	1.99	0.44
1:A:82:GLN:HE21	1:A:82:GLN:N	2.11	0.44
1:A:82:GLN:H	1:A:82:GLN:NE2	2.10	0.44
1:A:79:PHE:HB2	1:A:80:PRO:HD2	1.99	0.44
1:A:70:PRO:O	1:A:72:HIS:HD2	2.01	0.43
3:D:422:DG:H2"	3:D:423:DT:OP2	2.18	0.43
1:A:61:ILE:HD11	1:A:107:ILE:HD11	2.01	0.43
1:A:127:HIS:CD2	1:A:132[B]:THR:HG21	2.51	0.42
1:A:279:CYS:O	1:A:282[B]:VAL:HG22	2.20	0.41
3:D:424:DC:H2"	3:D:425:DA:C8	2.55	0.41
1:A:279:CYS:HA	1:A:282[B]:VAL:HG22	2.03	0.41
1:A:79:PHE:HB2	1:A:80:PRO:CD	2.51	0.41
1:A:251:LEU:HD11	1:A:282[B]:VAL:CG2	2.50	0.41
1:A:65:ASN:HB3	1:A:68:THR:HG23	2.03	0.41
1:A:21:LEU:HD23	1:A:48:VAL:HG11	2.03	0.41
1:A:269:GLY:N	6:A:2210:HOH:O	2.33	0.40
1:A:101:PHE:O	1:A:104:ILE:HB	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

There are no protein backbone outliers to report in this entry.

5.3.2 Protein sidechains (i)

There are no protein residues with a non-rotameric sidechain to report in this entry.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
3	2PR	D	426	2,3	17,23,24	1.12	2 (11%)	16,33,36	1.47	2 (12%)
2	5CM	C	407	2,3	17,21,22	0.74	1 (5%)	24,30,33	1.63	4 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	2PR	D	426	2,3	-	0/3/21/22	0/3/3/3
2	5CM	C	407	2,3	-	0/7/21/22	0/2/2/2

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	426	2PR	C6-N1	3.42	1.39	1.32
2	C	407	5CM	C4-N3	2.37	1.38	1.34
3	D	426	2PR	C8-N7	-2.01	1.31	1.34

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	C	407	5CM	C5-C4-N3	-5.27	115.98	121.67
3	D	426	2PR	C6-N1-C2	4.64	125.01	116.43
2	C	407	5CM	C4-N3-C2	4.58	126.88	120.69
3	D	426	2PR	C2-N3-C4	-2.84	112.11	115.36

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	C	407	5CM	N1-C2-N3	-2.30	114.62	118.81
2	C	407	5CM	O2-C2-N1	2.07	123.17	118.89

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	426	2PR	2	0
2	C	407	5CM	2	0

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
5	SO4	A	1330	-	4,4,4	1.84	2 (50%)	6,6,6	0.94	0
4	SAH	A	1328	-	24,28,28	1.34	3 (12%)	25,40,40	2.37	4 (16%)
5	SO4	A	1332	-	4,4,4	1.86	2 (50%)	6,6,6	0.91	0
5	SO4	A	1329	-	4,4,4	1.83	2 (50%)	6,6,6	0.90	0
5	SO4	A	1331	-	4,4,4	1.88	2 (50%)	6,6,6	0.88	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	SAH	A	1328	-	-	3/11/31/31	0/3/3/3

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1328	SAH	C4-N3	3.45	1.40	1.35
4	A	1328	SAH	C2'-C1'	-3.42	1.48	1.53
5	A	1331	SO4	O1-S	3.09	1.62	1.46
5	A	1332	SO4	O1-S	3.06	1.62	1.46
5	A	1330	SO4	O1-S	3.06	1.62	1.46
5	A	1329	SO4	O1-S	3.00	1.62	1.46
4	A	1328	SAH	C8-N7	-2.45	1.30	1.34
5	A	1331	SO4	O3-S	-2.09	1.30	1.47
5	A	1332	SO4	O3-S	-2.07	1.30	1.47
5	A	1329	SO4	O3-S	-2.04	1.31	1.47
5	A	1330	SO4	O3-S	-2.01	1.31	1.47

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
4	A	1328	SAH	OXT-C-O	-9.19	103.22	124.09
4	A	1328	SAH	O4'-C1'-C2'	-4.98	99.66	106.93
4	A	1328	SAH	CB-CA-N	2.87	117.68	110.17
4	A	1328	SAH	C5-C6-N6	2.83	124.66	120.35

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1328	SAH	OXT-C-CA-CB
4	A	1328	SAH	OXT-C-CA-N
4	A	1328	SAH	CB-CG-SD-C5'

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.

Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand SAH A 1328



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.