

Full wwPDB X-ray Structure Validation Report (i)

Mar 4, 2024 – 04:13 AM EST

PDB ID	:	1C81
Title	:	MICHAELIS COMPLEX OF FRUCTOSE-2,6-BISPHOSPHATASE
Authors	:	Lee, Y.H.; Olson, T.W.; McClard, R.W.; Witte, J.F.; McFarlan, S.C.; Ba-
		naszak, L.J.; Levitt, D.G.; Lange, A.J.
Deposited on	:	2000-04-03
Resolution	:	2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain	
1	А	191	70%	30% •



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 1822 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


• Molecule 1 is a protein called FRUCTOSE-2,6-BISPHOSPHATASE.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	А	191	Total 1555	C 984	N 268	O 292	S 11	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	250	MET	-	initiating methionine	UNP P07953

• Molecule 2 is 2,5-anhydro-1-deoxy-1-phosphono-6-O-phosphono-D-glucitol (three-letter code: FDQ) (formula: $C_6H_{14}O_{10}P_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	А	1	Total 18	С 6	O 10	Р 2	0	0

• Molecule 3 is water.



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	А	249	Total 249	O 249	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: FRUCTOSE-2,6-BISPHOSPHATASE





4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	80.10Å 51.60Å 49.30Å	Deperitor
a, b, c, α , β , γ	90.00° 95.70° 90.00°	Depositor
Resolution (Å)	18.00 - 2.50	Depositor
% Data completeness	79.1 (18.00-2.50)	Depositor
(in resolution range)	75.1 (10.00-2.50)	Depositor
R_{merge}	0.04	Depositor
R _{sym}	(Not available)	Depositor
Refinement program	X-PLOR 3.1	Depositor
R, R_{free}	0.209 , 0.287	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	1822	wwPDB-VP
Average B, all atoms $(Å^2)$	22.0	wwPDB-VP



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: FDQ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain		lengths	Bond angles		
IVI01	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	0.36	0/1587	0.59	0/2142	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	1555	0	1535	45	2
2	А	18	0	10	3	0
3	А	249	0	0	14	2
All	All	1822	0	1545	45	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)	
1:A:278:ARG:CZ	3:A:614:HOH:O	2.20	0.87	

Continued on next page...



Continued from prev		Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:A:353:ASP:CG	3:A:639:HOH:O	2.17	0.82
1:A:325:LEU:HD21	1:A:377:VAL:HG11	1.63	0.81
1:A:353:ASP:OD1	3:A:639:HOH:O	2.06	0.73
1:A:251:ARG:HH12	1:A:253:ILE:HD11	1.55	0.70
1:A:381:LEU:HD22	3:A:684:HOH:O	1.90	0.69
1:A:251:ARG:NH1	1:A:253:ILE:HD11	2.08	0.67
1:A:278:ARG:NH1	3:A:614:HOH:O	2.25	0.67
1:A:367:TYR:O	1:A:371:VAL:HG23	1.97	0.65
1:A:421:VAL:HG13	1:A:436:ILE:HB	1.83	0.60
1:A:371:VAL:HG22	3:A:478:HOH:O	2.02	0.60
1:A:338:TYR:HD2	1:A:338:TYR:H	1.48	0.59
1:A:286:LEU:O	1:A:290:ILE:HG12	2.04	0.57
1:A:253:ILE:HG13	3:A:680:HOH:O	2.04	0.57
1:A:289:PHE:CE1	1:A:436:ILE:HG21	2.40	0.56
1:A:266:ARG:HB3	3:A:600:HOH:O	2.07	0.54
1:A:305:MET:HG3	1:A:392:HIS:CE1	2.46	0.51
1:A:397:ARG:HH12	2:A:441:FDQ:P2	2.37	0.48
1:A:338:TYR:CD2	1:A:338:TYR:N	2.81	0.48
1:A:411:LEU:HB3	1:A:412:PRO:HD3	1.95	0.48
1:A:410:GLU:HB3	3:A:624:HOH:O	2.15	0.47
1:A:255:LEU:HD23	1:A:389:VAL:HB	1.97	0.47
1:A:328:ILE:HG13	1:A:365:GLU:HG3	1.97	0.46
1:A:303:SER:HA	1:A:322:TRP:O	2.16	0.45
1:A:305:MET:HB3	3:A:469:HOH:O	2.17	0.45
1:A:275:LEU:HD21	1:A:307:ARG:O	2.16	0.44
1:A:284:TYR:HE1	3:A:689:HOH:O	1.99	0.44
1:A:365:GLU:HB2	1:A:369:ASP:HB2	1.98	0.44
1:A:322:TRP:HB2	1:A:325:LEU:HD12	2.00	0.43
1:A:368:GLU:HG3	3:A:584:HOH:O	2.17	0.43
1:A:386:ASN:HD22	1:A:386:ASN:HA	1.66	0.43
1:A:370:LEU:HD13	1:A:373:ARG:NH2	2.34	0.43
1:A:269:ILE:HG22	2:A:441:FDQ:O4	2.18	0.42
1:A:377:VAL:HG12	3:A:684:HOH:O	2.18	0.42
1:A:356:LYS:HG3	1:A:413:TYR:OH	2.19	0.42
1:A:295:ILE:HD13	1:A:298:LEU:HB2	2.01	0.42
1:A:424:LEU:CD2	1:A:433:VAL:HG22	2.50	0.42
1:A:298:LEU:HD12	1:A:386:ASN:HB3	2.02	0.41
1:A:317:VAL:HB	1:A:318:PRO:HD2	2.02	0.41
1:A:426:PRO:HA	1:A:431:CYS:HA	2.03	0.41
1:A:304:HIS:CE1	3:A:603:HOH:O	2.74	0.41
1:A:406:LYS:HD2	1:A:410:GLU:OE2	2.20	0.41

Continued from previous page...

Continued on next page...



Atom-1			Clash overlap (Å)	
1:A:325:LEU:CD2	1:A:377:VAL:HG11	2.42	0.41	
1:A:257:ARG:HD2	2:A:441:FDQ:O3P	2.21	0.40	
1:A:362:PRO:O	1:A:363:LYS:HB2	2.21	0.40	

Continued from previous page...

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:267:GLY:N	3:A:621:HOH:O[2_456]	1.81	0.39
1:A:266:ARG:C	3:A:621:HOH:O[2_456]	2.15	0.05

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed Outliers		Percentiles	
1	А	189/191~(99%)	172 (91%)	15 (8%)	2(1%)	14 26	

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	А	405	ASP
1	А	428	ALA

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed Rotameric		Outliers	Percentiles	
1	А	168/168~(100%)	164 (98%)	4 (2%)	49 74	

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	256	CYS
1	А	332	VAL
1	А	338	TYR
1	А	384	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	281	GLN
1	А	310	GLN
1	А	354	GLN
1	А	386	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the



expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	e Chain R	Chain	Res	Link	Bo	ond leng	ths	B	ond ang	les
NIOI			nam res		Counts $ $ RMSZ $ $ $\# Z > 2$ $ $			Counts	RMSZ	# Z >2	
2	FDQ	А	441	-	17,18,18	7.78	3 (17%)	25,28,28	1.45	3 (12%)	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FDQ	А	441	-	-	10/11/27/27	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	$\mathrm{Ideal}(\mathrm{\AA})$
2	А	441	FDQ	P1-C21	-31.58	1.47	1.78
2	А	441	FDQ	P1-O3P	-3.38	1.47	1.54
2	А	441	FDQ	P1-O2P	-2.53	1.49	1.54

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	$Observed(^{o})$	$Ideal(^{o})$
2	А	441	FDQ	O1P-P1-C21	4.67	120.20	111.54
2	А	441	FDQ	P2-O6-C6	2.57	125.38	118.30
2	А	441	FDQ	C3-C4-C5	2.03	106.58	102.64

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	А	441	FDQ	C2-C21-P1-O1P
2	А	441	FDQ	C2-C21-P1-O2P
2	А	441	FDQ	O5-C2-C21-P1
2	А	441	FDQ	C6-O6-P2-O4P
2	А	441	FDQ	C6-O6-P2-O5P
2	А	441	FDQ	C6-O6-P2-O6P
2	А	441	FDQ	C4-C5-C6-O6
2	А	441	FDQ	C2-C21-P1-O3P
2	А	441	FDQ	O5-C5-C6-O6
2	А	441	FDQ	C5-C6-O6-P2




There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	А	441	FDQ	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers (i)

There are no such residues in this entry.



5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

