

Full wwPDB X-ray Structure Validation Report (i)

Nov 20, 2023 – 12:36 AM JST

PDB ID : 7C91

Title : Blasnase-T13A with D-asn

Authors: Lu, F.; Ran, T.; Jiao, L.; Wang, W.

Deposited on : 2020-06-04

Resolution : 1.98 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.36

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.98 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},{\rm resolution\ range}(\mathring{\rm A})) \end{array}$
R_{free}	130704	11647 (2.00-1.96)
Clashscore	141614	1014 (1.98-1.98)
Ramachandran outliers	138981	1006 (1.98-1.98)
Sidechain outliers	138945	1006 (1.98-1.98)
RSRZ outliers	127900	11410 (2.00-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
			11%		
1	A	376	77%	8%	15%
	_		5%		
1	В	376	78%	7%	15%



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 10292 atoms, of which 4967 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called L-asparaginase.

\mathbf{Mol}	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
1	В	321	Total 4971	C 1580	H 2484	N 420	O 479	S 8	114	0	0
1	A	320	Total 4929	C 1565		N 419	O 472	S 8	116	0	0

There are 108 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
В	-53	MET	-	initiating methionine	UNP A0A1Q8GMZ7
В	-52	HIS	-	expression tag	UNP A0A1Q8GMZ7
В	-51	HIS	-	expression tag	UNP A0A1Q8GMZ7
В	-50	HIS	-	expression tag	UNP A0A1Q8GMZ7
В	-49	HIS	-	expression tag	UNP A0A1Q8GMZ7
В	-48	HIS	-	expression tag	UNP A0A1Q8GMZ7
В	-47	HIS	-	expression tag	UNP A0A1Q8GMZ7
В	-46	SER	-	expression tag	UNP A0A1Q8GMZ7
В	-45	SER	-	expression tag	UNP A0A1Q8GMZ7
В	-44	GLY	-	expression tag	UNP A0A1Q8GMZ7
В	-43	LEU	-	expression tag	UNP A0A1Q8GMZ7
В	-42	VAL	-	expression tag	UNP A0A1Q8GMZ7
В	-41	PRO	-	expression tag	UNP A0A1Q8GMZ7
В	-40	ARG	-	expression tag	UNP A0A1Q8GMZ7
В	-39	GLY	-	expression tag	UNP A0A1Q8GMZ7
В	-38	SER	-	expression tag	UNP A0A1Q8GMZ7
В	-37	GLY	-	expression tag	UNP A0A1Q8GMZ7
В	-36	MET	-	expression tag	UNP A0A1Q8GMZ7
В	-35	LYS	-	expression tag	UNP A0A1Q8GMZ7
В	-34	GLU	-	expression tag	UNP A0A1Q8GMZ7
В	-33	THR	_	expression tag	UNP A0A1Q8GMZ7
В	-32	ALA	-	expression tag	UNP A0A1Q8GMZ7
В	-31	ALA	-	expression tag	UNP A0A1Q8GMZ7
В	-30	ALA	-	expression tag	UNP A0A1Q8GMZ7
В	-29	LYS	-	expression tag	UNP A0A1Q8GMZ7

Continued on next page...



Continued from previous page...

Chain	Residue	Modelled Modelled	Actual	Comment	Reference
В	-28	PHE	-	expression tag	UNP A0A1Q8GMZ7
В	-27	GLU	-	expression tag	UNP A0A1Q8GMZ7
В	-26	ARG	-	expression tag	UNP A0A1Q8GMZ7
В	-25	GLN	-	expression tag	UNP A0A1Q8GMZ7
В	-24	HIS	-	expression tag	UNP A0A1Q8GMZ7
В	-23	MET	-	expression tag	UNP A0A1Q8GMZ7
В	-22	ASP	-	expression tag	UNP A0A1Q8GMZ7
В	-21	SER	-	expression tag	UNP A0A1Q8GMZ7
В	-20	PRO	-	expression tag	UNP A0A1Q8GMZ7
В	-19	ASP	-	expression tag	UNP A0A1Q8GMZ7
В	-18	LEU	-	expression tag	UNP A0A1Q8GMZ7
В	-17	GLY	-	expression tag	UNP A0A1Q8GMZ7
В	-16	THR	-	expression tag	UNP A0A1Q8GMZ7
В	-15	ASP	-	expression tag	UNP A0A1Q8GMZ7
В	-14	ASP	-	expression tag	UNP A0A1Q8GMZ7
В	-13	ASP	-	expression tag	UNP A0A1Q8GMZ7
В	-12	ASP	-	expression tag	UNP A0A1Q8GMZ7
В	-11	LYS	-	expression tag	UNP A0A1Q8GMZ7
В	-10	ALA	-	expression tag	UNP A0A1Q8GMZ7
В	-9	MET	-	expression tag	UNP A0A1Q8GMZ7
В	-8	ALA	-	expression tag	UNP A0A1Q8GMZ7
В	-7	ASP	-	expression tag	UNP A0A1Q8GMZ7
В	-6	ILE	-	expression tag	UNP A0A1Q8GMZ7
В	-5	GLY	-	expression tag	UNP A0A1Q8GMZ7
В	-4	SER	-	expression tag	UNP A0A1Q8GMZ7
В	-3	GLU	-	expression tag	UNP A0A1Q8GMZ7
В	-2	PHE	-	expression tag	UNP A0A1Q8GMZ7
В	-1	GLU	-	expression tag	UNP A0A1Q8GMZ7
В	0	LEU	-	expression tag	UNP A0A1Q8GMZ7
A	-53	MET	-	initiating methionine	UNP A0A1Q8GMZ7
A	-52	HIS	-	expression tag	UNP A0A1Q8GMZ7
A	-51	HIS	-	expression tag	UNP A0A1Q8GMZ7
A	-50	HIS	-	expression tag	UNP A0A1Q8GMZ7
A	-49	HIS	-	expression tag	UNP A0A1Q8GMZ7
A	-48	HIS	-	expression tag	UNP A0A1Q8GMZ7
A	-47	HIS	-	expression tag	UNP A0A1Q8GMZ7
A	-46	SER	-	expression tag	UNP A0A1Q8GMZ7
A	-45	SER	-	expression tag	UNP A0A1Q8GMZ7
A	-44	GLY	-	expression tag	UNP A0A1Q8GMZ7
A	-43	LEU	-	expression tag	UNP A0A1Q8GMZ7
A	-42	VAL	-	expression tag	UNP A0A1Q8GMZ7
A	-41	PRO	-	expression tag	UNP A0A1Q8GMZ7

Continued on next page...




 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled	Actual	Comment	Reference
A	-40	ARG	-	expression tag	UNP A0A1Q8GMZ7
A	-39	GLY	-	expression tag	UNP A0A1Q8GMZ7
A	-38	SER	-	expression tag	UNP A0A1Q8GMZ7
A	-37	GLY	-	expression tag	UNP A0A1Q8GMZ7
A	-36	MET	-	expression tag	UNP A0A1Q8GMZ7
A	-35	LYS	-	expression tag	UNP A0A1Q8GMZ7
A	-34	GLU	-	expression tag	UNP A0A1Q8GMZ7
A	-33	THR	-	expression tag	UNP A0A1Q8GMZ7
A	-32	ALA	-	expression tag	UNP A0A1Q8GMZ7
A	-31	ALA	-	expression tag	UNP A0A1Q8GMZ7
A	-30	ALA	-	expression tag	UNP A0A1Q8GMZ7
A	-29	LYS	-	expression tag	UNP A0A1Q8GMZ7
A	-28	PHE	-	expression tag	UNP A0A1Q8GMZ7
A	-27	GLU	-	expression tag	UNP A0A1Q8GMZ7
A	-26	ARG	-	expression tag	UNP A0A1Q8GMZ7
A	-25	GLN	-	expression tag	UNP A0A1Q8GMZ7
A	-24	HIS	-	expression tag	UNP A0A1Q8GMZ7
A	-23	MET	-	expression tag	UNP A0A1Q8GMZ7
A	-22	ASP	-	expression tag	UNP A0A1Q8GMZ7
A	-21	SER	-	expression tag	UNP A0A1Q8GMZ7
A	-20	PRO	-	expression tag	UNP A0A1Q8GMZ7
A	-19	ASP	-	expression tag	UNP A0A1Q8GMZ7
A	-18	LEU	-	expression tag	UNP A0A1Q8GMZ7
A	-17	GLY	-	expression tag	UNP A0A1Q8GMZ7
A	-16	THR	-	expression tag	UNP A0A1Q8GMZ7
A	-15	ASP	-	expression tag	UNP A0A1Q8GMZ7
A	-14	ASP	-	expression tag	UNP A0A1Q8GMZ7
A	-13	ASP	-	expression tag	UNP A0A1Q8GMZ7
A	-12	ASP	-	expression tag	UNP A0A1Q8GMZ7
A	-11	LYS	-	expression tag	UNP A0A1Q8GMZ7
A	-10	ALA	-	expression tag	UNP A0A1Q8GMZ7
A	-9	MET	-	expression tag	UNP A0A1Q8GMZ7
A	-8	ALA	-	expression tag	UNP A0A1Q8GMZ7
A	-7	ASP	-	expression tag	UNP A0A1Q8GMZ7
A	-6	ILE	-	expression tag	UNP A0A1Q8GMZ7
A	-5	GLY	-	expression tag	UNP A0A1Q8GMZ7
A	-4	SER	-	expression tag	UNP A0A1Q8GMZ7
A	-3	GLU	-	expression tag	UNP A0A1Q8GMZ7
A	-2	PHE	-	expression tag	UNP A0A1Q8GMZ7
A	-1	GLU	-	expression tag	UNP A0A1Q8GMZ7
A	0	LEU	-	expression tag	UNP A0A1Q8GMZ7

 \bullet Molecule 2 is D-ASPARAGINE (three-letter code: DSG) (formula: $C_4H_8N_2O_3)$ (labeled as



"Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf		
2	В	1	Total 17					2	0
2	A	1	Total 17			N 2		2	0

• Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	В	3	Total Mg 3 3	0	0
3	A	2	Total Mg 2 2	0	0

• Molecule 4 is FORMIC ACID (three-letter code: FMT) (formula: CH_2O_2) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	H	0	1	0

• Molecule 5 is water.


\mathbf{Mol}	Chain	Residues	Atoms	$\mathbf{ZeroOcc}$	AltConf
5	В	193	Total O 193 193	0	0
5	A	155	Total O 155 155	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: L-asparaginase





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants	92.26Å 92.26Å 231.46Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	72.25 - 1.98	Depositor
Resolution (A)	72.14 - 1.98	EDS
% Data completeness	99.7 (72.25-1.98)	Depositor
(in resolution range)	99.7 (72.14-1.98)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.98 (at 1.98Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
D D.	0.182 , 0.211	Depositor
R, R_{free}	0.185 , 0.214	DCC
R_{free} test set	3524 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	39.8	Xtriage
Anisotropy	0.172	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.42 , 46.4	EDS
L-test for twinning ²	$ < L >=0.46, < L^2>=0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	10292	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.29% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: FMT, DSG, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
MIOI	Wioi Chain		# Z > 5	RMSZ	# Z > 5	
1	A	0.39	0/2510	0.54	0/3400	
1	В	0.40	0/2534	0.54	0/3432	
All	All	0.40	0/5044	0.54	0/6832	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2464	2465	2449	20	0
1	В	2487	2484	2476	21	0
2	A	9	8	7	1	0
2	В	9	8	7	0	0
3	A	2	0	0	0	0
3	В	3	0	0	0	0
4	A	3	2	1	0	0
5	A	155	0	0	2	0
5	В	193	0	0	4	1
All	All	5325	4967	4940	38	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including



hydrogen atoms). The all-atom clashscore for this structure is 4.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic	Clash
1 1 22 0110 1172	1 1 1 0 0 FF FF C C D 1	distance (Å)	overlap (Å)
1:A:38:CYS:HB3	1:A:126:TYR:CD1	2.06	0.91
1:A:34:LEU:HA	1:A:37:MET:HE3	1.58	0.84
1:A:38:CYS:HB3	1:A:126:TYR:HD1	1.52	0.74
1:B:227:ARG:NH1	5:B:502:HOH:O	2.13	0.70
1:B:121:GLN:OE1	5:B:501:HOH:O	2.11	0.67
1:A:131:HIS:HB3	1:A:182:ILE:HD13	1.80	0.64
1:A:321:CYS:O	5:A:501:HOH:O	2.16	0.61
1:B:13:THR:HA	1:B:16:SER:HB3	1.85	0.59
1:B:14:ILE:O	1:B:118:PRO:HG3	2.02	0.59
1:A:291:ILE:HG12	1:A:317:LYS:HG2	1.86	0.57
1:B:3:LYS:NZ	5:B:507:HOH:O	2.39	0.56
1:B:247:MET:HE3	1:B:251:ILE:HD11	1.87	0.55
1:B:19:THR:HG22	1:B:20:GLU:H	1.71	0.55
1:A:48:VAL:HG12	1:A:50:PRO:HD3	1.90	0.53
1:B:3:LYS:HD3	1:B:137:CYS:HB3	1.91	0.52
1:B:276:TYR:HB3	1:B:278:TYR:CE2	2.45	0.51
1:A:276:TYR:HB3	1:A:278:TYR:CE2	2.45	0.51
1:A:4:LYS:HG2	1:A:80:TYR:CD1	2.46	0.51
1:B:35:ALA:O	1:B:40:LEU:HD11	2.12	0.49
1:B:247:MET:CE	1:B:251:ILE:HD11	2.42	0.49
1:A:34:LEU:HA	1:A:37:MET:CE	2.36	0.49
1:B:24:LEU:HB2	1:A:276:TYR:CD1	2.48	0.49
1:A:226:VAL:HG22	1:A:259:VAL:HG21	1.95	0.48
1:B:19:THR:HG22	1:B:20:GLU:N	2.29	0.46
1:B:185:ASP:OD1	5:B:503:HOH:O	2.21	0.44
1:A:77:ASP:OD1	1:A:79:SER:OG	2.32	0.44
1:A:314:GLU:O	1:A:319:LYS:NZ	2.45	0.44
1:A:19:THR:HG22	1:A:23:ARG:O	2.18	0.44
1:B:245:PRO:HG3	1:A:58:HIS:CD2	2.52	0.43
1:B:251:ILE:HG23	1:B:261:ILE:HD12	2.01	0.43
1:B:235:LEU:HD22	1:B:247:MET:HE3	2.01	0.43
1:A:33:GLU:O	1:A:37:MET:HE2	2.19	0.42
1:B:27:GLY:O	1:B:53:GLN:NE2	2.53	0.42
1:B:214:TYR:O	1:A:57:MET:HE1	2.20	0.42
1:B:18:LYS:H	1:B:18:LYS:HD2	1.85	0.41
1:A:196:ARG:NH1	5:A:502:HOH:O	2.16	0.41
1:A:13:THR:OG1	2:A:401:DSG:N	2.54	0.41
1:B:18:LYS:HE3	1:B:24:LEU:CD2	2.51	0.41



All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$\begin{array}{c} \text{Clash} \\ \text{overlap } (\text{\AA}) \end{array}$
5:B:553:HOH:O	5:B:553:HOH:O[7_555]	2.11	0.09

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	A	318/376 (85%)	313 (98%)	5 (2%)	0	100	100
1	В	319/376~(85%)	314 (98%)	5 (2%)	0	100	100
All	All	637/752 (85%)	627 (98%)	10 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	A	260/310 (84%)	257 (99%)	3 (1%)	71 67		
1	В	265/310~(86%)	261 (98%)	4 (2%)	65 59		
All	All	525/620~(85%)	518 (99%)	7 (1%)	69 64		

All (7) residues with a non-rotameric sidechain are listed below:



Mol	Chain	Res	Type
1	В	18	LYS
1	В	52	PHE
1	В	57	MET
1	В	322	TYR
1	A	18	LYS
1	A	52	PHE
1	A	75	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 8 ligands modelled in this entry, 5 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trino	Chain	Res	Link	В	ond leng	$_{ m gths}$	В	ond ang	gles
MIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	DSG	A	401	-	6,8,8	0.91	0	8,10,10	1.19	1 (12%)
4	FMT	A	402	3	2,2,2	0.62	0	1,1,1	0.21	0
2	DSG	В	401	-	6,8,8	1.09	0	8,10,10	1.27	1 (12%)



In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	DSG	A	401	-	-	2/8/8/8	-
2	DSG	В	401	-	-	2/8/8/8	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
2	A	401	DSG	OXT-C-O	-2.65	118.08	124.09
2	В	401	DSG	OXT-C-O	-2.53	118.33	124.09

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	В	401	DSG	OXT-C-CA-CB
2	A	401	DSG	OXT-C-CA-CB
2	В	401	DSG	O-C-CA-CB
2	A	401	DSG	O-C-CA-CB


There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	DSG	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\#\mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q<0.9
1	A	320/376~(85%)	1.15	42 (13%) 3 3	28, 40, 76, 105	0
1	В	321/376~(85%)	0.83	19 (5%) 22 24	26, 38, 65, 92	0
All	All	641/752 (85%)	0.99	61 (9%) 8 9	26, 39, 71, 105	0

All (61) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	19	THR	11.7
1	A	21	SER	9.5
1	A	41	PRO	9.4
1	A	20	GLU	9.4
1	A	34	LEU	7.2
1	A	22	GLY	7.1
1	A	25	ALA	6.9
1	A	38	CYS	5.8
1	В	322	TYR	5.7
1	A	28	ALA	5.4
1	A	35	ALA	5.3
1	A	26	ALA	5.1
1	В	276	TYR	5.0
1	A	40	LEU	5.0
1	A	24	LEU	4.8
1	A	17	ARG	4.7
1	A	39	SER	4.6
1	A	27	GLY	4.6
1	A	29	ILE	4.5
1	A	44	VAL	4.3
1	A	185	ASP	4.1
1	A	33	GLU	4.1
1	A	42	GLU	4.1
1	В	42	GLU	4.0

Continued on next page...



Continued from previous page...

Mol	nuea fron Chain	Res	Type	RSRZ	
1	В	20	GLU	4.0	
1	A	23	ARG	3.7	
1	A	37	MET	3.6	
1	В	26	ALA	3.5	
1	A	18	LYS	3.4	
1	A	121	GLN	3.3	
1	A	119	GLU	3.1	
1	В	43	ASP	3.1	
1	A	120	GLN	3.1	
1	A	126	TYR	3.0	
1	В	19	THR	3.0	
1	A	117	ALA	2.9	
1	A	2	LYS	2.8	
1	A	14	ILE	2.6	
1	A	106	GLU	2.6	
1	В	18	LYS	2.6	
1	В	21	SER	2.6	
1	В	275	THR	2.5	
1	A	186	LYS	2.5	
1	В	273	TYR	2.4	
1	A	36	GLU	2.4	
1	В	52	PHE	2.4	
1	В	10	THR	2.3	
1	A	15	ALA	2.2	
1	В	38	CYS	2.2	
1	В	277	ASP	2.2	
1	A	3	LYS	2.2	
1	A	122	GLY	2.1	
1	A	43	ASP	2.1	
1	A	197	ASP	2.1	
1	В	113	GLY	2.1	
1	A	65	LEU	2.1	
1	В	127	THR	2.1	
1	A	16	SER	2.1	
1	В	27	GLY	2.1	
1	A	275	THR	2.0	
1	В	114	SER	2.0	

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



6.3 Carbohydrates (i)

There are no monosaccharides in this entry.


6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
3	MG	A	403	1/1	0.86	0.17	49,49,49,49	0
3	MG	A	404	1/1	0.86	0.12	59,59,59,59	0
2	DSG	В	401	9/9	0.89	0.21	34,40,47,47	2
2	DSG	A	401	9/9	0.91	0.20	41,49,60,60	2
3	MG	В	404	1/1	0.91	0.07	50,50,50,50	0
3	MG	В	403	1/1	0.92	0.25	56,56,56,56	0
3	MG	В	402	1/1	0.95	0.12	53,53,53,53	0
4	FMT	A	402	3/3	0.97	0.09	39,47,47,47	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



















Electron density around DSG A 401: 2mF_o-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)





















6.5 Other polymers (i)

There are no such residues in this entry.

