

## wwPDB EM Validation Summary Report (i)

#### Nov 11, 2023 – 08:28 pm GMT

PDB ID : 8CAM

EMDB ID : EMD-16530

Title: Evernimic bound to the 50S subunit

Authors : Paternoga, H.; Crowe-McAuliffe, C.; Novacek, J.; Wilson, D.N.

Deposited on : 2023-01-24

Resolution : 1.86 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev70

Mogul : 1.8.4, CSD as541be (2020)

MolProbity : 4.02b-467 buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

MapQ : 1.9.9

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

### 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $ELECTRON\ MICROSCOPY$ 

The reported resolution of this entry is 1.86 Å.


Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	${ m EM\ structures} \ (\#{ m Entries})$
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826
RNA backbone	4643	859

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion <40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	0	55	91%	9%
2	1	46	96%	• •
3	2	65	97%	
4	3	38	100%	
5	a	2904	81% 11%	7%
6	b	120	27% • 72%	
7	c	273	99%	





 $Continued\ from\ previous\ page...$ 

Mol	Chain	Length	Quality of chain
8	d	209	99%
9	e	201	9%
10	g	177	43%
11	h	149	9%
			21% • 77%
12	i	142	98%
13	j	123	98%
14	k	144	99%
15	l	136	100%
16	m	127	93% 7%
17	О	115	97%
18	p	118	98%
19	q	103	98%
20	r	110	98%
21	s	100	9% • 7%
22	t	104	89% · 10%
23	u	94	80% 20%
24	V	85	8%
25		78	<u> </u>
	W		96% · • • 25%
26	X	63	92% • 5% 7%
27	У	59	98%
28	Z	57	95% 5%



## 2 Entry composition (i)

There are 34 unique types of molecules in this entry. The entry contains 86648 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Large ribosomal subunit protein bL33.

Mol	Chain	Residues		Aton	ıs		AltConf	Trace
1	0	50	Total 413	C 267	N 75	O 71	0	0

• Molecule 2 is a protein called Large ribosomal subunit protein bL34.

Mol	Chain	Residues		Ato	ms			AltConf	Trace
9	1	45	Total	С	N	О	S	0	0
	1	40	367	222	88	55	2	0	U

• Molecule 3 is a protein called Large ribosomal subunit protein bL35.

Mol	Chain	Residues	${f Atoms}$					AltConf	Trace
3	2	64	Total 504		N 105	O 74	S	0	0

• Molecule 4 is a protein called Large ribosomal subunit protein bL36A.

Mol	Chain	Residues		Ato	$\mathbf{m}\mathbf{s}$			AltConf	Trace
4	3	38	Total 302	C 185	N 65	O 48	S 4	0	0

• Molecule 5 is a RNA chain called 23S rRNA.

Mol	Chain	Residues				AltConf	Trace		
5	a	2690	Total 57788	C 25784	N 10657	O 18657	P 2690	0	0

• Molecule 6 is a RNA chain called 5S rRNA.

Mol	Chain	Residues		A	toms			AltConf	Trace
6	b	34	Total 734	C 326	N 134	O 240	P 34	0	0



• Molecule 7 is a protein called Large ribosomal subunit protein uL2.

Mol	Chain	Residues		Ato	oms			AltConf	Trace
7	c	271	Total 2082	C 1288	N 423	O 364	S 7	0	0

• Molecule 8 is a protein called Large ribosomal subunit protein uL3.

Mol	Chain	Residues		At	oms			AltConf	Trace
Q	d	207	Total	С	N	О	S	0	0
0	u	207	1552	972	286	291	3	0	U

• Molecule 9 is a protein called Large ribosomal subunit protein uL4.

Mol	Chain	Residues		At	oms			AltConf	Trace
9	e	197	Total 1524	C 959	N 278	O 282	S 5	0	0

• Molecule 10 is a protein called Large ribosomal subunit protein uL6.

Mol	Chain	Residues		At	oms			AltConf	Trace
10	œ	174	Total	С	N	О	S	0	0
10	g	174	1304	820	239	243	2	U	U

• Molecule 11 is a protein called Large ribosomal subunit protein bL9.

N	/Iol	Chain	Residues		Ato	ms			AltConf	Trace
	11	h	34	Total 257	C 167	N 46	O 43	S 1	0	0

• Molecule 12 is a protein called Large ribosomal subunit protein uL13.

Mol	Chain	Residues		At	oms	AltConf	Trace		
12	i	141	Total 1121	C 709	N 211	O 198	S 3	0	0

• Molecule 13 is a protein called Large ribosomal subunit protein uL14.

Mol	Chain	Residues		At	oms	AltConf	Trace		
13	j	123	Total 946	C 593	N 181	O 166	S 6	0	0

• Molecule 14 is a protein called 50S ribosomal protein L15.



Mol	Chain	Residues		At	oms	AltConf	Trace		
1.4	1,	144	Total	С	N	О	S	0	0
14	K	144	1053	654	207	190	2	0	

• Molecule 15 is a protein called Large ribosomal subunit protein uL16.

Mol	Chain	Residues		At	oms			AltConf	Trace
15	l	136	Total 1075	C 686	N 205	O 177	S 7	0	0

There is a discrepancy between the modelled and reference sequences:

Cha	n Residue	Modelled	Actual	Comment	Reference
1	82	MS6	MET	modified residue	UNP P0ADY7

• Molecule 16 is a protein called Large ribosomal subunit protein bL17.

Mol	Chain	Residues		At	oms			AltConf	Trace
16	m	118	Total 945		N 194	O 161	S 5	0	0

• Molecule 17 is a protein called Large ribosomal subunit protein bL19.

Mol	Chain	Residues		Atoms					Trace
17	О	114	Total 917	C 574	N 179	O 163	S 1	0	0

• Molecule 18 is a protein called Large ribosomal subunit protein bL20.

Mol	Chain	Residues		Ato	ms		AltConf	Trace
18	р	117	Total 947	C 604	N 192	O 151	0	0

• Molecule 19 is a protein called Large ribosomal subunit protein bL21.

Mol	Chain	Residues		At	oms	AltConf	Trace		
19	q	101	Total 803	C 509	N 151	O 141	S 2	0	0

• Molecule 20 is a protein called Large ribosomal subunit protein uL22.



Mol	Chain	Residues		At	oms	AltConf	Trace		
20	r	109	Total	С	N	О	S	0	0
20	1	109	845	526	162	154	3	0	

• Molecule 21 is a protein called Large ribosomal subunit protein uL23.

Mol	Chain	Residues	Atoms				AltConf	Trace	
91	g	93	Total	С	N	О	S	0	0
21	8	90	738	466	139	131	2	0	U

• Molecule 22 is a protein called Large ribosomal subunit protein uL24.

Mol	Chain	Residues		Ato	ms		AltConf	Trace
22	t	94	Total 724	C 457	N 136	O 131	0	0

• Molecule 23 is a protein called 50S ribosomal protein L25.

Mol	Chain	Residues		Atoms				AltConf	Trace
22	11	75	Total	С	N	О	S	0	0
23	u	10	607	391	111	103	2		U

• Molecule 24 is a protein called Large ribosomal subunit protein bL27.

Mol	Chain	Residues		Atoms				AltConf	Trace
24	V	74	Total 559	C 348	N 112	O 98	S 1	0	0

• Molecule 25 is a protein called Large ribosomal subunit protein bL28.

Mol	Chain	Residues		Atoms				AltConf	Trace
25	W	77	Total 625	C 388	N 129	O 106	S 2	0	0

• Molecule 26 is a protein called Large ribosomal subunit protein uL29.

Mol	Chain	Residues		Ato	ms			AltConf	Trace
26	X	60	Total 491	C 303	N 96	O 91	S 1	0	0

• Molecule 27 is a protein called Large ribosomal subunit protein uL30.



Mol	Chain	Residues	Atoms				AltConf	Trace	
27	**	50	Total	С	N	О	S	0	0
21	У	58	449	281	87	79	2	0	U


• Molecule 28 is a protein called Large ribosomal subunit protein bL32.

Mol	Chain	Residues		Ato	ms			AltConf	Trace
28	Z	54	Total 429	C 260	N 91	O 77	S 1	0	0

 $\bullet$  Molecule 29 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	AltConf
29	3	1	Total Zn 1 1	0

 $\bullet$  Molecule 30 is ACETATE ION (three-letter code: ACT) (formula:  $\mathrm{C_2H_3O_2}).$ 



Mol	Chain	Residues	Atoms	AltConf
30	a	1	Total C O 4 2 2	0

• Molecule 31 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

$\mathbf{Mol}$	Chain	Residues	Atoms	AltConf
31	a	237	Total Mg 237 237	0

Continued on next page...




Continued from previous page...

Mol	Chain	Residues	Atoms	AltConf
31	b	3	Total Mg 3 3	0
31	С	3	Total Mg 3 3	0
31	d	1	Total Mg 1 1	0
31	р	1	Total Mg 1 1	0
31	Z	1	Total Mg 1 1	0

• Molecule 32 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms	AltConf
32	a	68	Total K 68 68	0
32	c	2	Total K 2 2	0
32	е	1	Total K 1 1	0
32	t	1	Total K 1 1	0

• Molecule 33 is Evernimic in (three-letter code: 6O1) (formula:  $C_{70}H_{97}Cl_2NO_{38}$ ) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms				AltConf	
33		1	Total	С	Cl	N	О	0
33	a	1	111	70	2	1	38	0

### • Molecule 34 is water.

Mol	Chain	Residues	Atoms	AltConf
34	0	5	Total O 5 5	0
34	1	32	Total O 32 32	0
34	2	30	Total O 30 30	0
34	3	7	Total O 7 7	0
34	a	5251	Total O 5251 5251	0
34	b	63	Total O 63 63	0
34	С	115	Total O 115 115	0
34	d	74	Total O 74 74	0
34	e	51	Total O 51 51	0
34	g	1	Total O 1 1	0
34	h	2	Total O 2 2	0
34	i	36	Total O 36 36	0
34	j	32	Total O 32 32	0
34	k	50	Total O 50 50	0
34	1	45	Total O 45 45	0
34	m	38	Total O 38 38	0
34	О	25	Total O 25 25	0
34	р	51	Total O 51 51	0
34	q	29	Total O 29 29	0

Continued on next page...



 $Continued\ from\ previous\ page...$ 

Mol	Chain	Residues	Atoms	AltConf
34	r	48	Total O 48 48	0
34	s	12	Total O 12 12	0
34	t	11	Total O 11 11	0
34	u	6	Total O 6 6	0
34	V	23	Total O 23 23	0
34	W	23	Total O 23 23	0
34	X	4	Total O 4 4	0
34	у	11	Total O 11 11	0
34	Z	38	Total O 38 38	0




Chain a:

#### 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Large ribosomal subunit protein bL33




81%



11%

















 $\bullet$  Molecule 18: Large ribosomal subunit protein bL20

Chain p: 98%



• Molecule 19: Large ribosomal subunit protein bL21

Chain q: 98%



• Molecule 20: Large ribosomal subunit protein uL22

Chain r: 98% ...



• Molecule 21: Large ribosomal subunit protein uL23

Chain s: 90% . 7%



• Molecule 22: Large ribosomal subunit protein uL24

Chain t: 89% • 10%




• Molecule 23: 50S ribosomal protein L25

Chain u: 80% 20%



• Molecule 24: Large ribosomal subunit protein bL27







# 4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	514855	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE	Depositor
	CORRECTION	
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{Å}^2)$	44	Depositor
Minimum defocus (nm)	400	Depositor
Maximum defocus (nm)	1600	Depositor
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.414	Depositor
Minimum map value	-0.132	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.006	Depositor
Recommended contour level	0.0325	Depositor
Map size (Å)	390.144, 390.144, 390.144	wwPDB
Map dimensions	512, 512, 512	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.762, 0.762, 0.762	Depositor



## 5 Model quality (i)

### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: MEQ, K, ZN, 1MG, 5MC, G7M, PSU, 5MU, 2MA, 6O1, 4D4, ACT, OMU, 6MZ, MS6, 2MG, OMC, OMG, H2U, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Clasia.	Bond	lengths	Е	Bond angles
Mol	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	0	0.31	0/420	0.67	0/560
2	1	0.30	0/370	0.74	0/487
3	2	0.30	0/513	0.68	0/676
4	3	0.29	0/303	0.75	0/397
5	a	0.51	0/64215	1.12	45/100167 (0.0%)
6	b	0.52	0/821	1.15	0/1280
7	С	0.31	0/2121	0.70	0/2852
8	d	0.30	0/1562	0.66	0/2102
9	е	0.28	0/1542	0.62	0/2072
10	g	0.30	0/1324	0.68	0/1794
11	h	0.33	0/259	0.78	0/348
12	i	0.29	0/1144	0.65	0/1541
13	j	0.30	0/955	0.71	0/1279
14	k	0.33	0/1062	0.64	0/1413
15	1	0.29	0/1073	0.70	0/1433
16	m	0.30	0/958	0.68	0/1281
17	О	0.30	0/929	0.69	0/1242
18	p	0.29	0/960	0.66	0/1278
19	q	0.31	0/815	0.67	0/1087
20	r	0.29	0/852	0.67	0/1142
21	S	0.27	0/744	0.63	0/994
22	t	0.29	0/729	0.67	0/968
23	u	0.33	0/615	0.67	0/819
24	V	0.33	0/566	0.65	0/750
25	W	0.31	0/635	0.68	0/848
26	X	0.27	0/492	0.61	0/655
27	У	0.29	0/453	0.64	0/605
28	Z	0.33	0/435	0.71	0/581
All	All	0.47	0/86867	1.04	45/130651 (0.0%)



Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a maintain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	1	0	1
9	е	0	2
12	i	0	2
17	О	0	1
18	р	0	1
21	$\mathbf{s}$	0	1
24	V	0	1
25	W	0	1
26	X	0	2
All	All	0	12

There are no bond length outliers.

The worst 5 of 45 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$Observed(^o)$	$Ideal(^{o})$
5	a	512	G	O4'-C1'-N9	9.89	116.11	108.20
5	a	781	A	O3'-P-O5'	-6.96	90.77	104.00
5	a	2824	С	O5'-P-OP2	-6.74	99.64	105.70
5	a	1971	U	O3'-P-O5'	-6.73	91.22	104.00
5	a	395	U	O4'-C1'-N1	6.70	113.56	108.20

There are no chirality outliers.

5 of 12 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	1	41	ARG	Sidechain
9	е	21	ARG	Sidechain
9	е	88	ARG	Sidechain
12	i	95	ARG	Sidechain
12	i	96	ARG	Sidechain

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	0	413	0	448	0	0
2	1	367	0	405	0	0
3	2	504	0	572	0	0
4	3	302	0	340	0	0
5	a	57788	0	29065	0	0
6	b	734	0	367	0	0
7	С	2082	0	2153	0	0
8	d	1552	0	1601	0	0
9	е	1524	0	1596	0	0
10	g	1304	0	1345	0	0
11	h	257	0	283	0	0
12	i	1121	0	1150	0	0
13	j	946	0	1023	0	0
14	k	1053	0	1129	0	0
15	1	1075	0	1145	0	0
16	m	945	0	989	0	0
17	О	917	0	962	0	0
18	p	947	0	1019	0	0
19	q	803	0	829	0	0
20	r	845	0	909	0	0
21	s	738	0	807	0	0
22	t	724	0	773	0	0
23	u	607	0	621	0	0
24	V	559	0	575	0	0
25	W	625	0	652	0	0
26	X	491	0	523	0	0
27	у	449	0	488	0	0
28	Z	429	0	440	0	0
29	3	1	0	0	0	0
30	a	4	0	3	0	0
31	a	237	0	0	0	0
31	b	3	0	0	0	0
31	С	3	0	0	0	0
31	d	1	0	0	0	0
31	p	1	0	0	0	0
31	Z	1	0	0	0	0
32	a	68	0	0	0	0
32	c	2	0	0	0	0
32	e	1	0	0	0	0
32	t	1	0	0	0	0
33	a	111	0	0	0	0
34	0	5	0	0	0	0
34	1	32	0	0	0	0

Continued on next page...



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
34	2	30	0	0	0	0
34	3	7	0	0	0	0
34	a	5251	0	0	0	0
34	b	63	0	0	0	0
34	С	115	0	0	0	0
34	d	74	0	0	0	0
34	е	51	0	0	0	0
34	g	1	0	0	0	0
34	h	2	0	0	0	0
34	i	36	0	0	0	0
34	j	32	0	0	0	0
34	k	50	0	0	0	0
34	1	45	0	0	0	0
34	m	38	0	0	0	0
34	О	25	0	0	0	0
34	p	51	0	0	0	0
34	q	29	0	0	0	0
34	r	48	0	0	0	0
34	S	12	0	0	0	0
34	t	11	0	0	0	0
34	u	6	0	0	0	0
34	V	23	0	0	0	0
34	W	23	0	0	0	0
34	X	4	0	0	0	0
34	у	11	0	0	0	0
34	Z	38	0	0	0	0
All	All	86648	0	52212	0	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

There are no clashes within the asymmetric unit.

There are no symmetry-related clashes.

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.



The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	0	48/55~(87%)	47 (98%)	1 (2%)	0	100	100
2	1	43/46 (94%)	43 (100%)	0	0	100	100
3	2	62/65~(95%)	60 (97%)	2 (3%)	0	100	100
4	3	$36/38 \; (95\%)$	36 (100%)	0	0	100	100
7	c	269/273 (98%)	258 (96%)	11 (4%)	0	100	100
8	d	204/209 (98%)	197 (97%)	7 (3%)	0	100	100
9	e	193/201 (96%)	188 (97%)	5 (3%)	0	100	100
10	g	172/177 (97%)	158 (92%)	13 (8%)	1 (1%)	25	12
11	h	30/149 (20%)	25 (83%)	4 (13%)	1 (3%)	4	0
12	i	139/142 (98%)	138 (99%)	1 (1%)	0	100	100
13	j	121/123 (98%)	115 (95%)	6 (5%)	0	100	100
14	k	142/144 (99%)	139 (98%)	2 (1%)	1 (1%)	22	9
15	1	132/136 (97%)	126 (96%)	6 (4%)	0	100	100
16	m	116/127 (91%)	110 (95%)	6 (5%)	0	100	100
17	О	112/115 (97%)	108 (96%)	4 (4%)	0	100	100
18	p	115/118 (98%)	115 (100%)	0	0	100	100
19	q	97/103 (94%)	97 (100%)	0	0	100	100
20	r	107/110 (97%)	102 (95%)	5 (5%)	0	100	100
21	s	91/100 (91%)	88 (97%)	3 (3%)	0	100	100
22	t	90/104 (86%)	89 (99%)	1 (1%)	0	100	100
23	u	64/94 (68%)	63 (98%)	1 (2%)	0	100	100
24	V	72/85 (85%)	70 (97%)	2 (3%)	0	100	100
25	W	75/78 (96%)	74 (99%)	1 (1%)	0	100	100
26	X	58/63 (92%)	58 (100%)	0	0	100	100
27	У	56/59~(95%)	55 (98%)	1 (2%)	0	100	100
28	Z	52/57 (91%)	51 (98%)	1 (2%)	0	100	100
All	All	2696/2971 (91%)	2610 (97%)	83 (3%)	3 (0%)	54	36

All (3) Ramachandran outliers are listed below:



Mol	Chain	Res	Type
10	g	61	GLY
11	h	7	ASP
14	k	29	LYS

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	0	46/49~(94%)	46 (100%)	0	100	100
2	1	37/38~(97%)	37 (100%)	0	100	100
3	2	51/52~(98%)	50 (98%)	1 (2%)	55	40
4	3	34/34 (100%)	34 (100%)	0	100	100
7	c	$216/218\ (99\%)$	216 (100%)	0	100	100
8	d	162/163~(99%)	162 (100%)	0	100	100
9	e	162/165 (98%)	160 (99%)	2 (1%)	71	62
10	g	135/138 (98%)	133 (98%)	2 (2%)	65	53
11	h	27/114 (24%)	26 (96%)	1 (4%)	34	17
12	i	115/116 (99%)	114 (99%)	1 (1%)	78	72
13	j	104/104 (100%)	102 (98%)	2 (2%)	57	43
14	k	103/103 (100%)	103 (100%)	0	100	100
15	1	107/107 (100%)	107 (100%)	0	100	100
16	m	$98/103\ (95\%)$	98 (100%)	0	100	100
17	0	99/100~(99%)	98 (99%)	1 (1%)	76	69
18	p	89/90~(99%)	89 (100%)	0	100	100
19	q	83/84~(99%)	83 (100%)	0	100	100
20	r	$92/93\ (99\%)$	91 (99%)	1 (1%)	73	65
21	S	80/84 (95%)	78 (98%)	2 (2%)	47	31
22	t	77/85 (91%)	76 (99%)	1 (1%)	69	58
23	u	63/78 (81%)	63 (100%)	0	100	100
24	V	55/63 (87%)	55 (100%)	0	100	100

Continued on next page...



Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
25	W	67/68 (98%)	66 (98%)	1 (2%)	65 53
26	X	54/55~(98%)	54 (100%)	0	100 100
27	у	48/49 (98%)	48 (100%)	0	100 100
28	Z	46/48 (96%)	46 (100%)	0	100 100
All	All	2250/2401 (94%)	2235 (99%)	15 (1%)	84 79

5 of 15 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
13	j	58	LEU
22	t	24	LYS
13	j	105	ARG
25	W	44	LYS
21	S	64	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
7	С	90	ASN
9	е	195	GLN
23	u	12	GLN
26	X	58	ASN

#### 5.3.3 RNA (i)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
5	a	2679/2904 (92%)	292 (10%)	0
6	b	33/120 (27%)	2 (6%)	0
All	All	2712/3024 (89%)	294 (10%)	0

5 of 294 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
5	a	10	A
5	a	12	U
5	a	13	A
5	a	15	G
5	a	34	U



There are no RNA pucker outliers to report.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

23 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Т	Clasica	Dag	T 2 1-	Во	ond leng	ths	В	ond ang	les
Mol	Type	Chain	Res	Link	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	2MA	a	2503	31,32,5	17,25,26	1.04	2 (11%)	17,37,40	1.05	1 (5%)
5	5MU	a	1939	32,5	19,22,23	0.34	0	28,32,35	0.43	0
5	1MG	a	745	5	18,26,27	1.02	2 (11%)	19,39,42	0.47	0
15	4D4	1	81	15	9,11,12	0.53	0	8,13,15	0.88	0
5	OMU	a	2552	5	19,22,23	0.26	0	26,31,34	0.46	0
5	6MZ	a	1618	5	18,25,26	0.70	0	16,36,39	0.91	1 (6%)
5	2MG	a	1835	5	18,26,27	1.06	2 (11%)	16,38,41	0.61	0
5	PSU	a	2504	32,5	18,21,22	0.95	1 (5%)	22,30,33	0.75	0
5	G7M	a	2069	5	20,26,27	1.11	2 (10%)	17,39,42	0.62	0
5	PSU	a	2604	5	18,21,22	0.95	1 (5%)	22,30,33	0.73	1 (4%)
5	PSU	a	2580	32,5	18,21,22	1.03	1 (5%)	22,30,33	0.79	1 (4%)
5	H2U	a	2449	5	18,21,22	0.61	0	21,30,33	0.67	0
5	OMG	a	2251	32,5	18,26,27	0.99	2 (11%)	19,38,41	0.81	0
5	2MG	a	2445	5	18,26,27	0.98	1 (5%)	16,38,41	0.82	1 (6%)
5	5MC	a	1962	5	18,22,23	0.38	0	26,32,35	0.70	0
8	MEQ	d	150	8	8,9,10	0.45	0	5,10,12	1.33	0
5	PSU	a	746	31,5	18,21,22	1.02	1 (5%)	22,30,33	0.67	0
5	PSU	a	2457	5	18,21,22	1.03	1 (5%)	22,30,33	0.69	0
5	PSU	a	955	5	18,21,22	0.91	1 (5%)	22,30,33	0.65	0
5	OMC	a	2498	31,5	19,22,23	0.30	0	26,31,34	0.54	0
5	PSU	a	2605	5	18,21,22	0.98	1 (5%)	22,30,33	0.91	1 (4%)
5	6MZ	a	2030	5	18,25,26	0.72	0	16,36,39	0.90	1 (6%)
5	5MU	a	747	5	19,22,23	0.29	0	28,32,35	0.44	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the



Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	2MA	a	2503	31,32,5	-	2/3/25/26	0/3/3/3
5	5MU	a	1939	32,5	-	0/7/25/26	0/2/2/2
5	1MG	a	745	5	=	0/3/25/26	0/3/3/3
15	4D4	1	81	15	-	1/11/12/14	-
5	OMU	a	2552	5	ı	0/9/27/28	0/2/2/2
5	6MZ	a	1618	5	-	0/5/27/28	0/3/3/3
5	2MG	a	1835	5	-	0/5/27/28	0/3/3/3
5	PSU	a	2504	32,5	-	0/7/25/26	0/2/2/2
5	G7M	a	2069	5	-	2/3/25/26	0/3/3/3
5	PSU	a	2604	5	-	0/7/25/26	0/2/2/2
5	PSU	a	2580	32,5	=	0/7/25/26	0/2/2/2
5	H2U	a	2449	5	-	0/7/38/39	0/2/2/2
5	OMG	a	2251	32,5	-	0/5/27/28	0/3/3/3
5	2MG	a	2445	5	-	1/5/27/28	0/3/3/3
5	5MC	a	1962	5	ı	0/7/25/26	0/2/2/2
8	MEQ	d	150	8	-	2/8/9/11	-
5	PSU	a	746	31,5	=	1/7/25/26	0/2/2/2
5	PSU	a	2457	5	-	0/7/25/26	0/2/2/2
5	PSU	a	955	5	=	0/7/25/26	0/2/2/2
5	OMC	a	2498	31,5	-	0/9/27/28	0/2/2/2
5	PSU	a	2605	5	-	0/7/25/26	0/2/2/2
5	6MZ	a	2030	5	-	1/5/27/28	0/3/3/3
5	5MU	a	747	5	-	0/7/25/26	0/2/2/2

The worst 5 of 18 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$\operatorname{Observed}(\text{\AA})$	$Ideal(\AA)$
5	a	2580	PSU	C6-C5	4.09	1.40	1.35
5	a	2457	PSU	C6-C5	4.08	1.40	1.35
5	a	746	PSU	C6-C5	4.01	1.40	1.35
5	a	2605	PSU	C6-C5	3.85	1.39	1.35
5	a	2504	PSU	C6-C5	3.83	1.39	1.35

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
5	a	2503	2MA	CM2-C2-N1	3.20	123.34	116.23
5	a	2580	PSU	C3'-C2'-C1'	2.39	104.42	101.64
5	a	1618	6MZ	C2-N1-C6	2.30	118.56	116.59
5	a	2605	PSU	C2'-C3'-C4'	-2.17	98.42	102.64

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
5	a	2030	6MZ	C2-N1-C6	2.13	118.42	116.59

There are no chirality outliers.

5 of 10 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
8	d	150	MEQ	NE2-CD-CG-CB
8	d	150	MEQ	OE1-CD-CG-CB
5	a	2069	G7M	C4'-C5'-O5'-P
5	a	2030	6MZ	O4'-C4'-C5'-O5'
5	a	746	PSU	O4'-C1'-C5-C6

There are no ring outliers.

No monomer is involved in short contacts.

### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry (i)

Of 321 ligands modelled in this entry, 319 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
MIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
30	ACT	a	3001	-	3,3,3	0.95	0	3,3,3	0.78	0
33	6O1	a	3307	-	117,123,123	1.43	5 (4%)	155,191,191	1.81	20 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
33	6O1	a	3307	-	-	11/50/234/234	0/13/13/13

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$\operatorname{Observed}(\text{\AA})$	$\operatorname{Ideal}( ext{\AA})$
33	a	3307	6O1	O71-N65	12.52	1.49	1.22
33	a	3307	6O1	C04-C05	3.74	1.46	1.40
33	a	3307	6O1	O53-C49	3.69	1.43	1.40
33	a	3307	6O1	C56-C61	3.46	1.45	1.40
33	a	3307	6O1	O20-C16	2.58	1.44	1.41

The worst 5 of 20 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^o)$
33	a	3307	6O1	O50-C50-C51	-7.94	99.98	106.63
33	a	3307	6O1	C16-O20-C20	7.57	119.05	112.16
33	a	3307	6O1	C53-O53-C49	7.35	116.62	111.59
33	a	3307	6O1	O03-C03-C02	-7.20	109.33	120.05
33	a	3307	6O1	C10-O19-C19	6.67	128.15	114.66

There are no chirality outliers.

5 of 11 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
33	a	3307	601	C64-C63-O12-C12
33	a	3307	6O1	O67-C63-O12-C12
33	a	3307	6O1	C53-C52-O52-C55
33	a	3307	6O1	C18-C19-O19-C10
33	a	3307	6O1	C11-C10-O19-C19

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.



The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

### 5.7 Other polymers (i)

There are no such residues in this entry.

### 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.




### 6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-16530. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

### 6.1 Orthogonal projections (i)

#### 6.1.1 Primary map



#### 6.1.2 Raw map



The images above show the map projected in three orthogonal directions.



### 6.2 Central slices (i)

#### 6.2.1 Primary map



#### 6.2.2 Raw map



The images above show central slices of the map in three orthogonal directions.



### 6.3 Largest variance slices (i)

#### 6.3.1 Primary map



#### 6.3.2 Raw map




The images above show the largest variance slices of the map in three orthogonal directions.




### 6.4 Orthogonal standard-deviation projections (False-color) (i)

#### 6.4.1 Primary map



#### 6.4.2 Raw map



The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.




### 6.5 Orthogonal surface views (i)

#### 6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.0325. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

#### 6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.

### 6.6 Mask visualisation (i)


This section was not generated. No masks/segmentation were deposited.



## 7 Map analysis (i)

This section contains the results of statistical analysis of the map.


### 7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



### 7.2 Volume estimate (i)




The volume at the recommended contour level is  $338~\mathrm{nm}^3$ ; this corresponds to an approximate mass of  $305~\mathrm{kDa}$ .

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



### 7.3 Rotationally averaged power spectrum (i)




\*Reported resolution corresponds to spatial frequency of 0.538  $\rm \AA^{-1}$ 



### 8 Fourier-Shell correlation (i)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

### 8.1 FSC (i)



\*Reported resolution corresponds to spatial frequency of 0.538  $\rm \mathring{A}^{-1}$ 



### 8.2 Resolution estimates (i)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)			
resolution estimate (A)	0.143	0.5	Half-bit	
Reported by author	1.86	-	-	
Author-provided FSC curve	-	-	-	
Unmasked-calculated*	2.29	2.81	2.37	


<sup>\*</sup>Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 2.29 differs from the reported value 1.86 by more than 10 %



## 9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-16530 and PDB model 8CAM. Per-residue inclusion information can be found in section 3 on page 12.


### 9.1 Map-model overlay (i)



The images above show the 3D surface view of the map at the recommended contour level 0.0325 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.




### 9.2 Q-score mapped to coordinate model (i)



The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


### 9.3 Atom inclusion mapped to coordinate model (i)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.0325).



### 9.4 Atom inclusion (i)



At the recommended contour level, 88% of all backbone atoms, 86% of all non-hydrogen atoms, are inside the map.



### 9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (0.0325) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.8620	0.7310
0	0.6760	0.6480
1	0.9800	0.8210
2	0.9530	0.7890
3	0.8910	0.7350
a	0.8740	0.7300
b	0.9210	0.7250
С	0.9220	0.7830
d	0.8890	0.7610
е	0.8140	0.7490
g	0.4800	0.5680
h	0.5160	0.6270
i	0.9170	0.7710
j	0.8740	0.7660
k	0.8630	0.7600
1	0.8640	0.7310
m	0.9270	0.7450
О	0.8230	0.7220
p	0.9460	0.8080
q	0.8230	0.7460
r	0.8800	0.7520
S	0.7420	0.6710
t	0.7450	0.6850
u	0.7290	0.6550
V	0.8070	0.6900
W	0.8650	0.7560
X	0.6220	0.6460
У	0.8470	0.7380
Z	0.8500	0.7320



