



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 18, 2023 – 02:50 PM EDT

PDB ID : 5CCX
Title : Structure of the product complex of tRNA m1A58 methyltransferase with tRNA³Lys as substrate
Authors : Finer-Moore, J.; Czudnochowski, N.; O'Connell III, J.D.; Wang, A.L.; Stroud, R.M.
Deposited on : 2015-07-02
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION




The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)
RNA backbone	3102	1000 (2.54-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	289	
2	B	497	
3	N	77	

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 12320 atoms, of which 5155 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called tRNA (adenine(58)-N(1))-methyltransferase catalytic subunit TRMT61A.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	283	4337	1374	2156	394	402	11	0	2	0

- Molecule 2 is a protein called tRNA (adenine(58)-N(1))-methyltransferase non-catalytic subunit TRM6.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
2	B	371	5894	1871	2956	520	533	14	0	2	0

- Molecule 3 is a RNA chain called tRNA3Lys.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	P			
3	N	77	1734	763	24	306	561	80	0	6	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
N	74	C	-	insertion	GB 339572
N	75	C	-	insertion	GB 339572
N	76	A	-	insertion	GB 339572

- Molecule 4 is S-ADENOSYL-L-HOMOCYSTEINE (three-letter code: SAH) (formula: C₁₄H₂₀N₆O₅S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
4	A	1	Total	C	H	N	O	S	0	0
			45	14	19	6	5	1		

- Molecule 5 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	N	1	Total	Na	0	0
			1	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	132	Total	O	0	0
			132	132		
6	B	102	Total	O	0	0
			102	102		
6	N	75	Total	O	0	0
			75	75		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: tRNA (adenine(58)-N(1))-methyltransferase catalytic subunit TRMT61A



- Molecule 2: tRNA (adenine(58)-N(1))-methyltransferase non-catalytic subunit TRM6



- Molecule 3: tRNA³Lys



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	136.77Å 136.77Å 177.16Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	54.13 – 2.10 54.13 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.9 (54.13-2.10) 89.0 (54.13-2.10)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.78 (at 2.10Å)	Xtrriage
Refinement program	PHENIX 1.8.1_1168	Depositor
R, R_{free}	0.207 , 0.237 0.209 , 0.239	Depositor DCC
R_{free} test set	4964 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	39.5	Xtrriage
Anisotropy	0.124	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 52.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	12320	wwPDB-VP
Average B, all atoms (Å ²)	66.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.06% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SAH, NA, 1MA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.27	0/2240	0.47	0/3046
2	B	0.28	0/2999	0.46	0/4034
3	N	0.20	0/1858	0.73	0/2894
All	All	0.26	0/7097	0.55	0/9974

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2181	2156	2155	21	0
2	B	2938	2956	2955	51	0
3	N	1710	24	866	51	0
4	A	26	19	19	0	0
5	N	1	0	0	0	0
6	A	132	0	0	0	1
6	B	102	0	0	1	0
6	N	75	0	0	2	0
All	All	7165	5155	5995	121	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 10.

All (121) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:N:59[A]:G:H1'	3:N:60[A]:U:H5'	1.39	1.04
3:N:17:C:OP2	3:N:19:G:N2	2.01	0.93
3:N:75:C:H2'	3:N:75:C:O2	1.72	0.89
2:B:220:ARG:NH2	6:B:501:HOH:O	2.11	0.84
3:N:75:C:O2'	3:N:76:A:H5'	1.79	0.82
1:A:1:MET:HB3	1:A:2:SER:HA	1.64	0.80
3:N:8:U:H3	3:N:21:A:H2	1.28	0.79
3:N:76:A:N3	3:N:76:A:C5	2.44	0.79
3:N:76:A:N3	3:N:76:A:H3'	1.99	0.78
3:N:59[A]:G:H1'	3:N:60[A]:U:C5'	2.13	0.78
3:N:69:G:OP2	6:N:201:HOH:O	2.04	0.75
2:B:351[B]:GLU:OE2	2:B:354:ARG:NH1	2.20	0.74
2:B:384:LEU:HD12	2:B:410:CYS:SG	2.27	0.73
2:B:99:ASP:OD1	3:N:53:G:O2'	2.06	0.71
3:N:17:C:P	3:N:19:G:H22	2.17	0.68
3:N:17:C:P	3:N:19:G:N2	2.67	0.68
3:N:59[A]:G:C1'	3:N:60[A]:U:H5'	2.20	0.67
3:N:17:C:H3'	3:N:17:C:OP1	1.94	0.67
3:N:36:U:H2'	3:N:36:U:O2	1.95	0.66
3:N:76:A:N3	3:N:76:A:N9	2.44	0.65
3:N:75:C:O2	3:N:75:C:C2'	2.43	0.64
1:A:243:VAL:CG2	1:A:265:PHE:CE2	2.80	0.64
3:N:59[A]:G:C4'	3:N:60[A]:U:OP1	2.47	0.62
1:A:243:VAL:HG23	1:A:265:PHE:CZ	2.34	0.61
3:N:17:C:O2'	3:N:18:G:H5'	2.01	0.60
3:N:19:G:H2'	3:N:19:G:N3	2.16	0.60
3:N:0:G:O6	3:N:75:C:N4	2.35	0.60
3:N:34:U:OP2	6:N:202:HOH:O	2.15	0.59
3:N:73:G:O2'	3:N:75:C:H5	1.84	0.59
3:N:36:U:O2	3:N:36:U:C2'	2.51	0.58
1:A:243:VAL:CG2	1:A:265:PHE:CZ	2.86	0.58
2:B:46:THR:CG2	2:B:51:TRP:CH2	2.87	0.58
2:B:57:VAL:HG23	2:B:66:PHE:CZ	2.39	0.58
3:N:56:C:C5	3:N:58[A]:1MA:HM11	2.41	0.56
3:N:71:G:O2'	3:N:72:C:H5'	2.04	0.56
2:B:54:LEU:O	2:B:57:VAL:HG12	2.07	0.54
2:B:176:GLU:HB3	2:B:179:LYS:HG3	1.88	0.54
2:B:64:THR:HG22	2:B:66:PHE:CE2	2.43	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:N:57:A:H4'	3:N:58[B]:1MA:OP2	2.09	0.52
2:B:66:PHE:HA	2:B:76:PRO:HD3	1.92	0.51
2:B:343:ILE:O	2:B:347:GLN:HG2	2.10	0.51
1:A:24:ALA:O	1:A:25:MET:HG3	2.11	0.51
2:B:354:ARG:O	2:B:358:LEU:HD13	2.10	0.50
2:B:65:ALA:O	2:B:66:PHE:CD2	2.65	0.50
2:B:432:ASN:ND2	2:B:443:LYS:HE3	2.26	0.50
2:B:184:ARG:NH2	2:B:450:GLY:HA3	2.27	0.50
3:N:59[A]:G:H4'	3:N:60[A]:U:OP1	2.10	0.50
3:N:73:G:O2'	3:N:74:C:O5'	2.29	0.50
2:B:26:VAL:CG1	2:B:160:VAL:HG13	2.43	0.49
3:N:4:C:O2'	3:N:5:G:H5'	2.13	0.49
1:A:262:THR:O	1:A:262:THR:HG22	2.13	0.49
3:N:73:G:O2'	3:N:75:C:C5	2.65	0.49
2:B:46:THR:HG23	2:B:51:TRP:CZ3	2.47	0.49
3:N:15[A]:G:C6	3:N:19:G:C8	3.00	0.49
1:A:66:VAL:HG12	1:A:67:TYR:N	2.28	0.49
3:N:76:A:N3	3:N:76:A:C3'	2.72	0.49
2:B:114:LEU:HD22	2:B:114:LEU:N	2.29	0.48
1:A:66:VAL:CG1	1:A:67:TYR:N	2.78	0.47
2:B:405:GLU:HB2	2:B:406:PRO:HD3	1.97	0.47
3:N:17:C:C4	3:N:19:G:C6	3.02	0.47
1:A:1:MET:HG3	1:A:14:ASP:HA	1.97	0.47
2:B:26:VAL:CG1	2:B:27:VAL:N	2.77	0.47
2:B:68:VAL:HG12	2:B:69:THR:N	2.29	0.47
2:B:384:LEU:HD11	2:B:399:VAL:HG21	1.95	0.47
2:B:110:ASP:O	2:B:114:LEU:CD2	2.63	0.47
2:B:405:GLU:N	2:B:406:PRO:CD	2.77	0.47
3:N:59[A]:G:O4'	3:N:60[A]:U:OP1	2.33	0.46
2:B:115:LYS:O	2:B:118:GLY:N	2.48	0.46
3:N:59[A]:G:O2'	3:N:60[A]:U:H5'	2.14	0.46
3:N:66:U:C4	3:N:67:U:C4	3.03	0.46
2:B:207:GLU:HG3	2:B:209:CYS:H	1.80	0.46
1:A:1:MET:HG2	1:A:71:PRO:HG3	1.98	0.45
2:B:46:THR:CG2	2:B:51:TRP:CZ3	3.00	0.45
1:A:198:VAL:HG22	1:A:199:GLU:OE2	2.17	0.45
3:N:21:A:N3	3:N:21:A:O4'	2.50	0.44
2:B:184:ARG:HD2	2:B:186:ASP:OD1	2.17	0.44
3:N:54:U:O2'	3:N:55:U:H5'	2.18	0.44
3:N:73:G:H1'	3:N:75:C:H41	1.82	0.43
3:N:8:U:N3	3:N:21:A:H2	2.05	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:N:64:U:O2'	3:N:65:G:H5'	2.18	0.43
1:A:239:ASN:ND2	1:A:271:MET:HG2	2.34	0.43
2:B:29:LYS:HE3	2:B:31:GLU:O	2.19	0.43
1:A:83:ARG:HD3	1:A:143:LYS:HG2	2.01	0.42
2:B:52:PHE:CE1	2:B:68:VAL:CG2	3.02	0.42
3:N:42:G:O2'	3:N:43:A:H5'	2.18	0.42
2:B:65:ALA:C	2:B:66:PHE:CD2	2.93	0.42
2:B:108:GLN:O	2:B:111:ILE:HG22	2.18	0.42
2:B:184:ARG:CZ	2:B:450:GLY:HA3	2.49	0.42
2:B:346:LYS:O	2:B:350:GLN:HG3	2.19	0.42
1:A:134:VAL:HG21	1:A:169:PHE:HA	2.01	0.42
2:B:26:VAL:HG11	2:B:160:VAL:HG13	2.02	0.42
2:B:107:THR:HG22	2:B:108:GLN:N	2.35	0.42
2:B:158:ILE:N	2:B:158:ILE:CD1	2.83	0.42
3:N:20:U:O2	3:N:20:U:O4'	2.37	0.42
2:B:184:ARG:HB2	2:B:450:GLY:HA2	2.01	0.42
3:N:73:G:O2'	3:N:74:C:C5'	2.67	0.42
3:N:59[A]:G:H2'	3:N:59[A]:G:N3	2.35	0.42
2:B:157:ILE:C	2:B:158:ILE:HD12	2.40	0.42
3:N:39:U:H2'	3:N:40:C:C6	2.55	0.42
2:B:158:ILE:N	2:B:158:ILE:HD12	2.35	0.41
1:A:87:LEU:HD12	1:A:117:VAL:HG21	2.01	0.41
1:A:252:THR:HG22	1:A:253:GLY:N	2.35	0.41
2:B:29:LYS:HB3	2:B:159:THR:OG1	2.20	0.41
2:B:176:GLU:HB3	2:B:179:LYS:CG	2.50	0.41
3:N:67:U:H2'	3:N:68:C:C6	2.55	0.41
2:B:57:VAL:O	2:B:57:VAL:HG13	2.20	0.41
1:A:19:SER:HA	1:A:25:MET:HG3	2.03	0.41
3:N:59[A]:G:C2'	3:N:60[A]:U:H5'	2.51	0.41
1:A:108:CYS:HB3	1:A:169:PHE:CD1	2.55	0.41
2:B:106:LEU:C	2:B:106:LEU:HD12	2.41	0.41
2:B:384:LEU:HD13	2:B:414:LEU:HD11	2.01	0.41
3:N:1:G:O2'	3:N:76:A:N6	2.53	0.41
2:B:199:ALA:HB1	2:B:224:PHE:HB2	2.03	0.41
1:A:6:TYR:HB3	2:B:395:ARG:NH2	2.35	0.41
2:B:52:PHE:HE1	2:B:68:VAL:HG21	1.86	0.41
3:N:74:C:O5'	3:N:74:C:H6	2.04	0.41
1:A:83:ARG:CD	1:A:143:LYS:HG2	2.51	0.40
2:B:46:THR:CG2	2:B:51:TRP:CZ2	3.04	0.40
1:A:243:VAL:HG21	1:A:265:PHE:CE2	2.55	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the sym-

metry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:405:HOH:O	6:A:405:HOH:O[7_555]	2.14	0.06

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	281/289 (97%)	270 (96%)	11 (4%)	0	100	100
2	B	366/497 (74%)	349 (95%)	17 (5%)	0	100	100
All	All	647/786 (82%)	619 (96%)	28 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	236/238 (99%)	232 (98%)	4 (2%)	60	67
2	B	310/428 (72%)	309 (100%)	1 (0%)	92	95
All	All	546/666 (82%)	541 (99%)	5 (1%)	78	84

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	82	HIS
1	A	83	ARG
1	A	136	PHE
2	B	198	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
3	N	73/77 (94%)	12 (16%)	1 (1%)

All (12) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
3	N	18	G
3	N	19	G
3	N	21	A
3	N	37	A
3	N	48	C
3	N	56	C
3	N	57	A
3	N	59[A]	G
3	N	60[A]	U
3	N	73	G
3	N	74	C
3	N	76	A

All (1) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
3	N	59[A]	G

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	1MA	N	58[A]	3	16,25,26	1.38	3 (18%)	18,37,40	1.27	2 (11%)
3	1MA	N	58[B]	3	16,25,26	1.32	3 (18%)	18,37,40	1.29	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	1MA	N	58[A]	3	-	2/3/25/26	0/3/3/3
3	1MA	N	58[B]	3	-	3/3/25/26	0/3/3/3

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	N	58[A]	1MA	C2-N3	3.17	1.32	1.29
3	N	58[B]	1MA	C2-N3	2.84	1.32	1.29
3	N	58[B]	1MA	C5-C4	-2.69	1.36	1.43
3	N	58[A]	1MA	C5-C4	-2.61	1.36	1.43
3	N	58[A]	1MA	C6-N6	2.52	1.33	1.27
3	N	58[B]	1MA	C6-N6	2.51	1.33	1.27

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	N	58[B]	1MA	N1-C2-N3	-4.29	121.02	126.02
3	N	58[A]	1MA	N1-C2-N3	-4.09	121.25	126.02
3	N	58[B]	1MA	C5-C6-N1	2.71	117.94	113.90
3	N	58[A]	1MA	C5-C6-N1	2.63	117.82	113.90

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	N	58[B]	1MA	O4'-C4'-C5'-O5'

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	N	58[B]	1MA	C3'-C4'-C5'-O5'
3	N	58[A]	1MA	C4'-C5'-O5'-P
3	N	58[B]	1MA	C4'-C5'-O5'-P
3	N	58[A]	1MA	O4'-C4'-C5'-O5'

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	N	58[A]	1MA	1	0
3	N	58[B]	1MA	1	0

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	SAH	A	301	-	24,28,28	1.13	3 (12%)	25,40,40	1.60	4 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	SAH	A	301	-	-	0/11/31/31	0/3/3/3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	301	SAH	C2-N3	3.55	1.37	1.32
4	A	301	SAH	C2-N1	2.23	1.38	1.33
4	A	301	SAH	OXT-C	-2.02	1.23	1.30

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	301	SAH	N3-C2-N1	-5.36	120.30	128.68
4	A	301	SAH	C5'-SD-CG	-3.27	92.46	102.27
4	A	301	SAH	OXT-C-O	-2.37	118.71	124.09
4	A	301	SAH	OXT-C-CA	2.10	120.53	113.38


There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	283/289 (97%)	0.21	8 (2%) 53 59	32, 44, 80, 115	0
2	B	371/497 (74%)	0.82	69 (18%) 1 1	36, 60, 117, 152	0
3	N	76/77 (98%)	-0.49	3 (3%) 39 45	41, 59, 122, 160	1 (1%)
All	All	730/863 (84%)	0.44	80 (10%) 5 7	32, 54, 110, 160	1 (0%)

All (80) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1	MET	10.1
2	B	342	TYR	9.1
2	B	42	ARG	6.8
2	B	53	TYR	6.5
1	A	260	SER	6.4
2	B	341	ASP	5.4
2	B	43	LYS	5.3
2	B	52	PHE	5.0
2	B	463	ASN	4.9
2	B	54	LEU	4.8
2	B	57	VAL	4.8
2	B	51	TRP	4.8
2	B	343	ILE	4.7
2	B	80	ARG	4.6
2	B	346	LYS	4.6
1	A	253	GLY	4.4
2	B	119	ILE	4.4
2	B	120	LYS	4.3
2	B	348	ARG	4.2
2	B	111	ILE	4.1
2	B	74	LEU	4.1
2	B	349	ARG	4.0
2	B	351[A]	GLU	3.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	117	LYS	3.9
2	B	347	GLN	3.7
2	B	70	SER	3.7
3	N	18	G	3.7
2	B	124	ILE	3.6
2	B	271	SER	3.5
2	B	66	PHE	3.4
2	B	69	THR	3.3
2	B	345	GLU	3.3
2	B	40	GLN	3.3
2	B	113	ALA	3.3
2	B	344	GLN	3.2
2	B	354	ARG	3.1
2	B	39	VAL	3.1
2	B	350	GLN	3.0
2	B	462	ASP	3.0
2	B	268	GLY	3.0
2	B	71	GLY	3.0
2	B	102	LYS	2.9
2	B	122	GLU	2.9
2	B	429	TRP	2.9
2	B	44	LYS	2.9
2	B	79	LYS	2.9
2	B	88	LYS	2.9
2	B	41	ARG	2.9
3	N	76	A	2.9
2	B	266	LEU	2.9
2	B	234	GLY	2.8
2	B	358	LEU	2.8
2	B	49	LYS	2.8
2	B	18[A]	HIS	2.8
2	B	59	GLY	2.6
2	B	108	GLN	2.6
2	B	353	GLN	2.6
1	A	63	GLY	2.6
2	B	270	PHE	2.6
2	B	89	GLU	2.5
2	B	112	LYS	2.5
2	B	114	LEU	2.5
2	B	355	LYS	2.5
1	A	261	ASP	2.4
2	B	352	GLU	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	115	LYS	2.4
2	B	428	THR	2.3
1	A	233	VAL	2.3
2	B	68	VAL	2.3
2	B	56	ASN	2.3
1	A	235	PRO	2.3
3	N	75	C	2.3
2	B	105	LYS	2.3
2	B	116	ASP	2.3
2	B	136	ARG	2.3
1	A	251	GLY	2.2
2	B	55	ASP	2.2
2	B	126	GLN	2.2
2	B	19	ARG	2.1
2	B	118	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	1MA	N	58[A]	23/24	0.94	0.19	46,57,68,69	35
3	1MA	N	58[B]	23/24	0.94	0.19	42,53,69,69	35

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	SAH	A	301	26/26	0.96	0.13	32,40,46,55	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	NA	N	101	1/1	0.98	0.04	42,42,42,42	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.