



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 4, 2024 – 05:44 am GMT

PDB ID : 5CDP  
Title : 2.45Å structure of etoposide with S.aureus DNA gyrase and DNA  
Authors : Bax, B.D.; Srikannathasan, V.; Chan, P.F.  
Deposited on : 2015-07-04  
Resolution : 2.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1544 (2.48-2.44)
Clashscore	141614	1613 (2.48-2.44)
Ramachandran outliers	138981	1598 (2.48-2.44)
Sidechain outliers	138945	1598 (2.48-2.44)
RSRZ outliers	127900	1523 (2.48-2.44)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	483	94% (green), 6% (yellow)
1	C	483	95% (green), . (grey)
2	B	190	2% (red), 94% (green), 6% (yellow)
2	D	190	5% (red), 91% (green), 8% (yellow), . (grey)
3	E	8	75% (green), 25% (yellow)

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	F	8	 100%
4	G	13	 77% 15% 8%
4	H	13	 69% 23% 8%

## 2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 12642 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA gyrase subunit A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	483	Total	C	N	O	S	0	14	0
			3957	2454	726	760	17			
1	C	481	Total	C	N	O	S	0	10	0
			3906	2420	722	748	16			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	123	PHE	TYR	engineered mutation	UNP Q99XG5
C	123	PHE	TYR	engineered mutation	UNP Q99XG5

- Molecule 2 is a protein called DNA gyrase subunit B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	190	Total	C	N	O	S	0	8	0
			1539	966	266	298	9			
2	D	189	Total	C	N	O	S	0	15	0
			1596	996	275	316	9			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	544	THR	-	linker	UNP P66937
B	545	GLY	-	linker	UNP P66937
D	544	THR	-	linker	UNP P66937
D	545	GLY	-	linker	UNP P66937

- Molecule 3 is a DNA chain called DNA (5'-D(\*AP\*GP\*CP\*CP\*GP\*TP\*AP\*G\*GP\*GP\*TP\*AP\*CP\*CP\*TP\*AP\*CP\*GP\*GP\*CP\*T)-3').

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
3	E	8	Total	C	N	O	P	0	0	0
			163	78	33	45	7			
3	F	8	Total	C	N	O	P	0	0	0
			163	78	33	45	7			

- Molecule 4 is a DNA chain called DNA (5'-D(\*AP\*GP\*CP\*CP\*GP\*TP\*AP\*G\*GP\*GP\*TP\*AP\*CP\*CP\*TP\*AP\*CP\*GP\*GP\*CP\*T)-3').

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
4	G	12	Total	C	N	O	P	0	0	0
			244	116	43	73	12			
4	H	12	Total	C	N	O	P	0	2	0
			256	118	46	78	14			


- Molecule 5 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Na	0	0
			1	1		
5	H	1	Total	Na	0	0
			1	1		

- Molecule 6 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	1	Total	Mn	0	0
			1	1		
6	D	1	Total	Mn	0	1
			2	2		
6	G	1	Total	Mn	0	0
			1	1		

- Molecule 7 is (5S,5aR,8aR,9R)-9-(4-hydroxy-3,5-dimethoxyphenyl)-8-oxo-5,5a,6,8,8a,9-hexahydrofuro[3',4':6,7]naphtho[2,3-d][1,3]dioxol-5-yl 4,6-O-[(1R)-ethylidene]-beta-D-glucopyranoside (three-letter code: EVP) (formula: C<sub>29</sub>H<sub>32</sub>O<sub>13</sub>).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	H	1	Total	C O	0	0
			42	29 13		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	299	Total	O	0	0
			299	299		
8	B	61	Total	O	0	0
			61	61		
8	C	298	Total	O	0	1
			299	299		
8	D	33	Total	O	0	2
			35	35		
8	E	17	Total	O	0	0
			17	17		
8	G	24	Total	O	0	0
			24	24		
8	F	13	Total	O	0	0
			13	13		
8	H	22	Total	O	0	0
			22	22		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA gyrase subunit A




- Molecule 1: DNA gyrase subunit A



- Molecule 2: DNA gyrase subunit B



- Molecule 2: DNA gyrase subunit B



- Molecule 3: DNA (5'-D(\*AP\*GP\*CP\*CP\*GP\*TP\*AP\*G\*GP\*GP\*TP\*AP\*CP\*CP\*TP\*AP\*CP\*GP\*GP\*CP\*T)-3')






- Molecule 3: DNA (5'-D(\*AP\*GP\*CP\*CP\*GP\*TP\*AP\*G\*GP\*GP\*TP\*AP\*CP\*CP\*TP\*AP\*CP\*GP\*GP\*CP\*T)-3')

Chain F:  100%

There are no outlier residues recorded for this chain.

- Molecule 4: DNA (5'-D(\*AP\*GP\*CP\*CP\*GP\*TP\*AP\*G\*GP\*GP\*TP\*AP\*CP\*CP\*TP\*AP\*CP\*GP\*GP\*CP\*T)-3')

Chain G:  77% 15% 8%



- Molecule 4: DNA (5'-D(\*AP\*GP\*CP\*CP\*GP\*TP\*AP\*G\*GP\*GP\*TP\*AP\*CP\*CP\*TP\*AP\*CP\*GP\*GP\*CP\*T)-3')

Chain H:  69% 23% 8%





## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	93.36Å 93.36Å 411.24Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	39.78 – 2.45 39.78 – 2.45	Depositor EDS
% Data completeness (in resolution range)	95.9 (39.78-2.45) 96.0 (39.78-2.45)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.38 (at 2.45Å)	Xtrriage
Refinement program	BUSTER-TNT BUSTER 2.11.5	Depositor
R, $R_{free}$	0.177 , 0.211 0.176 , 0.211	Depositor DCC
$R_{free}$ test set	2860 reflections (4.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	32.2	Xtrriage
Anisotropy	0.146	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 51.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.056 for h,-h-k,-l	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	12642	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	37.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.55% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, EVP, MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.43	0/4008	0.63	0/5391
1	C	0.42	0/3954	0.62	0/5325
2	B	0.42	0/1570	0.66	0/2119
2	D	0.42	0/1623	0.64	0/2192
3	E	1.03	0/183	0.96	0/281
3	F	1.00	0/183	0.99	0/281
4	G	1.14	0/272	0.99	0/417
4	H	1.10	0/284	1.13	0/434
All	All	0.50	0/12077	0.68	0/16440

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3957	0	4004	18	0
1	C	3906	0	3938	11	0
2	B	1539	0	1507	5	0
2	D	1596	0	1548	10	0
3	E	163	0	91	3	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	F	163	0	90	0	0
4	G	244	0	136	3	0
4	H	256	0	134	3	0
5	A	1	0	0	0	0
5	H	1	0	0	0	0
6	B	1	0	0	0	0
6	D	2	0	0	0	0
6	G	1	0	0	0	0
7	H	42	0	32	1	0
8	A	299	0	0	1	0
8	B	61	0	0	0	0
8	C	299	0	0	0	0
8	D	35	0	0	1	0
8	E	17	0	0	0	0
8	F	13	0	0	0	0
8	G	24	0	0	0	0
8	H	22	0	0	0	0
All	All	12642	0	11480	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:447:ARG:HD3	2:D:454:ILE:HD11	1.59	0.85
2:D:443:THR:HG22	2:D:454:ILE:HG12	1.67	0.76
2:D:583:LEU:HD12	8:D:804:HOH:O	1.91	0.71
1:A:232[A]:ARG:HH11	1:A:232[A]:ARG:CG	2.05	0.70
1:A:232[A]:ARG:HH11	1:A:232[A]:ARG:HG3	1.62	0.63
2:D:447:ARG:HD3	2:D:454:ILE:CD1	2.32	0.58
2:D:458[A]:ARG:HG2	4:H:2010:DT:H5'	1.84	0.58
2:B:459:GLY:HA2	3:E:8:DG:H2''	1.90	0.53
1:A:232[A]:ARG:HG3	1:A:232[A]:ARG:NH1	2.21	0.53
2:D:626:VAL:HG11	4:G:2017:DG:H3'	1.90	0.52
2:D:458[B]:ARG:HG3	4:H:2009[B]:DG:H2''	1.91	0.52
2:B:626:VAL:HG11	4:H:2017:DG:H3'	1.91	0.52
1:A:64:TYR:HB3	1:A:125:GLU:HB3	1.92	0.52
1:C:64:TYR:HB3	1:C:125:GLU:HB3	1.92	0.52
1:C:225:LEU:HD21	1:C:244:ARG:HD2	1.92	0.51
1:A:225:LEU:HD21	1:A:244:ARG:HD2	1.90	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:387:ALA:HA	1:C:394:ILE:HG13	1.94	0.49
1:A:391:ILE:O	1:A:395:ILE:HG12	2.12	0.49
1:C:122:ARG:HH21	4:G:2009:DG:H5''	1.77	0.49
1:A:404:ASP:OD2	1:C:431:ARG:NH2	2.46	0.47
1:C:391:ILE:O	1:C:395:ILE:HG12	2.13	0.47
1:C:388:LEU:HD13	1:C:438:ARG:HG2	1.98	0.46
1:A:377:ARG:HG3	1:A:448:LEU:HD11	1.98	0.46
1:A:58:MET:HE3	8:A:770:HOH:O	2.17	0.45
1:A:387:ALA:HA	1:A:394:ILE:HG13	1.99	0.44
1:C:234:TYR:O	1:C:347:ASN:HB2	2.18	0.43
2:D:444:LYS:HG3	2:D:454:ILE:HD13	2.00	0.43
2:D:502:LYS:HG2	2:D:538:TYR:CE1	2.54	0.43
2:B:459:GLY:CA	3:E:8:DG:H2''	2.48	0.43
1:A:76:GLY:HA3	1:C:69:ARG:HD2	2.01	0.43
4:G:2009:DG:N3	7:H:2101:EVP:H21A	2.35	0.42
1:A:232[A]:ARG:CG	1:A:232[A]:ARG:NH1	2.72	0.42
2:B:543:PRO:HB2	2:B:580[A]:TYR:OH	2.20	0.42
1:A:234:TYR:O	1:A:347:ASN:HB2	2.19	0.42
1:A:272:ARG:CZ	3:E:3:DC:H5''	2.50	0.41
1:A:381:LEU:HD12	1:A:448:LEU:HD12	2.03	0.41
1:A:365:ARG:HA	1:A:368:GLN:HE21	1.85	0.41
1:C:365:ARG:HA	1:C:368:GLN:HE21	1.86	0.40
1:A:100:ARG:HG3	1:A:101:TYR:CE2	2.56	0.40
2:B:430:GLU:HG2	2:B:502:LYS:HB2	2.03	0.40
1:A:109:ASN:HB3	1:A:119:ALA:HB2	2.03	0.40
1:C:100:ARG:HG3	1:C:101:TYR:CE2	2.57	0.40
2:D:430:GLU:HG2	2:D:502:LYS:HB2	2.04	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	495/483 (102%)	483 (98%)	12 (2%)	0	100	100
1	C	489/483 (101%)	479 (98%)	10 (2%)	0	100	100
2	B	196/190 (103%)	194 (99%)	2 (1%)	0	100	100
2	D	202/190 (106%)	192 (95%)	10 (5%)	0	100	100
All	All	1382/1346 (103%)	1348 (98%)	34 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	430/417 (103%)	421 (98%)	9 (2%)	53	66
1	C	422/417 (101%)	418 (99%)	4 (1%)	78	86
2	B	161/158 (102%)	155 (96%)	6 (4%)	34	45
2	D	169/158 (107%)	163 (96%)	6 (4%)	35	46
All	All	1182/1150 (103%)	1157 (98%)	25 (2%)	53	66

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	62	LYS
1	A	122	ARG
1	A	208	GLU
1	A	232[A]	ARG
1	A	232[B]	ARG
1	A	285	LYS
1	A	299	ARG
1	A	404	ASP
1	A	453	SER
2	B	427	GLU
2	B	442	SER
2	B	445	SER
2	B	590	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	B	595	THR
2	B	597	ASN
1	C	122	ARG
1	C	232	ARG
1	C	285	LYS
1	C	298	LEU
2	D	445	SER
2	D	474	ASN
2	D	501	HIS
2	D	581	LYS
2	D	590	GLN
2	D	595	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	324	GLN
1	A	368	GLN
2	B	541	GLN
2	B	597	ASN
2	B	605	GLN
1	C	324	GLN
1	C	368	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 6 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
7	EVP	H	2101	-	48,48,48	0.78	1 (2%)	72,73,73	1.68	12 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	EVP	H	2101	-	-	0/12/76/76	0/7/7/7

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	H	2101	EVP	C6-C9	-2.20	1.50	1.54

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	H	2101	EVP	C20-O5-C16	-5.19	109.70	117.53
7	H	2101	EVP	C21-O7-C18	-4.73	110.39	117.53
7	H	2101	EVP	C1-O1-C13	-4.49	99.51	105.34
7	H	2101	EVP	C1-O2-C2	-4.24	99.83	105.34
7	H	2101	EVP	O7-C18-C17	3.60	118.17	114.54
7	H	2101	EVP	O2-C1-O1	3.25	113.28	108.08
7	H	2101	EVP	C11-C4-C5	3.13	122.67	114.40
7	H	2101	EVP	C22-O8-C10	-3.02	110.08	114.99
7	H	2101	EVP	O5-C16-C17	2.89	117.45	114.54
7	H	2101	EVP	O3-C7-O4	2.72	124.24	121.42
7	H	2101	EVP	O1-C13-C2	2.56	112.69	109.78
7	H	2101	EVP	O4-C7-C6	-2.28	126.45	129.38

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	H	2101	EVP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.



## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	483/483 (100%)	-0.64	2 (0%) 92 93	19, 31, 50, 80	0
1	C	481/483 (99%)	-0.66	0 100 100	20, 31, 48, 81	0
2	B	190/190 (100%)	-0.31	4 (2%) 63 60	23, 40, 64, 100	0
2	D	189/190 (99%)	0.02	10 (5%) 26 23	31, 48, 74, 88	0
3	E	8/8 (100%)	-0.71	0 100 100	27, 30, 45, 47	0
3	F	8/8 (100%)	-0.33	0 100 100	32, 37, 51, 57	0
4	G	12/13 (92%)	-0.49	0 100 100	31, 39, 61, 63	0
4	H	12/13 (92%)	-0.28	0 100 100	25, 50, 66, 68	0
All	All	1383/1388 (99%)	-0.50	16 (1%) 79 77	19, 34, 58, 100	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	580[A]	TYR	6.9
2	B	545[A]	GLY	3.9
2	D	544	THR	3.4
2	B	640	ALA	3.4
2	D	491	GLY	2.6
2	D	473	LEU	2.6
2	D	476[A]	ASN	2.6
2	D	486	GLY	2.5
2	D	490	GLY	2.5
1	A	490[A]	LEU	2.5
2	B	544[A]	THR	2.2
2	D	580	TYR	2.1
1	A	491	GLY	2.1
2	D	638	VAL	2.1
2	D	493	PHE	2.0
2	D	495	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
7	EVP	H	2101	42/42	0.92	0.14	42,48,50,51	0
5	NA	H	2102	1/1	0.96	0.18	28,28,28,28	0
6	MN	D	701[B]	1/1	0.97	0.11	37,37,37,37	1
6	MN	D	701[A]	1/1	0.97	0.11	24,24,24,24	1
5	NA	A	501	1/1	0.98	0.05	33,33,33,33	0
6	MN	G	2101	1/1	0.99	0.17	54,54,54,54	0
6	MN	B	1001	1/1	1.00	0.07	36,36,36,36	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [i](#)

There are no such residues in this entry.