



wwPDB X-ray Structure Validation Summary Report ⓘ

May 19, 2020 – 05:44 pm BST

PDB ID : 2CF2
Title : Architecture of mammalian fatty acid synthase
Authors : Maier, T.; Jenni, S.; Ban, N.
Deposited on : 2006-02-14
Resolution : 4.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3146 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called FATTY ACID SYNTHASE, KS DOMAIN.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
1	A	406	Total C 406 406	0	0	406
1	J	406	Total C 406 406	0	0	406

- Molecule 2 is a protein called FATTY ACID SYNTHASE, MAT DOMAIN.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
2	B	304	Total C 304 304	0	0	304
2	K	304	Total C 304 304	0	0	304

- Molecule 3 is a protein called FATTY ACID SYNTHASE, DH DOMAIN.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
3	C	342	Total C 342 342	0	0	342
3	L	342	Total C 342 342	0	0	342

- Molecule 4 is a protein called FATTY ACID SYNTHASE, ER DOMAIN.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
4	D	295	Total C 295 295	0	0	295
4	M	295	Total C 295 295	0	0	295

- Molecule 5 is a protein called FATTY ACID SYNTHASE, KR DOMAIN.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	Trace
5	E	226	Total 226	C 226	0	0	226
5	N	226	Total 226	C 226	0	0	226


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: FATTY ACID SYNTHASE, KS DOMAIN

Chain A:  98%



- Molecule 1: FATTY ACID SYNTHASE, KS DOMAIN

Chain J:  98%



- Molecule 2: FATTY ACID SYNTHASE, MAT DOMAIN

Chain B:  99%



- Molecule 2: FATTY ACID SYNTHASE, MAT DOMAIN

Chain K:  99%




- Molecule 3: FATTY ACID SYNTHASE, DH DOMAIN

Chain C:  92% 8%




- Molecule 3: FATTY ACID SYNTHASE, DH DOMAIN

Chain L:  91% 9%




- Molecule 4: FATTY ACID SYNTHASE, ER DOMAIN

Chain D:  98%



- Molecule 4: FATTY ACID SYNTHASE, ER DOMAIN

Chain M:  99%



- Molecule 5: FATTY ACID SYNTHASE, KR DOMAIN

Chain E:  100%



- Molecule 5: FATTY ACID SYNTHASE, KR DOMAIN

Chain N:  100%



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	96.10Å 245.00Å 134.10Å 90.00° 101.20° 90.00°	Depositor
Resolution (Å)	50.00 – 4.30	Depositor
% Data completeness (in resolution range)	(Not available) (50.00-4.30)	Depositor
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
Refinement program		Depositor
R, R_{free}	(Not available) , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3146	wwPDB-VP
Average B, all atoms (Å ²)	60.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

There are no protein, RNA or DNA chains available to summarize Z scores of covalent bonds and angles.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	406	0	0	10	0
1	J	406	0	0	8	0
2	B	304	0	0	1	0
2	K	304	0	0	1	0
3	C	342	0	0	26	0
3	L	342	0	0	30	0
4	D	295	0	0	4	0
4	M	295	0	0	2	0
5	E	226	0	0	1	0
5	N	226	0	0	1	0
All	All	3146	0	0	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

The worst 5 of 49 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:121:ARG:CA	3:C:1005:GLU:CA	1.75	1.60
1:A:43:GLY:CA	3:L:1020:LEU:CA	1.83	1.55
1:J:126:LEU:CA	3:L:1014:ALA:CA	1.77	1.53
3:L:2137:ARG:CA	4:M:52:ARG:CA	1.87	1.50
3:C:1047:GLY:CA	3:L:1127:ARG:CA	1.92	1.47

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

There are no protein backbone outliers to report in this entry.

5.3.2 Protein sidechains [i](#)

There are no protein residues with a non-rotameric sidechain to report in this entry.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.