

Full wwPDB X-ray Structure Validation Report (i)

Jun 14, 2020 - 04:07 am BST

PDB ID	:	1CGD
Title	:	HYDRATION STRUCTURE OF A COLLAGEN PEPTIDE
Authors	:	Bella, J.; Brodsky, B.; Berman, H.M.
Deposited on		
Resolution	:	1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
Clashscore	141614	2625 (1.86-1.86)
Ramachandran outliers	138981	2592(1.86-1.86)
Sidechain outliers	138945	2592 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain	
1	А	30	83%	17%
1	В	30	87%	13%
1	С	30	67% 30%	•



2 Entry composition (i)


There are 3 unique types of molecules in this entry. The entry contains 745 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	Δ	30	Total	С	Ν	Ο	0	0	0
	Л	50	192	121	30	41			
1	В	30	Total	С	Ν	Ο	0	0	Ο
	I D	50	192	121	30	41	0	0	U
1	С	30	Total	С	Ν	Ο	0	0	0
		C 30	192	121	30	41	0	U	U

• Molecule 1 is a protein called COLLAGEN-LIKE PEPTIDE.

• Molecule 2 is ACETIC ACID (three-letter code: ACY) (formula: $C_2H_4O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 4 2 2 \end{array}$	0	0
2	А	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 4 2 2 \end{array}$	0	0
2	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0

Continued on next page...



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 4 2 2 \end{array}$	0	0
2	В	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 4 2 2 \end{array}$	0	0
2	С	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 4 2 2 \end{array}$	0	0
2	С	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 4 2 2 \end{array}$	0	0

• Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	53	Total O 53 53	0	0
3	В	41	Total O 41 41	0	0
3	С	47	TotalO4747	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: COLLAGEN-LIKE PEPTIDE

Chain A:	83%	17%
P1 P13 P14 A15 P28 P29 G30		
• Molecule 1:	COLLAGEN-LIKE PEPTIDE	
Chain B:	87%	13%
P31 P32 P50 P59 G60		
• Molecule 1:	COLLAGEN-LIKE PEPTIDE	
Chain C:	67% 30	% •
P61 P62 663 P64 P71 A75 A75	P10 P10 P10 P10 P10 P10 P10 P10 P10 P10	



4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants	173.46Å 14.06 Å 25.31 Å	Depositor
a, b, c, α , β , γ	90.00° 95.82° 90.00°	Depositor
Resolution (Å)	12.50 - 1.85	Depositor
% Data completeness	72.1 (12.50-1.85)	Depositor
(in resolution range)	72.1 (12.00-1.00)	Depositor
R_{merge}	(Not available)	Depositor
R _{sym}	(Not available)	Depositor
Refinement program	X-PLOR	Depositor
R, R_{free}	0.172 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	745	wwPDB-VP
Average B, all atoms $(Å^2)$	10.0	wwPDB-VP



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: HYP, ACY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
	Cham	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	0.59	0/111	0.80	1/138~(0.7%)	
1	В	0.58	0/111	0.90	1/138~(0.7%)	
1	С	0.59	0/111	0.85	1/138~(0.7%)	
All	All	0.59	0/333	0.85	3/414~(0.7%)	

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
1	А	1	PRO	CA-N-CD	-5.69	103.53	111.50
1	С	61	PRO	CA-N-CD	-5.44	103.89	111.50
1	В	31	PRO	CA-N-CD	-5.34	104.02	111.50

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	192	0	174	4	0
1	В	192	0	172	5	0
1	С	192	0	172	13	0
2	А	16	0	12	0	0
2	В	4	0	3	0	0

Continued on next page...



Mol	ě	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	С	8	0	6	1	0
3	А	53	0	0	1	0
3	В	41	0	0	0	0
3	С	47	0	0	2	0
All	All	745	0	539	14	0

Continued from previous page...

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (14) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:32:HYP:HB3	1:C:63:GLY:HA2	1.51	0.92
1:C:89:HYP:HD23	3:C:169:HOH:O	1.85	0.76
1:C:65:HYP:HB3	2:C:301:ACY:O	1.92	0.69
1:B:32:HYP:HA	1:C:61:PRO:HD3	1.86	0.58
1:C:61:PRO:HA	1:C:62:HYP:C	2.36	0.55
1:A:28:PRO:HG2	1:C:89:HYP:HB2	1.93	0.51
1:A:28:PRO:CG	1:C:89:HYP:HB2	2.43	0.48
1:B:50:HYP:HA	1:C:79:PRO:O	2.12	0.48
1:B:59:HYP:HA	1:C:88:PRO:HG2	1.96	0.47
3:A:121:HOH:O	1:C:77:HYP:HB2	2.17	0.44
1:A:15:ALA:HB2	1:C:75:ALA:HB3	2.00	0.43
1:B:32:HYP:HB2	1:C:62:HYP:O	2.19	0.42
1:A:13:PRO:HA	1:A:14:HYP:HD23	1.93	0.41
1:C:71:HYP:HD23	3:C:124:HOH:O	2.20	0.41

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	18/30~(60%)	18~(100%)	0	0	100	100
1	В	18/30~(60%)	18 (100%)	0	0	100	100
1	С	18/30~(60%)	18 (100%)	0	0	100	100
All	All	54/90~(60%)	54 (100%)	0	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	10/10~(100%)	10~(100%)	0	100 100
1	В	10/10~(100%)	10~(100%)	0	100 100
1	С	10/10~(100%)	10 (100%)	0	100 100
All	All	30/30~(100%)	30~(100%)	0	100 100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

30 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the



Mal	T	Chain	Dag	T : 1-	B	ond leng	gths	B	ond ang	gles
Mol	Type	Chain	\mathbf{Res}	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	HYP	С	83	1	6,8,9	0.72	0	5,10,12	0.78	0
1	HYP	В	53	1	6,8,9	0.59	0	5,10,12	0.79	0
1	HYP	С	86	1	6,8,9	0.49	0	5,10,12	0.80	0
1	HYP	А	5	1	6,8,9	0.68	0	5,10,12	0.80	0
1	HYP	С	89	1	6,8,9	0.52	0	5,10,12	0.87	0
1	HYP	А	11	1	6,8,9	0.55	0	5,10,12	0.89	0
1	HYP	В	32	1	6,8,9	0.46	0	5,10,12	1.02	0
1	HYP	В	44	1	6,8,9	0.49	0	5,10,12	1.48	1 (20%)
1	HYP	С	65	1	6,8,9	0.56	0	5,10,12	0.70	0
1	HYP	А	29	1	6,8,9	0.46	0	5,10,12	0.91	0
1	HYP	А	17	1	6,8,9	0.62	0	5,10,12	0.86	0
1	HYP	С	80	1	6,8,9	0.61	0	5,10,12	0.77	0
1	HYP	А	2	1	6,8,9	0.48	0	5,10,12	0.93	0
1	HYP	С	62	1	6,8,9	0.52	0	5,10,12	0.91	0
1	HYP	В	50	1	6,8,9	0.57	0	5,10,12	1.06	0
1	HYP	В	38	1	6,8,9	0.45	0	5,10,12	1.09	0
1	HYP	С	68	1	6,8,9	0.52	0	5,10,12	0.97	0
1	HYP	А	26	1	6,8,9	0.50	0	5,10,12	0.95	0
1	HYP	А	20	1	6,8,9	0.54	0	5,10,12	1.03	0
1	HYP	В	59	1	6,8,9	0.57	0	5,10,12	1.07	0
1	HYP	С	74	1	6,8,9	0.54	0	5,10,12	0.76	0
1	HYP	А	14	1	6,8,9	0.60	0	5,10,12	0.74	0
1	HYP	А	8	1	6,8,9	0.70	0	$5,\!10,\!12$	0.94	0
1	HYP	В	47	1	6,8,9	0.49	0	5,10,12	0.91	0
1	HYP	С	71	1	6,8,9	0.44	0	$5,\!10,\!12$	0.86	0
1	HYP	С	77	1	6,8,9	0.53	0	$5,\!10,\!12$	1.17	0
1	HYP	В	35	1	6,8,9	0.62	0	$5,\!10,\!12$	0.75	0
1	HYP	А	23	1	6,8,9	0.67	0	$5,\!10,\!12$	0.84	0
1	HYP	В	56	1	6, 8, 9	0.53	0	$5,\!10,\!12$	0.76	0
1	HYP	В	41	1	6,8,9	0.35	0	5,10,12	1.03	0

expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	HYP	С	83	1	-	0/0/11/13	0/1/1/1
1	HYP	В	53	1	-	0/0/11/13	0/1/1/1
1	HYP	С	86	1	-	0/0/11/13	0/1/1/1

Continued on next page...



Conti	nued from	m previou	ıs page				
Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	HYP	А	5	1	-	0/0/11/13	0/1/1/1
1	HYP	С	89	1	-	0/0/11/13	0/1/1/1
1	HYP	А	11	1	_	0/0/11/13	0/1/1/1
1	HYP	В	32	1	-	0/0/11/13	0/1/1/1
1	HYP	В	44	1	-	0/0/11/13	0/1/1/1
1	HYP	С	65	1	_	0/0/11/13	0/1/1/1
1	HYP	А	29	1	-	0/0/11/13	0/1/1/1
1	HYP	А	17	1	-	0/0/11/13	0/1/1/1
1	HYP	С	80	1	-	0/0/11/13	0/1/1/1
1	HYP	А	2	1	-	0/0/11/13	0/1/1/1
1	HYP	С	62	1	-	0/0/11/13	0/1/1/1
1	HYP	В	50	1	-	0/0/11/13	0/1/1/1
1	HYP	В	38	1	-	0/0/11/13	0/1/1/1
1	HYP	С	68	1	-	0/0/11/13	0/1/1/1
1	HYP	А	26	1	-	0/0/11/13	0/1/1/1
1	HYP	А	20	1	-	0/0/11/13	0/1/1/1
1	HYP	В	59	1	-	0/0/11/13	0/1/1/1
1	HYP	С	74	1	-	0/0/11/13	0/1/1/1
1	HYP	А	14	1	-	0/0/11/13	0/1/1/1
1	HYP	А	8	1	-	0/0/11/13	0/1/1/1
1	HYP	В	47	1	-	0/0/11/13	0/1/1/1
1	HYP	С	71	1	-	0/0/11/13	0/1/1/1
1	HYP	С	77	1	-	0/0/11/13	0/1/1/1
1	HYP	В	35	1	-	0/0/11/13	0/1/1/1
1	HYP	А	23	1	-	0/0/11/13	0/1/1/1
1	HYP	В	56	1	-	0/0/11/13	0/1/1/1
1	HYP	В	41	1	-	0/0/11/13	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
1	В	44	HYP	CG-CB-CA	2.54	107.17	103.96

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

9 monomers are involved in 13 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	С	89	HYP	3	0
1	В	32	HYP	3	0
1	С	65	HYP	1	0
1	С	62	HYP	2	0
1	В	50	HYP	1	0
1	В	59	HYP	1	0
1	А	14	HYP	1	0
1	С	71	HYP	1	0
1	С	77	HYP	1	0

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Tune	Chain	Res	Link	B	Bond lengths			Bond angles		
	Type	Chain	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
2	ACY	С	305	-	$1,\!3,\!3$	0.44	0	$_{0,3,3}$	0.00	-	
2	ACY	А	306	-	$1,\!3,\!3$	1.06	0	$_{0,3,3}$	0.00	-	
2	ACY	С	301	-	$1,\!3,\!3$	1.39	0	$_{0,3,3}$	0.00	-	
2	ACY	В	302	-	$1,\!3,\!3$	0.83	0	$_{0,3,3}$	0.00	-	
2	ACY	А	307	-	$1,\!3,\!3$	1.84	0	$_{0,3,3}$	0.00	-	
2	ACY	А	304	-	$1,\!3,\!3$	0.15	0	$_{0,3,3}$	0.00	-	
2	ACY	А	303	-	$1,\!3,\!3$	1.50	0	$_{0,3,3}$	0.00	_	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.



1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	С	301	ACY	1	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

